arXiv:hep-ph/0109255v2 29 Nov 2001

Abstract

Title of Dissertation: A STUDY OF
NUCLEON SPIN STRUCTURE
FROM QUANTUM CHROMODYNAMICS

Jonathan Andrew Osborne, Doctor of Philosophy, 2000

Dissertation directed by: Associate Professor Xiangdong Ji
Department of Physics

Quantum Chromodynamics (QCD) is thought to be the correct microscopic theory of the strong force. Composed only of eight colored gluons and thirty-six colored quarks, its dynamics are believed to generate the observed hadronic spectrum of more than a thousand particles and account for many of their interactions - from the formation of nuclei to the rate at which a ρ-meson decays. Unfortunately, its predictions are often beyond the reach of of our current understanding of quantum field theory. Although the theory is perturbative at high energy, the phenomenon of confinement makes direct QCD calculations of physical observables quite intractable.

Through a process called factorization, one systematically separates the soft, incalculable physics from the perturbative high-energy physics. The soft physics is then grouped into matrix elements which can be measured in experiment. Once fixed, these matrix elements can be used to predict the outcome of other experiments. This process allows one to indirectly test the validity of QCD in the perturbative regime. Moreover, the matrix elements themselves are of interest. Acting as a window to the nonperturbative physics of QCD, these objects allow us to link the quark and gluon degrees of freedom of the microscopic theory to macroscopic observables such as hadron momentum and spin.

This thesis presents a study of how these ideas can be used to understand the origins of nucleon spin. Beginning with a general overview of QCD as a quantum field theory in Chapter 1, we then move on to study inclusive Deeply-Inelastic Scattering (DIS) in Chapter 2. Inclusive DIS presents one of the earliest examples of a factorizable process. The soft physics relevant to this process is related to parton probability distributions, and provided some of the first evidence that protons have nontrivial internal structure. We discuss the one-loop corrections to this amplitude, the definitions and interpretations of the distribution functions, and a proof that the hard and soft physics can be consistently separated at high energy to all orders in the QCD coupling.

The original research presented here concerns Deeply-Virtual Compton Scattering (DVCS), a process closely related to DIS. The special kinematics of DVCS allow one to measure new hadronic parton distributions. As explained in Chapter 3, these distributions contain information on the origins of nucleon spin that DIS cannot provide. We calculate the one-loop coefficient functions of these new distributions as they appear in DVCS and show that the soft physics inherent to this process can be separated from the hard at this order. Armed with the methods of Chapter 2, we then argue that this property persists to all orders in the strong coupling. Using our result, we calculate an operator product expansion for the product of two electromagnetic currents that is valid for a certain class of off-diagonal matrix elements. This constitutes a generalization of the expansion first considered by Wilson in 1969, which has been used extensively over the past thirty years.

Chapter 4 is devoted to the complications associated with transversely polarized DIS. We discuss the special circumstances that arise here due to the introduction of a transverse direction. In particular, the gauge-invariance of QCD demands that these effects be inseparable from multiparton correlations within the nucleon. Among other things, this fact complicates the scale evolution of the relevant distributions. We show that these complications are removed at the one-loop level by taking the limit of large N_c. Although this simplification has been known for several years, its origins have remained a mystery. Our new approach to the calculation allows us to directly understand these origins. We discuss the possibility of similar simplifications at the higher loop level and for other important distributions.

A STUDY OF NUCLEON SPIN STRUCTURE FROM QUANTUM CHROMODYNAMICS

by

Jonathan Andrew Osborne

Dissertation submitted to the Faculty of the Graduate School
of The University of Maryland in partial fulfillment
of the requirements for the degree of
Doctor of Philosophy
2000

Advisory Committee:

Associate Professor Xiangdong Ji, Chair
Professor Manoj Banerjee
Associate Professor Elizabeth Beise
Professor Thomas Cohen
Professor William Walters

© Copyright by

Jonathan Andrew Osborne

2000

Acknowledgements

I will always be grateful to my relatives and friends for supporting me through difficult times and understanding my need to discuss physics at the most inappropriate times. My cousin, Tom, who was always there with a beer when I needed a break. My father, who always pushed me to do my best. My mother, who always made me feel I could do it.

Over the past eight years, I have benefitted greatly from discussions with fellow students. It is only through heated debate that one completely understands the true nature of physics, and only through teaching and explaining that one can fully assimilate new material. Thanks in particular to Michael Hartl, Eduardo Ponton, William Linch, and C. K. Chow for many useful discussions on the subject matter of this dissertation (as well as many long, enjoyable digressions).

Over the years, I have had the great pleasure of working with and studying under several exceptional physicists and mathematicians. Their guidance has helped me to understand physics in a simple way which is difficult to present in textbooks. My undergraduate advisor, Prof. Ivan Kramer, once told me to keep on pushing and never give up. I have remembered that sentiment more than a few times during the last few months during the preparation of this manuscript. Prof. Thomas Cohen,

who always has a simple explanation for complicated mathematical results, has helped me to see physics from another perspective.

Of course, the largest contributions to this work come from my advisor, mentor, and collaborator Xiangdong Ji. His enthusiasm for the subject and confidence in his insights have always motivated me to prove him wrong - which is (almost!) impossible. At the same time, his confidence in *my* insights and understanding drives me to work harder to prove him right - which is nearly as difficult. I have been told many times that one can hear both of our laughs from the other side of the building. His influence has touched nearly every part of this dissertation, from finding irreducible representations of $SO(3,1)$ to proving all-orders factorization for DVCS. Without his patient understanding of my many faults, this dissertation would not have been possible.

In this revised manuscipt, I must also pay homage to Loretta Robinette, the beloved matriarch of the Theoretical Quarks, Hadrons, and Nuclei group at the University of Maryland. Time and again, she has done far more than I expected to help me with the preparation of this corrected version.

Table of Contents

0.1 Introduction 1

1 The Basics of QCD **20**

1.1 The Lagrangian 20

1.2 The Road to Quantization 23

1.3 Feynman Rules and Perturbation Theory 35

1.4 Renormalization in Covariant Gauges 38

1.5 The Beta Function - Asymptotic Freedom 60

1.6 Renormalization in the Axial Gauge -
The Price of a Ghost-Free Theory 66

2 Deep-Inelastic Scattering **81**

2.1 Kinematics 83

2.2 Deep-Inelastic Scattering on Partonic Targets 88

2.3 Infrared Divergences and Parton Distributions 102

2.4 Factorization to All Orders in DIS 126

2.5 The Operator Product Expansion 134

2.6 Summary of DIS 148

3 Deeply Virtual Compton Scattering **150**

3.1 Motivation 152

3.2 Kinematics and Parton distributions 160

3.3 One-Loop Compton Amplitudes
on Quark and Gluon Targets 165

3.4 One-Loop Factorization in DVCS 170

3.5 All Order Factorization of DVCS . . . 173
3.6 Generalized Operator Product Expansion and Wilson's Coefficients to the NLO Order . . . 176
3.7 Summary of DVCS . . . 180

4 The Structure Function $f_T(x,\mu^2)$ - Higher Twist Evolution 182
4.1 DIS With Transverse Polarization . . . 183
4.2 The Evolution Equation . . . 189
4.3 Conclusions . . . 196

A Standard Conventions and Formulae 198
A.1 Dirac Algebra . . . 200
A.2 $SU(N)$ Algebra . . . 202
A.3 Covariant integration . . . 205
A.4 QCD and Feynman's Rules of Perturbation Theory . . . 209
A.5 Other Useful Relations . . . 213
A.6 General References . . . 216

B The Lorentz Group and its Representations 218
B.1 Relativity . . . 219
B.2 Unitary Representations of $SO(3,1)$ . . . 227
B.3 The Spinor Representation of $SO(3,1)$ . . . 235

C $SU(N)$ 251
C.1 General Properties . . . 251
C.2 Local $SU(N)$ . . . 261
C.3 The Limit of Large N_c . . . 264

D Quantum Mechanics 267
D.1 Canonical Quantization . . . 267
D.2 Spacetime Symmetries - The Generators of the Poincaré Group . . . 275

E Dimensional Regularization **287**

List of Figures

Number **Page**

1 Gell-Mann's Eightfold Way for (a) spin-0 and (b) spin-1 mesons. These mesons form the **8** of $\mathbf{3} \times \overline{\mathbf{3}}$. If $SU(3)$ were exact, each multiplet would be degenerate and would not mix with the **1** of $\mathbf{3} \times \overline{\mathbf{3}}$. However, since in reality $SU(3)$ is badly broken, the mathematical ideals η_8 and ω_8 mix with the singlets η_0 and ω_0 to form the physical states η, η', ω, and ϕ. (c) and (d) show the baryon spin-1/2 octet and spin-3/2 decouplet, respectively. In each octet there are two particles in the center : one with isospin-0 and one with isospin-1. The fact that they lie on top of each other is due exclusively to the fact that I have chosen to suppress I^2. 8

1.1 The fundamental quark-gluon interaction is represented graphically by a Feynman diagram. The factor associated with the emission of a gluon with polarization μ and color index a is $-igt^a\gamma^\mu$, along with a δ-function to ensure momentum conservation. 36

1.2 This diagram presents a prime example of the reason symmetry factors are sometimes necessary to avoid overcounting. The Feynman rule for the triple gluon vertex counts each of these contributions as separate. In reality, however, only (a) and (b) represent different processes. In order to obtain the correct result using Feynman rules, one must include a factor of 1/2. 37

1.3 Three different kinds of contributions to the quark self-energy. The gluon loop in (a) is uncoupled to the sources and so contributes only to the vacuum energy. The reducible graph in (b) does not contain any new information; its contributions can be re-summed with a knowledge of irreducible diagrams, like that in (c). 40
1.4 The one-loop correction to the quark self-energy. 43
1.5 The one-loop correction to the ghost self-energy. 53
1.6 The three contributions to the gluon field-strength renormalization at one-loop order in covariant QCD. 55
1.7 The two vertex correction diagrams at one-loop order in QCD. 59

2.1 The two leading-order diagrams for virtual photon-quark scattering. The crossing symmetry allows us to obtain one from the other. 92
2.2 Contributions to virtual photon-quark scattering at (a) leading and (b)-(e) next-to-leading order in the QCD coupling. 97
2.3 Leading diagrams for gluon-virtual photon scattering. 101
2.4 Feynman rules for eikonal line propagation. 107
2.5 Contributions to the 'quark distribution in a quark' at (a) leading and (b)-(d) next-to-leading order in QCD. The final state in (a) and (b) is the vacuum, while (c) and (d) represent gluonic external states. 108
2.6 A representation of the all-orders scattering process. The nucleon 'blob' is intended to represent an all-orders sum, and the external photons are connected in all possible ways to this blob. 127
2.7 The most general reduced diagram for DIS. Straight lines represent collinear (or soft) partons of *any* species. 131
2.8 A graphical statement of factorization in the axial gauge. 133
2.9 A graphical illustration of the use of Ward identities to show that scalar gluons generate the gauge link in our parton distribution functions. Dashed lines represent the propagation of a scalar gluon. 135

3.1 Diagrams for Compton scattering on an onshell quark to order α_s. 166

3.2 The momentum flow in the box diagram 166
3.3 Diagrams for gluon Compton scattering at one-loop. 168
3.4 General reduced diagrams for the DVCS process. 175

4.1 Leading diagrams for 'naive' scattering with transverse polarization. 185
4.2 The gluon correction required for QCD gauge-invariance in DIS with transverse polarization. 186
4.3 These one-particle reducible diagrams represent the true nature of figure 4.1. They are obtained by eliminating the 'bad' component of ψ via the light-cone equations of motion. 189
4.4 The (a) two- and (b) three-point 1PI Feynman diagrams contributing to the evolution of θ_n in the large-N_c limit. 192
4.5 Four-point 1PI Feynman diagrams contributing to the evolution of $\hat{O}$ in the large-N_c limit. 194
4.6 Some two-loop 1PI Feynman diagrams that might break the autonomy of the θ_n in the large-N_c limit. 195

A.1 Feynman's rules for a non-abelian gauge theory with covariant gauge-fixing. The straight line represents quark propagation, while the squiggly line represents gluons and the dashed ghosts. In order to avoid index pollution, most of the indices here have been suppressed. The proper pole-prescription is given by adding $i\varepsilon$ to each denominator. Along with each vertex, a δ-function ensuring energy-momentum conservation is implied. 210

C.1 An illustration of the large-N_c counting rules. (a) is enhanced by a power of N_c^2 because of the unconstrained color loops within the bubble. Since it is a two-loop graph, it will contribute to the leading large-N_c behavior. On the other hand, (b) contains only one unconstrained color loop. As a three-loop diagram, its contribution comes only at $\mathcal{O}(1/N_c^2)$. Notice that (a) is a planar diagram, while (b) must contain a self-intersection when forced into a plane. As it stands, the counting rules for more complicated correlation functions, like that represented in (c), are ambiguous. At first, we may think that it is planar. However, drawing all external lines *directly* to the open dot (representing infinity), reveals its true nature. 265

A STUDY OF
NUCLEON SPIN STRUCTURE
FROM QUANTUM CHROMODYNAMICS

Jonathan Andrew Osborne

February 1, 2008

This comment page is not part of the dissertation.

Typeset by LaTeX using the `dissertation` style by Pablo A. Straub, University of Maryland.

0.1 Introduction

In the beginning, God created the Heavens and the Earth. After that, it has been the goal of physicists to understand the laws they obey. The dream of physics is a single set of laws which describe all phenomena from the largest galaxies to the smallest atoms and beyond. Recently, physics has come to a plateau of understanding in terms of which there are only a few physical processes which cannot be explained quantitatively. The physics of the very large is governed by Einstein's General Relativity [1]. This theory has had tremendous success in accurately predicting the orbits of the planets in our solar system, the bending of light by our Sun, and the behavior of radiation emitted by distant galaxies. The physics of the very small is thought to be described by the Standard Model. First formulated in its present form in 1973[1], this model is well-known for its extraordinarily accurate predictions of the properties of many subatomic particles, the excitation spectra of atoms, and the existence and theory of radioactive decay. Perhaps the most famous example of a precision calculation within the Standard Model is the calculation of the anomalous magnetic moment of the electron to 11 decimal places [5]. This quantity was measured to 7 parts in a trillion [6] as a test of the Standard Model, making it the most precisely measured quantity in the history of the human race. Agreement between experiment and theory to this degree of precision is unheard of in any other branch of physics, making this aspect of the Standard Model the most precisely tested theory in the history of physics.

The Standard Model (SM) consists of three parts. These can be thought of as Electricity and Magnetism, the Weak Interactions, and the Strong Interactions. Electricity and Magnetism in the SM is a relativistic quantum version of the familiar classical Electrodynamics embodied by the Maxwell Equations and Lorentz force law. This is the sector of the SM which is largely responsible for all of the macroscopic objects we see around us : computers, light bulbs, cars, etc. It is also responsible for almost every branch of science. It is the driving force behind chemistry, metallurgy, condensed matter physics, biology, etc. Well understood as a classical theory, its marriage with quantum mechanics led to the first realistic quantum

[1]The 'electroweak' sector was formulated in pieces throughout the 1960's. The most notable contributions are from S. L. Glashow [2], S. Weinberg [3], and A. Salam [4]. The 'strong' sector was not written in its present form until 1973, as discussed below.

field theory. Quantum field theories have dramatic implications on the structure of matter in general, some of which were studied for the first time in the context of quantum electrodynamics (QED). In particular, QED employs photons not only as the physical quanta of light, but also as carriers of the electromagnetic force. In QED, charged particles interact by exchanging massless spin-1 photons. This idea of forces being *mediated* by vector bosons[2] plays a fundamental role in the other two sectors of the SM. Indeed, one can understand many aspects of these elusive forces simply by exploiting an analogy to QED.

The Weak Interactions manifest themselves mainly in certain forms of radioactive decay. The processes of beta decay, electron capture, and positron emission first studied in the late 1890's [3] all take place via the Weak Interaction. Their discovery and subsequent study provided the first evidence of the existence of this sector of the SM. However, it was not understood that these processes required the introduction of a new force until the discovery of the neutron in 1932. Before this monumentous discovery, it was believed that beta decay was simply the emission of one of the 'nuclear electrons' needed to get the observed nuclear charge and mass right in the contemporary model of the nucleus. The discovery of the neutron forced scientists to reconsider the process that leads to beta decay in the nucleus. More recently, many other manifestations of this interaction have been seen experimentally, most notably the decay of muons and pions and the existence of the so-called 'strange' particles. However, this force has so many intricate features and is so well-hidden in ordinary matter that it was not completed for more than twenty years after these discoveries. Its completion at the end of the 1960's won Glashow [2], Weinberg [3], and Salam [4] the Nobel Prize in Physics in 1979. G. 't Hooft and M. Velman's 1972 proof that the theory is internally consistent and complete [10] won a second Nobel Prize in Physics in 1999. The experimental discovery by C. Rubbia and S. van der Meer of the W- and Z-bosons [11] proposed to mediate the weak interactions won them

[2]To highlight the difference between two basic particle types, particles which obey Bose-Einstein statistics are called bosons and those which obey Fermi-Dirac statistics are called fermions. This behavior finds its origins in the particle's *spin*, or intrinsic angular momentum. Particles with integer spin are invariably bosons and those with half-integer spin are fermions.

[3]True matter radiation was first discovered by H. Becquerel [7] in 1896. Originally discovered in samples of uranium ore, his 'uranic rays' were not understood as properties of individual atoms until M. Curie [8]. For a comprehensive history of the discovery of radiation, see [9].

the Nobel Prize in 1984.

In this work, we will mainly be concerned with the final sector of the SM. The Strong Interactions are responsible for the structure of nuclei and some of their decay modes. Alpha decay and the rare decays of spontaneous fission and proton and neutron emission are mediated by this force. It was alpha decay that first permitted the discovery of the nucleus itself. In 1909, H. Geiger and E. Marsden performed an experiment in which they found that approximately one in eight thousand alpha particles scattered off of a gold foil are scattered at an angle greater than 90° [12]. The contemporary model of the atom, due to J. J. Thompson [13], involved many negatively charged electrons embedded in a jelly-like medium of smeared-out positive charge. Since electrons are many thousands of times lighter than alpha particles, this model did not provide any mechanism for the observed back-scatter.

E. Rutherford proposed that atoms are composed of a very heavy positively charged hard core, or nucleus, surrounded by Thompson's negatively charged electrons. This core was estimated to be approximately one hundred thousand times smaller than the atom [14]. He argued that the electric field of this heavy nucleus was responsible for the back-scattering observed in recent experiments, rather than the light electrons or the diffuse jelly. This had the unattractive consequence that the vast majority of the volume of an atom is empty space. However, future experiments proved the correctness of this model. Since the attraction of electrons to the nucleus is accomplished by the electromagnetic field, it seemed the only thing left to study was the nucleus itself.

Radioactivity became physicists' 'window on the nucleus' when it was discovered that almost all radioactive processes are nuclear in origin. Physicists were beginning to understand that the nucleus itself is a strongly interacting bound state. Rutherford's experiments in 1919 showed that one could obtain oxygen by bombarding nitrogen with alpha particles [15]. Through years of these experiments and intense theoretical work, it became clear that nuclei are composed of two heavy particles, collectively called nucleons.

The positively charged protons responsible for the electric charge of nuclei were discovered first partly because they exist independently of other nucleons. These objects are the nuclei of the lightest isotope of hydrogen and were called 'positive electrons' by Rutherford until it became clear that they are really quite different from electrons. Neutrons were the last

constituent of 'ordinary' matter to be discovered since they do not carry an electric charge and decay into protons when isolated from nuclei. Long thought to be an extremely tightly bound system of a proton and an electron, neutrons were not given separate identities until 1932. In that year, J. Chadwick showed that the neutral radiation resulting from the bombardment of beryllium with alpha particles was not simply electromagnetic radiation, as widely believed, but was in fact a new kind of particle which he called the neutron [16]. He received the Nobel Prize for this discovery in 1935. The neutron solved many of the paradoxes the new quantum theory brought to nuclear physics. At last, the nucleus was seen as a sensible quantum system. All that was needed for a complete theory was an interaction responsible for the formation and stability of the bound states observed.

Once it was understood that the nucleus is made of positively charged protons and neutral neutrons, it became apparent that the electromagnetic repulsion inherent in nuclei must be overcome by a tremendously strong force. Called the Strong Force for obvious reasons, this new interaction was proposed to involve only protons and neutrons. Neutron, proton, and alpha[4] scattering data on various nuclei gave scientists of the time a good amount of information on the excitation spectra, but contemporary models were mainly empirical in origin. A modest degree of success was obtained by these early models, most notably the composite nucleus model of N. Bohr [17] and the celebrated nuclear shell model [18], but there was much left to be desired.

A partial solution to the problem of the origins of the nuclear force came with the discovery of a new triplet of particles and the exploitation of an approximate symmetry seen in nuclear interactions. It was proposed by H. Yukawa in 1934 [19] that if this new force is indeed fundamental, a description based on the newly formulated theory of relativistic quantum electrodynamics may lead to new insights. He argued that since the long-range nature of the electromagnetic field is attributed to the masslessness of the photon, a short-range nuclear force may be generated by massive particle exchange. Obtaining the range of the strong force from scattering experiments, he found that the mass of this new particle, called a meson, should be about two hundred times the mass of the electron.

In reality, there are five particles in Yukawa's predicted mass range. Only three of them

[4] α-particles were long known to be ^{4}He nuclei.

have the right properties to be candidates for the meson required in his theory. The other two appear as decay products of *these* three,[5] and have nothing whatsoever to do with the strong nuclear force.[6] This created a great deal of confusion in the scientific community for about ten years, but in 1947 C. F. Powell and collaborators were finally able to show conclusive evidence testifying to the existence of two charged strongly interacting bosons with masses approximately 270 times the electron mass [20]. For his prediction of their existence, Yukawa received the Nobel Prize in 1949. A third meson in this mass range was discovered in 1948, this one with no electric charge. The existence of this triplet forced physicists to take another look at the structure of the strong interactions.

For many years, it had been a known fact that to a very good approximation the strong force does not distinguish between protons and neutrons. In fact, if we take Yukawa's meson exchange force seriously, exchanges of charged mesons actually *change* neutrons *into* protons and vice versa. This new symmetry led physicists to consider protons and neutrons as different manifestations of the *same particle*. In these models, protons and neutrons are considered as components of a two-dimensional vector in a new space. This internal space is called 'isospin space' since it is mathematically identical to ordinary quantum-mechanical spin. The mesons are considered a vector in isospin space and work to rotate the nucleon states. This symmetry is actually a very good approximation to the true form of strong interaction physics, and had been useful in classifying particles and relating certain strong processes to each other. It also became instrumental in revealing to us the true nature of the strong interaction.

Along with the 1950's came a new generation of accelerators in which one could liberate larger and larger amounts of energy, which nature used to form exotic new particles. At this point, it was expected that all of the fundamental particles were known and these higher energies would only reveal more detail about the structure of the strong interactions and the behavior of nuclear excited states. However, hundreds of new strongly interacting particles had been discovered by 1960. These new objects were classified according to their charge, spin, decay properties, and isospin in an effort to make some sense of this 'zoo' of particles. The

[5] well, two of them...

[6] These objects, the $\mu^{\pm}$, are actually heavier copies of electrons. They provide the first evidence of the second generation of fermions.

situation here is analogous to Mendeleev's arrangement of elements on his periodic table. Just as his arrangement led to a greater understanding of chemistry and eventually the atom, this arrangement of particles led to the creation of the SM. In order to accommodate all these new particles, the word 'meson' was extended to mean 'integer spin strongly interacting particle' and the mesons of Yukawa's theory were called π-mesons, or pions. The word 'baryon' was introduced for strongly interacting fermions and 'hadron' as a blanket term for all strongly interacting particles.[7]

One of the most striking patterns in the particle zoo emerges when we arrange them according to the length of time it takes for them to decay. This exercise reveals three distinct time scales, showing physicists clear evidence of three different forces. The shortest decay times are on the order of 10^{-21} to 10^{-24} seconds, and involve only strongly interacting particles. These are attributed to the strong force. The mid-range lies on the order of 10^{-17} to 10^{-19} seconds and is associated with the electromagnetic interaction. These decays are signaled by the appearance of photons in the final state. The longest lived particles have half lives on the order of 10^{-10} to 10^{-8} seconds.[8] These decays are attributed to the weak interaction. Weak decays are signaled by the existence of leptons (like electrons and muons) in the final state, as well as missing energy attributed to the elusive neutrinos postulated to conserve energy and momentum in the weak decay of the neutron.

Many of the observed decays fit nicely into this scheme and physicists were given confidence in this three force model. However, a small number of mesons and baryons were found to behave very strangely. These hadrons participate in decays that look very much like strong decays, but they live entirely too long and certain symmetries seen to a very high degree of accuracy in all true strong interactions were badly broken by these decays. For example the heavy K-mesons, with a mass of about a thousand electrons, were found to have the competing

[7] Nuclei are not included in these designations. Although the ^{14}N nucleus is indeed a boson which participates in the strong interaction, it is never referred to as a meson. The terms 'meson' and 'baryon' will be defined more clearly below.

[8] Two notable exceptions are the neutron and the $\mu^{\pm}$, both of which have uncharacteristically long lifetimes due to phase space suppression.

decay modes

$$\begin{aligned} K^+ &\rightarrow \mu^+ + \nu_\mu \\ &\rightarrow \pi^+ + \pi^0 \ . \end{aligned} \tag{1}$$

The first mode is unambiguously a weak mode signaled by the neutrino ν_μ. The second appears to be a strong mode since all of the objects participating are strongly interacting particles. If this is the case, it should dominate the decay and cause the half-life of the K^+ to be on the order of the characteristic time scale of the strong interactions. In reality, the half-life of the K^+ is known to be $\sim 10^{-8}\, s$, characteristic of a weak decay.

A closer inspection of the second decay mode reveals that it does not respect the isospin symmetry found to such a good approximation in the strong interactions. This is easy to see since the K-mesons (kaons) have isospin-1/2 while pions have isospin-1. The final state can only have isospin-2, while the initial state has isospin-1/2. All of these strange new decays were found to violate isospin. Physicists of the time had to concede that we were seeing a new manifestation of the weak interaction. It was postulated that these particles posses a new quantum number (strangeness) which must be conserved by the strong interactions, but may be changed by one unit via the weak interaction. Mesons could be found only with strangeness 0 and ± 1, but there were baryons with strangeness ± 2 and ± 3 as well. A baryon with strangeness $S = 2$ has to decay first to a baryon with $S = 1$, then to a baryon with $S = 0$.

Since the strong interaction is postulated to conserve strangeness and (approximately) isospin, classification of its states in terms of these quantum numbers can give us insight into its true form. This observation led physicists to plot the particles of a given spin on axes labeled by these two quantum numbers.[9] Figure 0.1 shows this arrangement for the lowest-lying spin-0, -1/2, -1, and -3/2 hadrons. The mass-splitting within these multiplets is small compared with the energy required to jump to the next group of hadrons with the same spin, so to a first approximation these objects can be considered degenerate. This is much like the approximation of isospin symmetry which proved so fruitful in the past. In fact, isospin multiplets (particles lying along a horizontal line in Fig. 0.1) are very close to degenerate. One can think of each jump away from the $S = 0$ axis as a shift of about 350 electron masses,

[9]There are in fact two quantum numbers associated with isospin, as shown in Appendix C. In this plot, the isospin casimir I^2 has been suppressed and only the 3-component of isospin, I_3, is considered.




Figure 1: Gell-Mann's Eightfold Way for (a) spin-0 and (b) spin-1 mesons. These mesons form the **8** of $\mathbf{3} \times \overline{\mathbf{3}}$. If $SU(3)$ were exact, each multiplet would be degenerate and would not mix with the **1** of $\mathbf{3} \times \overline{\mathbf{3}}$. However, since in reality $SU(3)$ is badly broken, the mathematical ideals η_8 and ω_8 mix with the singlets η_0 and ω_0 to form the physical states η, η', ω, and ϕ. (c) and (d) show the baryon spin-1/2 octet and spin-3/2 decouplet, respectively. In each octet there are two particles in the center : one with isospin-0 and one with isospin-1. The fact that they lie on top of each other is due exclusively to the fact that I have chosen to suppress I^2.

while the splitting along the horizontal axis is closer to 20 m_e.[10] If the analogy with isospin were exact, we would assign the spin-1/2 baryons to the fundamental representation of some internal 8-dimensional space. However, this procedure would not allow us to understand the reason why isospin is a much better symmetry than this new group. Ideally we would like to place these particles in representations of some symmetry group which not only allows us to account for all our new low-lying states in a natural way, [11] but also contains our isospin symmetry as a subgroup. This means that we can break the full symmetry group in such a way that isospin remains unbroken.

Our ideal was achieved in 1962 by M. Gell-Mann's Eightfold Way [21]. Gell-Mann found that the observed hadron spectrum could be placed into representations of the special unitary group in three dimensions, $SU(3)$. Many of the mathematical properties of this group are discussed in Appendix C. Here, it is important only to know that there are two inequivalent fundamental representations of $SU(3)$ from which one can build all other representations. Calling these two representations $\mathbf{3}$ and $\overline{\mathbf{3}}$, we find that the mesons of spin-0 and -1 form representations of $\mathbf{3} \times \overline{\mathbf{3}}$ and the baryons of spin-1/2 and -3/2 form representations of $\mathbf{3} \times \mathbf{3} \times \mathbf{3}$. If we postulate the existence of objects in the fundamental representations of this $SU(3)$, as M. Gell-Mann and G. Zweig did in 1964 [22], say quarks for $\mathbf{3}$ and antiquarks for $\overline{\mathbf{3}}$, the last sentence says that mesons are mathematically equivalent to bound states of one quark and one antiquark and baryons are mathematically equivalent to bound states of three quarks. At this point, the idea of quarks and antiquarks is merely a mathematical construct. In order to suggest the actual existence of these particles we would have to propose a full theory of them explaining, in particular, why we haven't seen them in the lab.

Even if they don't actually exist in nature, quarks provide a nice way to organize the known particles. If we name the quarks in the fundamental representation 'up', 'down', and 'strange' (u, d, and s), we find that the new strangeness quantum number is simply the number of strange quarks in the hadron (with a minus sign from convention). Thus a baryon with $S = -2$ is composed of two strange quarks and one other quark. If we suppose that the strange quark

[10] In contrast, the jump to the next multiplet of spin-1/2 baryons is approximately 1000 m_e.

[11] This is not easy in a group with an 8-dimensional fundamental representation since the spin-3/2 baryons form a 10-dimensional representation.

is heavier than the other two, breaking the $SU(3)$ invariance of the theory, the multiplets break down exactly into the isospin multiplets we observe. The residual isospin symmetry is nothing but the remaining symmetry between up and down quarks. The electromagnetic charge assignments of all known particles can be predicted from the quark content if d and s quarks are given a charge of $-1/3$ and u quarks are given $+2/3$, in units of the proton charge. Conservation of strangeness is easily understood in this model if we postulate that the strong and electromagnetic interactions cannot change the type of quark, but the weak interaction can. In fact, almost all of these strangeness changing weak decays can be understood in terms of the basic weak decay of an s quark into a u quark and a negatively charged state (like a π^-). Other weak decays (like that of the neutron) are understood as a change from d to u or vice versa. This idea completely changed the shape of the weak interactions in the years to come. Spin assignments can be understood by postulating that quarks (and antiquarks) have spin-1/2: Since baryons are representations of $\mathbf{3} \times \mathbf{3} \times \mathbf{3}$, they may have spin-1/2 or -3/2.[12] Mesons will have integer spin since they are representations of $\mathbf{3} \times \overline{\mathbf{3}}$.[13] Hence we see that if quarks actually exist as particles, many of our problems will be solved. Much in the way that the discovery of protons, neutrons, and electrons reduced the number of fundamental particles from the eighty or ninety different atoms plus photons and other radiation to four, the discovery of quarks would reduce the hundreds of strongly interacting particles to three fundamental objects.

However nice Gell-Mann and Zweig's theory of quarks looks at first sight, it has some major problems. One problem is that no mechanism is given for binding the quarks into hadrons. Many models were proposed in analogy with QED to solve this problem, but no acceptable solution could be found. The new force would have to be extremely strong to account for the fact that hadrons appeared to be fundamental to the experimentalists of the time and no free quarks had been seen in the lab. In fact, no particle of fractional charge had ever been seen in any experiment anywhere. The fact that it was necessary to give quarks fractional charges to account for the observed hadron charges was greatly ridiculed at the time.

[12] Higher spins can also be obtained in this model by adding orbital contributions; these are always integer, so do not change the fact that the state will be a fermion.

[13] Herein lies the true meaning of the terms *baryon* and *meson*. A baryon is a three-quark bound state while a meson is a bound state of a quark and an antiquark.

Even more devastating is the problem of quantum statistics. Suppose, for example, we wish to construct the wavefunctions of the lowest-lying spin-3/2 baryons assuming the existence of quarks. Since we consider ground states, it is natural to assume that our wavefunctions are not orbitally excited (i.e. the orbital angular momentum of our states is zero). The only way to obtain spin-3/2 from three quarks without the help of orbital angular momentum is to align all of their spins, making this part of the wavefunction fully symmetric under quark exchange. Fermi-Dirac statistics then demands antisymmetry of the final $SU(3)$ sector. However, the **10** of $SU(3)$ is a fully symmetric representation. Hence the observed spectrum seems to require us to disallow the Pauli exclusion principle[14] [23] and be satisfied with a quantum state which is *symmetric* under the exchange of identical fermions.[15] In the non-relativistic theory, this principle had been made independently of the rest of quantum mechanics and could be argued away as something that is not valid in this arena. However, it had been shown by Dirac that a relativistic theory in which fermions do not obey the Pauli principle does not make sense. It was this argument that seemed like the devastating blow to the quark model. Quarks could not exist as real particles because they did not obey the basic requirements of a sensible quantum theory.

In reality, this blow was not as devastating as it seemed. In 1964 it was argued by O. W. Greenberg, M. Y. Han and Y. Nambu [24] that an extension of the simple quark model could solve this problem. If one endowed quarks with one other quantum direction in which they could be antisymmetrized, a sensible quantum theory could be constructed. There would have to be at least three directions in this new internal space since each quark in a baryon has to be associated with a different direction. Furthermore, since this new quantum number had not manifested itself in experiment, all observable objects would have to have zero charge. This

[14] Alternatively, we could concede that the spin-3/2 baryons are indeed orbitally excited. However, we would then be forced to explain why we see no lower-energy $SU(3)$ singlet state which is not orbitally excited.

[15] Even without taking the limit of exact $SU(3)$ symmetry, this problem exists. Consider the Ω baryon with its spin aligned along the z-axis. Experimentally, the Ω is seen to have $S = -3$; in the quark model, it is viewed as a bound state of three s quarks. In the absence of orbital excitations, all of our quarks must have their spins aligned along the z-axis to account for the observed spin-3/2 state. Hence we have three identical fermions in the same quantum state - an explicit violation of Pauli's exclusion principle without reference to $SU(3)$.

requires the number of internal directions to be exactly three, since otherwise three quarks could not form a singlet. Since the quarks must always be antisymmetrized in this internal direction the rest of their wavefunction will be symmetric, exactly as observed. Therefore, at the expense of introducing a new quantum number, we have disarmed the most devastating argument against the esthetically pleasing quark model. Of course, this postulate seems ludicrous at first glance. If the esthetics of the quark model were not so great, this theory may never have been advanced. However, the tremendous beauty of the quark model makes us want to interpret it physically rather than concede that it is nothing but mathematics. However, we cannot simply say that the quantum number is invisible and go home. It must have observable consequences. Furthermore, our theory still lacks the mechanism required to bind quarks inside hadrons.

These problems persisted for many years since no believable candidate for a theory of dynamical quarks appeared. At the time, people were still trying to make sense of quantum field theory itself, so there wasn't even a believable playing field on which to make a model of the strong interactions. In his book on current algebra, Gell-Mann suggested that quantum field theory be used only as a means to, "...construct a mathematical theory of the strongly interacting particles, which may or may not have anything to do with reality, find suitable algebraic relations that hold in that model, postulate their validity, and then throw away the model."[25]. This was the attitude of the time. Most of the activity in the field in the late 1960's was in the area of current algebra. Current algebra is a method advanced by Gell-Mann which allows one to derive experimentally verifiable relationships between observables from the symmetries underlying a theory without using the full formalism of quantum field theory. While the agreement between theory and experiment during this time was impressive, it was only Gell-Mann's $SU(3)$ that was being tested.

Real progress in the strong interactions continued in 1969 when E. D. Bloom et al. measured high energy electron-proton deep-inelastic scattering at SLAC for the first time [26]. Their data showed definitively that the proton behaved at high energy as though it was made of pointlike constituents. This prompted R. P. Feynman to advance his parton picture of deep-inelastic scattering (DIS) [27], in which he assumes that hadrons are made of pointlike *partons*.[16] This

[16]He is careful not to use 'quarks' in his model.

model has the advantage of describing DIS without assuming much about the nature of partons or their interactions. Further experiments at SLAC showed that the partons probed in DIS have spin-1/2. It seemed that a revival of the quark model was inevitable.

Another result from the SLAC data had an even more profound effect on the future of the strong interactions. At large angles, certain functions which parameterize the scattering were found to *scale* with energy. Rather than depending separately on the energy and momentum transfer to the proton, they depend only on a dimensionless ratio involving these two quantities. This behavior is predicted in the current algebra of Gell-Mann, as shown by C. G. Callan, D. J. Gross and J. D. Bjorken [28], and in the operator product expansion techniques of Wilson [29], which will be discussed in detail in Section 2.5. However, the details of renormalization in quantum field theories usually destroy this property. Intending to use the experimental results to finally put these nonsensical theories to rest, Gross attempted to prove that there exist no interacting quantum field theories which admit scaling. In a paper with Callan, he showed that in order for a field theory to scale it must have a property known as 'asymptotic freedom', that is it must behave like a free theory in the region of interest (the asymptotically large energy region in which scaling is predicted)[30]. Shortly thereafter, Gross and S. Coleman were able to show that the only kind of theory known which could be asymptotically free is a non-abelian gauge theory [31]. In order to complete his proof that quantum field theory cannot describe the strong interactions, Gross and his student F. Wilczek tackled the task of calculating directly whether or not these theories could admit asymptotic freedom. Much to their surprise, the calculation revealed that non-abelian gauge theories can indeed be asymptotically free [32].[17]

This result suggests that the symmetry associated with the quantum number introduced by Greenberg, Han and Nambu be used as the local symmetry required for a non-abelian quantum field theory of quark interactions. A physical motivation for the introduction of this new quantum number, called color because there are three primary colors, had been found in the total rate of $e^+e^- \to hadrons$, which was experimentally seen to be enhanced by a factor of three over the expectation from only three quarks. The rate of the electromagnetic decay $\pi^0 \to 2\gamma$ is similarly enhanced. This marks the first appearance of the theory of the strong interactions known today as quantum chromodynamics (QCD), or the quantum dynamics of

[17] H. D. Politzer also published this result simultaneously [33].

the color field.

At this time, it is advantageous for us to leave the historical treatment of the strong interactions and focus on QCD. Modern QCD views hadrons as highly relativistic bound states of nearly massless quarks and massless spin-1 objects known as gluons.[18] The strong interactions between hadrons are viewed as residual interactions, much like the molecular forces between neutral atoms. Obviously, this aspect of the modern theory of strong interactions complicates matters greatly - especially since we still do not fully understand the way it works in QED. Quarks interact by exchanging gluons in exactly the same way that electrons and protons interact by exchanging photons. However, in QCD gluons also exchange gluons. This new property comes from the fact that QCD is a non-abelian theory as opposed to QED, an abelian theory. This will be explained in detail in Chapter 1. For now, we explore some of the qualitative aspects of QCD.

The most important consequence of the non-abelian nature of QCD is the phenomenon of asymptotic freedom which led to its discovery. This property causes QCD to become a weakly interacting theory at short distances. Looking at this the other way, we see that QCD becomes strongly interacting at long distances. Since we don't know how to solve a strongly interacting quantum field theory, we cannot work out what this implies quantitatively. However, we can qualitatively say that if the interaction becomes stronger and stronger as color charges are separated, we can never completely liberate a color charge. Since a consistent quantum field theory has no choice but to allow the vacuum to create pairs of particles and antiparticles from ambient energy, the energy associated with separation of color charges will at some point become great enough to permit the creation of a pair of opposite color charges from the vacuum. At this point, the charges we were separating have happily combined with the new vacuum charges to form color singlets, and we find ourselves with two hadrons where we once had only one. This property of QCD is called color confinement : the color is physically confined in singlet arrangements. Of course, once the coupling gets large enough, the calculations required to show asymptotic freedom are no longer valid. It is for this reason that we cannot *prove* the existence of color confinement in QCD and it remains one of the most poorly understood aspects of the strong interactions, and, more generally, the entire SM.

[18] These are the analogue of QED's photons.

Quark	Constituent Mass(m_e)	Current Mass(m_e)
u	550-710	3-10
d	640-710	6-18
s	1000-1050	120-330

Table 1: Constituent versus current quark masses for the three lightest quarks.

One of the biggest mysteries of contemporary QCD is the approximate validity of naive quark models. Several phenomenological models of quarks were employed in the 1960's to get the observed hadron spectrum. If one assumes that a proton is made of two u quarks and a d quark, a Λ of a u, d, and an s, a K^+ of a u and an $\overline{s}$, et cetera, one can make naive predictions of the masses of the hadrons in terms of the quark masses. A spin-spin interaction proportional to the inverse masses of the quarks can account for the observed splitting between the spin-1/2 and spin-3/2 baryons and give us even more experimental data with which to compare. Once we have fit the few parameters of our phenomenological theory to data, we can predict the rest of the observed mass spectrum. These predictions are rather accurate, to the credit of our naive models. The fits also determine effective masses for the quarks.[19] In addition, one can obtain the masses of quarks in QCD from the current algebra approach mentioned above. These two methods give very different results, as shown in Table 1.

Another red flag seen in this approach is that it has completely ignored gluons. Gluons would certainly be expected to contribute to the energy of these highly relativistic bound states. Furthermore, the wavefunctions used to calculate the splitting between hadrons of different spin have assumed that the spin of hadrons comes only from quarks. However, since gluons are vector particles they are expected to contribute to the spin of the hadrons as well. In fact, asymptotically one expects approximately 50% of the spin of the proton to be carried by gluons. These models have also ignored the expected contribution from the so-called 'sea' quarks. It is well-known that relativistic quantum theories allow the vacuum to require create

[19]The best fits require one to use different quark masses for mesons and baryons. The values obtained from meson data are always smaller than those from baryons, and the fits are somewhat better in the baryon sector ($\sim 1\%$) than in the meson (as bad as $\sim 10\%$).

pairs of particles and antiparticles from the vacuum. It would certainly be naive of us to imagine that the highly relativistic strongly bound hadronic states consist only of three quarks or a quark and an antiquark. The simple fact that these systems are so strongly bound gives the system quite enough ambient energy to create many quark-antiquark pairs out of the vacuum. Such vacuum fluctuations are called the 'sea of quarks' in the hadron. Any realistic hadronic model must take this sea into account as well.

After all of this analysis, we are faced with the question of why these models work so well. It seems that somehow the interactions of QCD arrange the quarks, antiquarks, and gluons in such a way that a description of hadrons in terms of naive valence quarks is appropriate for certain quantities. These effective degrees of freedom are commonly called *constituent* quarks. The standard belief is that while these constituent quarks have the effective masses listed in the first column of Table 1, the true quarks of QCD have the much smaller 'current' masses in the second column. Of course, due to the effect of confinement, the masses of quarks and gluons are not well-defined within quantum field theory. Strictly speaking, the mass of an object is defined only as that object travels asymptotically far away from any other object. Since colored objects cannot do this, the 'mass' of a colored object has no meaning in this sense.[20]

One of the major difficulties with QCD is the fact that its degrees of freedom are not the same as those we see in the laboratory. This is an immediate consequence of confinement. If we would like to understand *any* strong interaction scattering process, we must first understand the complicated bound states involved in the process. This is an extremely difficult path to take; we still do not fully understand the scattering of complicated atoms. Nuclear bound states are even more difficult. Using QCD to understand even simple nuclei is undescribably complicated. For this reason, physicists have worked to find a theory which is expressed in terms of the low energy degrees of freedom. Since such a theory has composite objects as its fundamental fields, it cannot be expected to make sense for arbitrarily large momenta. However, if the momenta involved are small, the effects of the internal structure of the fundamental fields can be absorbed into unknown constants and fit to the data. Such theories are required to have the

[20] This is actually a property of any unbroken non-abelian gauge symmetry, whether or not it is asymptotically free.

same symmetries as the underlying theory, but are otherwise unconstrained. This naturally leads to a tremendously complicated theory whose lagrangian has infinitely many terms. The hope is that there is some small parameter in the theory which allows one to calculate only to finite order.

There are several theories which have achieved a limited degree of success in the past, all of which are derived from chiral symmetry. Chiral symmetry is an isospin-like symmetry possessed by QCD in the limit of small quark mass. An effective theory based on this symmetry ends up being an expansion in the ratio of external momentum or the pion mass to a scale which turns out to be near the proton mass. This means the expansion parameter in chiral perturbation theory is about 1/7. One interesting thing about Chiral Perturbation Theory (χPT) is that it restores the pions to their original place as carriers of the strong interaction. In χPT, one sees neutrons and protons exchanging pions in a way equivalent to that originally envisioned by Yukawa. However, this theory is much more complicated. The infinitely many terms in its lagrangian make power counting - the systematic truncation of the lagrangian - the most important game in town. In all expansions, one hopes that the next (uncalculated) coefficient is not unnaturally large. Even with a very small expansion parameter, this can lead to trouble. Unfortunately, many quantities which have been calculated in χPT are plagued with large coefficients. This fact of life has made low energy strong interaction physics very difficult. One wonders whether or not a sensible effective theory exists for this domain of nuclear physics.

Fortunately, the high energy side of QCD does not have these difficulties. Asymptotic freedom assures us that at high energy the QCD coupling is small and we may use perturbation theory as we do in QED. On the other hand, confinement tells us that we cannot escape the low energy physics we do not understand. Any physical process involving the strong interaction must necessarily also involve complicated bound states since they are the only physically acceptable external states of the theory. For example, DIS can be considered as the collision of a virtual photon and a proton. If we consider low energy scattering, the proton collectively absorbs the photon and recoils. At very high energy, the photon does not see the proton as a whole. Rather, it resolves the bound state into its constituents and is absorbed by one of the quarks. To properly calculate the amplitude for this process, we would have to

consider high energy scattering on an arbitrary parton in the proton and then average over the existing species of parton. High energy scattering on arbitrary partons can be done in QCD perturbation theory. However, the averaging process requires knowledge of the elusive proton wavefunction. Although we cannot calculate it, this object appears in many experiments. Since a proton is a proton regardless of what experiment we decide to subject it to, this information could in principle be used to predict the outcome of other experiments. More importantly, if we can use the perturbative calculation of the scattering to extract this information from experiment, we will have quantities which relate the physical degrees of freedom to those of our microscopic theory. This somewhat roundabout way of using perturbation theory to extract hadronic structural information from experiment is the topic of this dissertation.

In Chapter 1, I give a somewhat detailed introduction to the basics of perturbative quantum field theory as applied to QCD. This chapter is meant to provide a mathematical foundation for the rest of the thesis, as well as familiarize the reader with some of my conventions.

Chapter 2 is devoted to the aforementioned process of DIS. First considered in the late 1960's, this process provides the perfect first step for a look at factorization theorems, parton distribution functions and operator product expansions. It is here that the theoretical groundwork for my original research on deeply virtual Compton scattering (DVCS) is laid.

Chapter 3 encompasses the process of DVCS. Introduced by X. Ji because of the structural information it probes [34], DVCS can be used in conjunction with polarized DIS to extract the contribution to proton spin due to quark *orbital* angular momentum. This quantity cannot be isolated by any other experiment proposed in the literature. Using the treatment of Chapter 2 as a model, we calculate the one-loop QCD corrections to DVCS at the leading twist level and use the results to generalize Wilson's operator product expansion of two electromagnetic currents. More importantly, we show that this process is factorizable at this level to all orders in QCD perturbation theory. The material presented in this chapter was first published by X. Ji and I in [35].

Chapter 4 concerns the one-loop evolution of the twist-3 structure function $f_T(x)$ measured in DIS with transverse polarization. After a brief review of some of the complications that arise in transverse processes, we discuss the problem of its horrendous anomalous dimension matrix. The fact that this distribution mixes freely with the more general two-parton cor-

relators $G_1(x,y)$ and $G_2(x,y)$ causes immense complications in the analysis of experimental data. These other distributions are known to decouple in the limit of large N_c at the one-loop level [36], leaving only the simple evolution experienced by twist-2 distributions. This result is studied from a new standpoint in which the reason for the simplification becomes obvious. Our technique is extended to predict the outcome of the two-loop large-N_c result for twist-3 distributions and the one-loop twist-4. Unfortunately, this analysis reveals contributions which will surely destroy the simplification in these sectors. The material presented in this chapter is based on a paper published last year by X. Ji and I in [37].

Chapter 1

The Basics of QCD

In the introduction, I have outlined QCD qualitatively as a renormalizable quantum field theory describing the interactions of spin-1/2 quarks and spin-1 gluons. The form of this theory is governed by a gauge principle based on a local $SU(3)$ symmetry. This chapter is dedicated to explaining quantitatively what that means.

1.1 The Lagrangian

To build the lagrangian of QCD here, I will employ certain naturalness arguments which I believe to be a nice way to look at the SM. These arguments are certainly not intended as proofs that this is the only way the theory of strong interactions can be formulated. The validity of the theory is placed solely in the hands of experiment. However, it is interesting to note that one can build QCD in this way given only a knowledge of quantum field theory and an exact $SU(3)$ symmetry for some Dirac fields.

We begin with a theory of free spin-1/2 quarks. Assuming n_f different types (flavors) of colored quarks, we write our free lagrangian density as

$$\mathcal{L} = \sum_{f=1}^{n_f} \overline{\psi}_f (i \not{\partial} - m_f) \psi_f \ . \tag{1.1}$$

ψ_f denotes the color triplet of Dirac fields corresponding to quarks of flavor f. We note that quarks of different flavor are uncoupled. Our implicit assumption that color is entirely an internal symmetry is reflected in the fact that the quark mass does not depend on the color. The fact that this symmetry is internal means that there is no difference between quark

colors and our theory should be invariant under arbitrary reparameterizations of color. Since reparametrizations must not change the normalization of our fields, we restrict transformations to 3×3 unitary rotations, i.e. $\psi_f \to U\psi_f$ with U an arbitrary unitary 3×3 matrix. If the SM consisted only of this sector, these transformations could be made separately for each flavor. However, the weak sector breaks this larger symmetry group by coupling quarks of different flavors. In some sense, the weak sector forces us to choose the same color convention for all flavors; the remaining symmetry is concerned with which convention was chosen.

The group of unitary 3×3 matrices is called $U(3)$. Certain elements of $U(3)$ do not distinguish between different colors; they merely give an overall phase to the entire triplet. Such transformations are not associated with color reparametrization invariance[1], so we exclude them from our group. The remaining objects form the group of *special* unitary 3×3 matrices, $SU(3)$. Many of the mathematical properties of $SU(3)$ are discussed in Appendix C. At present we need only the form of a general[2] $SU(3)$ matrix,

$$U = e^{i\theta^a t^a} \ . \tag{1.2}$$

The t^a, a=1 to 8, are the $SU(3)$ generators introduced in the appendix. For now, their explicit form is unimportant. We need only know that they are hermitian and traceless. The θ^a are arbitrary real numbers which we will take to be small for most of our discussion.

When we rotate our quark fields, we do so in the same way at every point in spacetime. From the standpoint of special relativity, this seems like a strange procedure. We don't transmit any information during this process, so it does not violate causality. However, it still is not a natural thing to do. Ideally, we would like to define conventions separately at each point in spacetime so that the physics we describe is local. This means we wish our lagrangian to be invariant under arbitrary *spacetime dependent* $SU(3)$ transformations. Such local transformations are called gauge transformations.

Since a derivative compares field values at two infinitesimally close, but separate, spacetime

[1] These are are precisely the transformations which lead one to electromagnetic interactions.

[2] The colorblind $SU(3)$ transformations $e^{2i\pi/3}$ and $e^{4i\pi/3}$ cannot be written in this form. However, since these transformations are indeed colorblind, there are rotations in the other sectors which will compensate for them. For this reason, we may completely ignore these troublesome phases with the rest of the electroweak sector of the SM.

points without taking into account a difference in convention, our present lagrangian does not admit this local symmetry. However, if we decide that we really want this symmetry to be local, we can alter our theory slightly to admit it. The reason why the derivative term is not invariant under the local symmetry is that $\partial\psi$ does not transform in the same way as ψ. There is an extra additive term here which reflects the spacetime change in the transformation itself. If we add an extra term to the derivative to cancel this change in the transformation, our theory will be invariant. With this in mind, we introduce a covariant derivative $\mathcal{D} \equiv \partial + ig\mathcal{A}$.

Hermiticity of the lagrangian[3] requires $g\mathcal{A}$ hermitian. Since we want $\mathcal{D}\psi$ to transform in the same way as ψ itself, we require $\mathcal{D} \to U\mathcal{D}U^\dagger$. This implies that

$$\mathcal{A} \to U\mathcal{A}U^\dagger - \frac{i}{g}\, U\partial U^\dagger \ . \tag{1.3}$$

Any matrix $\mathcal{A}$ with a nonzero trace can be written as the sum of a traceless matrix and a multiple of the identity. As mentioned above, the identity is colorblind and contributes only to the electroweak sector of the SM. For this reason, we consider $\mathcal{A}$ traceless.[4] All traceless hermitian 3×3 matrices can be written as a linear combination of the eight $SU(3)$ generators, so $\mathcal{A}(x)$ contains eight hermitian vector fields - one for each generator of $SU(3)$. Objects which are related to the generators of a group in this way are said to be in the *adjoint* representation of the group.

We have promoted our $SU(3)$ color symmetry to a local symmetry at the cost of introducing new vector fields. Since these fields transform under the symmetry, it is natural to consider them as dynamical fields in our theory. In order to do this, we must endow them with the ability to propagate. In other words, we must give them a kinetic term. It is immediately obvious that the standard kinetic term, $1/2\, \partial_\mu \mathcal{A}_\nu \partial^\mu \mathcal{A}^\nu$, is not invariant under the local $SU(3)$ symmetry we have worked so hard to install. The nonlinear transformation (1.3) makes it difficult to construct an invariant kinetic term for $\mathcal{A}$. However, since $\mathcal{D}$ contains $\mathcal{A}$ and transforms linearly under the action of the gauge group, we can use it to form a covariant object which involves derivatives of $\mathcal{A}$:

$$\mathcal{F}^{\mu\nu} \equiv -\frac{i}{g}\,[\mathcal{D}^\mu, \mathcal{D}^\nu] \ . \tag{1.4}$$

[3] up to a total derivative

[4] This property is not altered by the gauge transformation (1.3), as can be seen by inspection.

The scalar $\mathcal{F}^{\mu\nu}\mathcal{F}_{\mu\nu}$ contains the appropriate terms for a vector boson kinetic energy. It's trace is also invariant under the gauge symmetry. Actually, this term and the quark piece above are the *only* gauge-invariant terms of dimension four or smaller[5] involving only these fields.[6] In particular, there can be no term that corresponds to a mass for the $\mathcal{A}$ fields.

Collecting these pieces and normalizing the kinetic term of our vector bosons, we write the full QCD lagrangian density as

$$\mathcal{L} = \sum_{f=1}^{n_f} \overline{\psi}_f (i\not{\mathcal{D}} - m_f)\psi_f - \frac{1}{2}\mathrm{Tr}\,\mathcal{F}^{\mu\nu}\mathcal{F}_{\mu\nu} \ . \tag{1.5}$$

This expression looks nearly identical to that of QED. In fact, the only difference is in the gauge group. Let us calculate $\mathcal{F}$:

$$\mathcal{F}^{\mu\nu} = \partial^\mu \mathcal{A}^\nu - \partial^\nu \mathcal{A}^\mu + ig[\mathcal{A}^\mu, \mathcal{A}^\nu] \ . \tag{1.6}$$

The first two terms are familiar from QED, while the last is not. Since we can express the gauge field $\mathcal{A}$ as a linear combination of generators of the gauge group, the last term expresses the commutator of two generators of $SU(3)$. For abelian groups, where all of the generators commute, this term is not present. This is trivially the case in QED, where there is only one generator. In non-abelian theories, like QCD, the inclusion of this term is unavoidable. We will see that it leads to self-interactions among gluons, the quanta of the $\mathcal{A}$ fields. It is these interactions which cause the theory to be asymptotically free, as shown in Section 1.4.

1.2 The Road to Quantization

Now that we have a lagrangian, we would like to quantize the theory. In scalar field theory, this is implemented merely by imposing commutation relations on the fields and their conjugate momenta. One introduces creation and annihilation operators[7] and proceeds with a straightforward evaluation of matrix elements. In QED, it is not quite that simple. Gauge symmetry

[5] This is required for renormalizability, as will be mentioned in Chapter 4.

[6] There is one other term which is thought to contribute to physics beyond perturbation theory. Its inclusion violates the discrete spacetime symmetries of parity and time-reversal, and its coefficient is experimentally seen to be extremely small.

[7] See Appendix D.1.

requires the introduction of unphysical degrees of freedom. This is obvious because a gauge transformation is by definition an unphysical thing. It corresponds merely to a redefinition of convention. For each configuration of the photon field, there exist infinitely many physically identical configurations which are obtained by transforming the original. Taking all of these into account separately is highly redundant and creates divergences which must be removed before quantization can make sense.[8] In order to make sense of a gauge theory, one must first choose a gauge so that every physically distinct field configuration is taken into account exactly once. The way this gauge is chosen is arbitrary, but certain choices make the resulting formalism more complicated than others.

To fix a gauge, one chooses a function of the gauge field and sets it equal to a certain function of spacetime. Normally, people choose a linear function to simplify the manipulations necessary to fix the gauge. A standard example is the axial gauge. This gauge choice is made by requiring $n \cdot \mathcal{A}(x) = \omega(x)$ for some function ω and some fixed vector n. If n^* satisfies $n^* \cdot n = 1$, then

$$\mathcal{A}(x) \to \mathcal{A}(x) + \partial \int^x (\omega(z) - n \cdot \mathcal{A}(z)) n^* \cdot dz \tag{1.7}$$

is an abelian gauge transformation which takes an arbitrary field configuration into one that satisfies the gauge condition. Hence this condition can be satisfied by all physically distinct configurations.[9] However, this gauge choice does not completely specify the gauge. If v satisfies $v \cdot n = 0$, the gauge transformation

$$\mathcal{A}(x) \to \mathcal{A}(x) + \partial f(v \cdot x) \tag{1.10}$$

does not change $n \cdot \mathcal{A}$. This is called a residual gauge transformation. If one wishes to completely specify the gauge, one must enforce a further gauge restriction to specify the residual gauge.

[8] This is purely an artifact of our mathematical incompetence. If we were smarter, we would not have to do such violence to gauge theories in order to make sense of them.

[9] In a non-abelian gauge theory, the required transformation is somewhat more involved. An explicit solution is

$$U = (\mathrm{P} e^{-ig \int^x \omega(z) n^* \cdot dz})(\overline{\mathrm{P}} e^{ig \int^x n \cdot \mathcal{A}(z) n^* \cdot dz}) \ , \tag{1.8}$$

where ($\overline{\mathrm{P}}$) P denotes (anti-)path ordering of the exponential. This ordering is constructed so that

$$n \cdot \partial U = ig(U n \cdot \mathcal{A} - \omega U) \ . \tag{1.9}$$

This is not usually necessary. In many cases, the ambiguities associated with residual gauge transformations are not enough to interfere with the quantization process. With the choice of axial gauge, the residual degrees of freedom exist in a special corner of momentum space. We will see that our gauge-fixed theory regards this corner as a singular region of momentum space, and special care must be taken near it.

In the axial gauge QCD and QED are very similar, but the problems associated with its residual gauge transformations can be quite severe. The axial gauge also breaks Lorentz invariance explicitly by introducing the special direction n during the quantization process. For these reasons, another gauge-fixing function is often employed. The 'covariant gauges' are obtained by specifying $\partial \cdot \mathcal{A}(x) = \omega(x)$. Once again, this gauge can be shown to accommodate all physical configurations since the abelian transformation

$$\mathcal{A}^\mu(x) \to \mathcal{A}^\mu(x) + \partial^\mu \int d^4z \, \frac{d^4q}{(2\pi)^4} e^{iq\cdot(x-z)} \frac{\partial_\nu \mathcal{A}^\nu(z) - \omega(z)}{q^2} \tag{1.11}$$

takes any configuration into one which satisfies our gauge condition.[10] This gauge choice also contains an ambiguity since any transformation of the form

$$\mathcal{A}^\mu(x) \to \mathcal{A}^\mu(x) + \partial^\mu \chi(x) \tag{1.12}$$

with $\chi(x)$ harmonic[11] leaves $\partial \cdot \mathcal{A}$ invariant. Immediately, one can see that the residual transformations associated with this gauge choice occupy a much smaller region of momentum space than those of the axial gauge. In the previous case, one has an infinite three-dimensional subspace of momentum to contend with. Here, only the light-cone itself is a singular region.[12] This region is the three-dimensional surface $(q^0)^2 = |\vec{q}|^2$, which is compact for each value of $|q^0|$. Furthermore, the divergences arising from this ambiguity in axial gauge require special treatment. Those appearing in covariant gauges can be treated along with other divergences which appear naturally in quantum field theory. We will see this in detail in the following sections.

Let us now proceed with the quantization process. To illustrate the similarities between

[10] Once again, this transformation is more involved for non-abelian groups. In this case, the explicit transformation is too complicated to present here.

[11] I.e. $\partial_\mu \partial^\mu \chi(x) = 0$

[12] The statement $\partial \cdot \partial \chi = 0$ reads $q^2 = 0$ in momentum space.

QED and QCD, we will not specify the gauge group immediately. In my view, the best way to see the process of gauge fixing lies in the path integral approach.[13] In this approach to quantum field theory, vacuum matrix elements of an operator are understood as weighted averages of the operator over all possible field configurations. The weight given to a particular configuration is the exponential of its action, $\mathcal{S}[\Phi] \equiv \int d^4x\, \mathcal{L}[\Phi]$:

$$\langle \mathrm{T}\mathcal{O}(x_1, \ldots, x_n)\rangle_{vac} = \frac{\int \mathcal{D}[\Phi]\mathcal{O}(x_1, \ldots, x_n)e^{i\mathcal{S}[\Phi]}}{\int \mathcal{D}[\Phi]e^{i\mathcal{S}[\Phi]}} \quad , \tag{1.13}$$

where Φ stands for all fields in the theory and $\mathrm{T}\mathcal{O}$ is any n-point correlation function of any of the fields Φ appropriately time-ordered.[14] The path integral takes *all* field configurations into account, which means that it is redundantly integrating over a continuous infinity of physically identical gauge configurations. The idea behind gauge fixing is to make this redundant integration explicit so it will cancel in the ratios appearing in physical matrix elements. Let us study only the denominator of Eq.(1.13). Our derivation will follow for the numerator as long as $\mathcal{O}$ is gauge-invariant. Parameterizing a general gauge transformation by real functions $\theta^a(x)$ as in Eq.(1.2), we write the identity

$$1 = \int \mathcal{D}\theta^a \delta(\mathcal{G}^a[\mathcal{A}^\theta] - \omega^a) \left| \det\left(\frac{\delta \mathcal{G}^a[\mathcal{A}^\theta]}{\delta \theta^b} \right) \right|_{\mathcal{A}} \quad . \tag{1.15}$$

This equation says that there exists a unique[15] gauge transformation parameterized by $\theta^a(x)$ which will transform a given arbitrary field configuration $\mathcal{A}$ into a new configuration $\mathcal{A}^\theta$ which satisfies the gauge conditions $\mathcal{G}^a[\mathcal{A}^\theta] = \omega^a$.[16] The determinant is the Jacobian of the change of variables between $\mathcal{G}$ and θ. Inserting this into the denominator of Eq.(1.13)

[13] For a derivation of the path integral and a discussion of its technical points and physical meaning, see Ref. [38]. Essentially, the path integral is an instruction to sum the contributions of every field configuration. This is done by integrating over the value of a field $\Phi(x)$ at *every* spacetime point x.

[14] One defines the *time ordering operator*, T, such that

$$\mathrm{T}\left\{\Phi(x_1)\Phi(x_2)\right\} \equiv \Phi(x_1)\Phi(x_2)\Theta(x_1^0 - x_2^0) \pm \Phi(x_2)\Phi(x_1)\Theta(x_2^0 - x_1^0) \quad . \tag{1.14}$$

The + refers to bosons while the − refers to fermions.

[15] I note here that neither of the gauge conditions specified above satisfy this property, as exemplified by Eqs.(1.10,1.12). What we really need is for this expression to be independent of $\mathcal{A}$ and ω^a. Both the axial gauge and the covariant gauge satisfy this requirement. The actual number of such gauge transformations will cancel in the calculation of any physical matrix element as long as it is the *same* for *all* configurations.

[16] Note that we have one gauge condition for each generator.

and performing the inverse gauge transformation parameterized by $-\theta^a$ on the integration variables,[17] we obtain

$$\int \mathcal{D}\theta^a \, \mathcal{D}[\Phi] \left| \det \left(\frac{\delta \mathcal{G}^a[\mathcal{A}^\theta]}{\delta \theta^b} \right) \right|_{\mathcal{A}^{-\theta}} \delta(\mathcal{G}^a[\mathcal{A}] - \omega^a) e^{i\mathcal{S}} \ . \tag{1.16}$$

It is easily seen by inspection that the dependence of the determinant on θ is entirely illusory, so the integration over θ factorizes completely and cancels in the ratio (1.13). This factor directly represents the redundancy that led to the need for gauge fixing in the first place.

Since the $\omega^a(x)$ were introduced as arbitrary functions, we may consider a weighted average over all possible ω^a :

$$\frac{1}{\int \mathcal{D}\omega^a \, f[\omega^a]} \int \mathcal{D}\omega^a \int \mathcal{D}[\Phi] f[\omega^a] \left| \det \left(\frac{\delta \mathcal{G}^a[\mathcal{A}^\theta]}{\delta \theta^b} \right) \right|_{\mathcal{A}^{-\theta}} \delta(\mathcal{G}^a[\mathcal{A}] - \omega^a) e^{i\mathcal{S}} \ . \tag{1.17}$$

The integration over ω cancels independently of the form of the functional $f[\omega^a(x)]$ if we use the fact that Eq.(1.16) does not actually depend on ω^a. On the other hand, we may consider the denominator an unimportant (though infinite) constant and use the δ-function to do the ω^a integral in the numerator :

$$\frac{1}{\int \mathcal{D}\omega^a \, f[\omega^a]} \int \mathcal{D}[\Phi] f[\mathcal{G}^a[\mathcal{A}]] \left| \det \left(\frac{\delta \mathcal{G}^a[\mathcal{A}^\theta]}{\delta \theta^b} \right) \right|_{\mathcal{A}^{-\theta}} e^{i\mathcal{S}} \ . \tag{1.18}$$

Note that although the constraint is no longer explicit, it is obvious from our construction that in some sense $\mathcal{G}^a[\mathcal{A}]$ *does not depend on* $\mathcal{A}$.[18]

All that is left is the calculation of the determinant. Since this depends nontrivially on the form of the gauge condition and the specific theory under consideration, we cannot go any further without specification.

Since QED is less complicated, we will consider it first. In an axial gauge,

$$\mathcal{G}[\mathcal{A}^\theta] = n \cdot \mathcal{A} + 1/g \, n \cdot \partial\theta \ . \tag{1.19}$$

Hence we are asked to evaluate the determinant of $1/g \, n \cdot \partial$. The meaning of this object is not obvious. If we really wanted to evaluate it, we would have to consider the actual definition of

[17] Since this transformation is unitary on the fermionic fields and the gauge fields (plus a trivial constant shift), the measure of the path integral is invariant. If it weren't, the path integral itself wouldn't be gauge invariant.

[18] This statement is true in the sense that, once all physically indistinguishable field configurations are taken into account, the contribution from each physically distinct set of configurations is identical.

the path integral. Such things are considered in [38]. Here, we do not care about this infinite constant. Since it is independent of all the relevant fields, it will cancel in the calculation of physical quantities (much like every other constant we have ignored in this derivation). Hence in this case the determinant may be completely ignored.

In a covariant gauge, the Jacobian, $\det(1/g\,\partial\cdot\partial)$, is also constant. This leaves the function $f[\omega]$ as the sole remnant of our gauge-fixing in QED. Since we must live with it, we intend to use our freedom to choose the form of $f[\omega]$ to the best of our advantage. This is accomplished by choosing a function which will add a quadratic term to the lagrangian. Any other choice would lead to new vertices for us to take into account in perturbation theory.[19] The function $f[\omega] = \exp[-i\int d^4x\,\omega(x)^2/2\xi]$ suits our purpose.

The only effect of all this work is the simple shift in the lagrangian density

$$\mathcal{L}^{axial}_{\rm QED} = \mathcal{L}^{0}_{\rm QED} - \frac{(n\cdot\mathcal{A})^2}{2\xi} \tag{1.20}$$

for the axial gauge and

$$\mathcal{L}^{cov}_{\rm QED} = \mathcal{L}^{0}_{\rm QED} - \frac{(\partial\cdot\mathcal{A})^2}{2\xi} \tag{1.21}$$

for the covariant gauge. There is one degree of freedom left from the choice of weighting function - the number ξ. In principle, this object could be a function of spacetime since the weighting was *completely* arbitrary. However, again the limitations of our mathematical abilities force us to be less general than the derivation requires. As it is, we take ξ as an arbitrary constant. This degree of freedom is useful as a check of lengthy calculations since its dependence must cancel in all gauge-invariant quantities.

We turn now to QCD. The similarities between QCD and QED are more apparent in the axial gauge, so we begin with this choice. Our first task is to rigorously define what is meant by the Jacobian under consideration. The matrix $\delta\mathcal{G}/\delta\theta$ is unambiguously given by

$$\mathcal{G}^a[\mathcal{A}^{\theta(x)+\delta\theta(x)}(x)] - \mathcal{G}^a[\mathcal{A}^{\theta(x)}(x)] = \int d^4y\,(\delta\mathcal{G}/\delta\theta)^{a\,b}(x,y)\delta\theta^b(y)\;. \tag{1.22}$$

As mentioned above, our Jacobian does not actually depend on θ. Hence we may take $\theta = 0$

[19] Actually, any other choice would lead to disaster in the sense that one still could not define a propagator. In this way, the choice of $f[\omega]$ is made for us by the way we know how to do quantum field theory. Of course, in principle the choice is arbitrary. However, we must know how to live with whatever choice we make.

to simplify our task. For infinitesimal $\delta\theta$, Eq.(1.3) reduces to

$$\mathcal{A} \to \mathcal{A} - \frac{1}{g}[\mathcal{D}, \delta\theta] \; . \tag{1.23}$$

The second term in this expression is the covariant derivative of $\delta\theta$:

$$\begin{aligned} [\mathcal{D}, \theta] &\equiv t^a \, \mathcal{D}^{a\,b} \, \theta^b \;\; ; \\ \mathcal{D}^{a\,b} &= \partial \, \delta^{a\,b} + g \, f^{a\,b\,c} \mathcal{A}^c \;\; , \end{aligned} \tag{1.24}$$

where $f^{a\,b\,c}$ are the structure constants of $SU(3)$ defined in Appendix C. Our matrix is given by

$$-\frac{1}{g} \, n \cdot \mathcal{D}_x^{a\,b} \delta\theta^b(x) = \int d^4y (\delta\mathcal{G}/\delta\theta)^{a\,b}(x,y) \delta\theta^b(y) \;\; , \tag{1.25}$$

so

$$(\delta\mathcal{G}/\delta\theta)^{a\,b}(x,y) = -\frac{1}{g} \, n \cdot \mathcal{D}_x^{a\,b} \delta^{(4)}(x-y) \; . \tag{1.26}$$

This matrix appears to depend on $\mathcal{A}$ through the covariant derivative. However, the dependence of the matrix on $\mathcal{A}$ is of the form $n \cdot \mathcal{A}$. Since we have *fixed* this component of $\mathcal{A}$, it no longer depends on the physical configuration under consideration. This allows us to ignore the determinant once again, making the quantization process in axial gauge identical for QED and QCD. The only difference between the two theories appears in the gluonic self-coupling terms in the kinetic energy.

The situation is not so simple in covariant gauge. The derivation of the matrix goes through as before. We find

$$(\delta\mathcal{G}/\delta\theta)^{a\,b}(x,y) = -\frac{1}{g} \, \partial \cdot \mathcal{D}_x^{a\,b} \delta^{(4)}(x-y) \; . \tag{1.27}$$

This time, the dependence on $\mathcal{A}$ is not illusory. Since ∂ is a differential operator that will act on the δ-function as well as the gauge field, the determinant of this matrix will contribute to the dynamics of our gauge-fixed theory. We must figure out some way to take it into account in our calculations if we want to use this gauge.

By direct analogy to N-dimensional integrals, one can perform Gaussian path integrals exactly. For any nonsingular hermitian matrix $\mathcal{M}$, we have

$$\int \mathcal{D}\phi^\dagger \mathcal{D}\phi e^{i \int d^4x \, d^4y \; \phi_i^\dagger(x) \mathcal{M}_{i\,j}(x,y) \phi(y)_j} = |\mathrm{det}\mathcal{M}|^{-1} \tag{1.28}$$

if the fields ϕ and $\phi^\dagger$ are bosonic (or commuting) fields and

$$\int \mathcal{D}\overline{\psi}\mathcal{D}\psi e^{i\int d^4x\, d^4y\, \overline{\psi}_i(x)\mathcal{M}_{i\,j}(x,y)\psi(y)_j} = |\mathrm{det}\mathcal{M}| \tag{1.29}$$

for fermionic (anticommuting) fields ψ, $\overline{\psi}$. Hence a simple way to take care of our problem is to utilize auxiliary anticommuting fields c and $\overline{c}$ to exponentiate the determinant. These fields are *not* physical. They are only mathematical artifacts we introduce to help us through the day. With this little snag out of the way, we write the QCD lagrangian with covariant gauge fixing as[20]

$$\mathcal{L}^{cov}_{\mathrm{QCD}} = \mathcal{L}^0_{\mathrm{QCD}} - \frac{(\partial\cdot\mathcal{A}^a)^2}{2\xi} - \overline{c}\,\partial\cdot\mathcal{D}\,c\,. \tag{1.30}$$

The new anticommuting 'ghosts' are very strange objects. Contrary to appearances, c and $\overline{c}$ are completely unrelated complex fields[21] which behave dynamically like scalars. Their sole function is to account for the Jacobian in Eq.(1.15). If it weren't for the gluon field appearing in the covariant derivative, they would not interact with any physical field and their fluctuations would simply shift the energy of the vacuum by a constant amount. These shifts would then cancel with the fluctuations of the denominator of Equation (1.13) for any properly normalized physical matrix element. However, their interaction with $\mathcal{A}$ allows an operator $\mathcal{O}$ to *polarize* these fluctuations so that they do not cancel its matrix elements. We will come back to the story of ghosts later when we consider the renormalization of the gluon fields.

At this point, it is convenient to introduce source terms into our action in order to define a generating functional for correlation functions :

$$\begin{aligned} \mathcal{S}[\Phi,\mathcal{J}] &\equiv \mathcal{S}[\Phi] + \int d^4x\,\mathcal{J}(x)\Phi(x) \\ \mathcal{Z}[\mathcal{J}] &\equiv \frac{\int\mathcal{D}[\Phi]e^{i\mathcal{S}[\Phi,\mathcal{J}]}}{\int\mathcal{D}[\Phi]e^{i\mathcal{S}[\Phi,0]}}\,. \end{aligned} \tag{1.31}$$

I have introduced a source for each field in the theory. Since taking functional derivatives with respect to the external source $\mathcal{J}$ and subsequently taking $\mathcal{J}=0$ can give us any correlation function,[22] the only object we need to consider is $\mathcal{Z}[\mathcal{J}]$. This is by no means a trivial task.

[20] The factors of i and g go into our overall normalization as before.

[21] The bar over $\overline{c}$ is merely convention. There is no γ^0 implied here.

[22] Of course, here we must worry about the interchange of functional integration and functional differentiation. While this interchange is extremely difficult if not impossible to justify, the theory it leads to is the same

The difficulties associated with QCD stem almost exclusively from an inability to calculate this object. A whole branch of QCD is dedicated to calculating this quantity numerically. Even in this approach there are many problems associated with discretization, finite lattice size, finite computing power, etc. Here, we will approach the problem in the same way it is approached for QED. Assuming the coupling g to be small, we solve the free theory and take interactions into account perturbatively. This method is certainly not without its shortcomings, some of which I will try to explain along the way.

Since we can solve the free theory using (1.28) and (1.29), it is advantageous for us to separate this part from the interaction part. We will do this explicitly for the covariant gauges since they are the most complicated. The construction is identical in axial gauge (ignoring the ghosts).

$$\begin{aligned}
\mathcal{L}^{cov}_{\rm QCD}[\Phi,\mathcal{J}] &= \mathcal{L}_{free}[\Phi]+\mathcal{L}_{int}[\Phi]+\mathcal{L}_{source}[\Phi,\mathcal{J}]\;;\\
\mathcal{L}_{free}[\Phi] &= \sum_f \overline{\psi}_f(i\,\partial\!\!\!/ - m_f)\psi_f\\
&\quad -\frac{1}{4}\,(\partial_\mu\mathcal{A}^a_\nu-\partial_\nu\mathcal{A}^a_\mu)(\partial^\mu\mathcal{A}^{a\,\nu}-\partial^\nu\mathcal{A}^{a\,\mu})-\frac{(\partial\cdot\mathcal{A}^a)^2}{2\xi}\\
&\quad -\overline{c}^a\partial^2 c^a\;,\\
\mathcal{L}_{int}[\Phi] &= -g\sum_f \overline{\psi}_f\mathcal{A}\!\!\!/\psi_f\\
&\quad +\frac{1}{2}\,g f^{a\,b\,c}\mathcal{A}^a_\mu\mathcal{A}^b_\nu\partial^\mu\mathcal{A}^{c\,\nu}-\frac{1}{4}g^2 f^{a\,b\,c}f^{a\,d\,e}\mathcal{A}^{b\,\mu}\mathcal{A}^{c\,\nu}\mathcal{A}^d_\mu\mathcal{A}^e_\nu\\
&\quad +g f^{a\,b\,c}\overline{c}^a\partial^\mu\mathcal{A}^b_\mu c^c\\
\mathcal{L}_{source}[\Phi,\mathcal{J}] &= \overline{\eta}\psi+\overline{\psi}\eta\\
&\quad +j^a_\mu\mathcal{A}^{a\,\mu}\\
&\quad +\overline{\zeta}^a c^a+\overline{c}^a\zeta^a\;.
\end{aligned}\tag{1.32}$$

As advertised, the free part of the lagrangian density is is quadratic in the fields and the interaction part vanishes with the coupling g.[23] We have argued above that *any* correlation function can be obtained by functional differentiation with respect to the sources. This means

as that obtained by other methods in all cases where there are other methods. Such technical issues are taken up to some extent in [38].

[23] Of course, this is the reason g was introduced in the first place.

that we can get *any* functional of the fields under the path integral by judicious use of the source terms. In particular, the functional $\exp[i\mathcal{S}_{int}[\Phi]]$:[24] :

$$\int \mathcal{D}[\Phi] e^{i\mathcal{S}[\Phi,\mathcal{J}]} = e^{i\mathcal{S}_{int}[\frac{1}{i}\frac{\delta}{\delta\mathcal{J}}]} \int \mathcal{D}[\Phi] e^{i(\mathcal{S}_{free}[\Phi]+\mathcal{S}_{source}[\Phi,\mathcal{J}])} \; . \tag{1.33}$$

This is as far as we can go without getting our hands dirty. At this point, we must do the path integral. Fortunately, our manipulations have resulted in a problem we can solve. Our path integral now splits into three integrals - one for each species of particle in our theory (counting ghosts and antighosts as the same species). Since it is only the source dependence that matters, we can complete the square using[25]

$$\Phi^\dagger \mathcal{M} \Phi + \mathcal{J}^\dagger \Phi + \Phi^\dagger \mathcal{J} = (\Phi^\dagger + \mathcal{J}^\dagger \mathcal{M}^{-1}) \mathcal{M} (\Phi + \mathcal{M}^{-1}\mathcal{J}) - \mathcal{J}^\dagger \mathcal{M}^{-1} \mathcal{J} \;\; , \tag{1.34}$$

shift the integration variables, and obtain[26]

$$\begin{aligned} \int \mathcal{D}[\Phi] e^{i\mathcal{S}_{free+source}[\Phi,\mathcal{J}]} \;\; &= \;\; e^{-i\int d^4x\, d^4y\, \mathcal{J}^\dagger(x)\mathcal{M}^{-1}(x,y)\mathcal{J}(y)} \\ &\times \int \mathcal{D}[\Phi] e^{i\int d^4x\, d^4y\, \Phi^\dagger(x)\mathcal{M}(x,y)\Phi(y)} \; . \end{aligned} \tag{1.35}$$

Once again, we ignore the infinite shift to the vacuum energy and obtain

$$\mathcal{Z}[\mathcal{J}] = \frac{e^{i\mathcal{S}_{int}[\frac{1}{i}\frac{\delta}{\delta\mathcal{J}}]} e^{-i\int d^4x\, d^4y\, \mathcal{J}^\dagger(x)\mathcal{M}^{-1}(x,y)\mathcal{J}(y)}}{e^{i\mathcal{S}_{int}[\frac{1}{i}\frac{\delta}{\delta\mathcal{J}}]} e^{-i\int d^4x\, d^4y\, \mathcal{J}^\dagger(x)\mathcal{M}^{-1}(x,y)\mathcal{J}(y)} \, |_{\mathcal{J}=0}} \; . \tag{1.36}$$

All that is left is a calculation of $\mathcal{M}^{-1}$ for each species of particle.

We start with the ghost fields since the inversion is simplest for them. The free ghost matrix is written

$$\mathcal{M}^{a\,b}_{ghost}(x,y) = -\partial^2 \delta^{a\,b} \delta^{(4)}(x-y) \; . \tag{1.37}$$

By definition, the inverse of this matrix satisfies

$$\int d^4z (\mathcal{M}^{-1})^{a\,c}(x,z) \mathcal{M}^{c\,b}(z,y) = \delta^{a\,b}\delta^{(4)}(x-y) \; . \tag{1.38}$$

[24] This is why we must fix our gauge *before* introducing sources. If we did not, we could only couple sources to gauge-invariant field combinations. While this is fine for physical quantities, the gauge-dependent separation of the action into 'free' and 'interaction' contributions would prevent us from introducing perturbation theory.

[25] This expression is valid for any *nonsingular* matrix $\mathcal{M}$. This is one reason we must fix our gauge before we quantize.

[26] This result is valid only for complex fields. For hermitian fields we use $1/2\,\Phi\mathcal{M}\Phi + \mathcal{J}\Phi = 1/2\,\Phi'\mathcal{M}\Phi' - 1/2\,\mathcal{J}\mathcal{M}^{-1}\mathcal{J}$.

Obviously, the $SU(3)$ inverse is trivial : $(\mathcal{M}^{-1})^{a\,b} \sim \delta^{a\,b}$. It is the spatial inversion that we must think about. Since we have not specified a coordinate system, $\mathcal{M}^{-1}$ cannot depend on one. Hence our inverse matrix is a function only of the difference $x-y$. Performing a Fourier transform on this variable, substituting the Fourier representation for the δ-function in $\mathcal{M}$, and inverting the transform we obtain

$$(\tilde{\mathcal{M}}^{-1}_{ghost})^{a\,b}(k) = \frac{\delta^{a\,b}}{k^2} \ , \tag{1.39}$$

or

$$(\mathcal{M}^{-1}_{ghost})^{a\,b}(x,y) = \int \frac{d^4k}{(2\pi)^4}\frac{\delta^{a\,b}}{k^2}e^{-ik\cdot(x-y)} \ . \tag{1.40}$$

In the other two sectors the calculation is similar, but complicated by the presence of spin indices.[27] The matrices

$$\begin{aligned}(\mathcal{M}_{quark})_{\alpha\beta}(x,y) &= \left(i\partial_\mu\gamma^\mu_{\alpha\beta} - m\delta_{\alpha\beta}\right)\delta^{\,(4)}(x-y) \\ (\mathcal{M}_{gluon})^{a\,b}_{\mu\nu} &= \left[g_{\mu\nu}\partial^2 - \left(1-\frac{1}{\xi}\right)\partial_\mu\partial_\nu\right]\delta^{a\,b}\delta^{\,(4)}(x-y)\end{aligned} \tag{1.41}$$

have inverses [28]

$$\begin{aligned}(\mathcal{M}^{-1}_{quark})_{\alpha\beta}(x,y) &= \int \frac{d^4p}{(2\pi)^4}\frac{(\not{p}_{\alpha\beta}+m\delta_{\alpha\beta})}{p^2-m^2}e^{-ip\cdot(x-y)} \\ (\mathcal{M}^{-1}_{gluon})^{a\,b}_{\mu\nu}(x,y) &= \int \frac{d^4q}{(2\pi)^4}\frac{-g_{\mu\nu}+(1-\xi)(q_\mu q_\nu/q^2)}{q^2}\delta^{a\,b}e^{-iq\cdot(x-y)} \ .\end{aligned} \tag{1.42}$$

The physical meaning of these objects becomes clear when we study the two-point correlation functions. For simplicity, we consider the free theory (g=0). The ghost correlation function

$$\begin{aligned}\left\langle \mathrm{T}c^a(x)\overline{c}^b(y)\right\rangle_{vac} &= \frac{1}{i}\frac{\delta}{\delta\overline{\zeta}^a(x)}\left(-\frac{1}{i}\right)\frac{\delta}{\delta\zeta^b(y)} \\ &\qquad \times\ e^{-i\int d^4w\,d^4z\,\overline{\zeta}(w)(\mathcal{M}^{-1}_{ghost})(w,z)\zeta(z)}\Big|_{\zeta=\overline{\zeta}=0} \\ &= i(\mathcal{M}^{-1}_{ghost})(x,y)\end{aligned} \tag{1.43}$$

[27] It is here that we run into trouble if we allow ξ to vary with spacetime. In that case, the Fourier transform does not go through in the same way. Presumably it can be done, but the calculation will be horribly complicated.

[28] Here, we have completely suppressed quark flavor and color indices. Barring the difference in quark mass, these matrices are proportional to the identity in both spaces.

is nothing but i times the inverse ghost matrix.[29] This tells us that the amplitude for a ghost to propagate from y to x is given by $i(\mathcal{M}^{-1}_{ghost})(x,y)$. If $y^0 < x^0$, the ghost is traveling forward in time and should have positive energy. To see which energies contribute, we attempt to do the k^0 integral via contour. The singularities at $k^0 = \pm|\,\vec{k}\,|$ occur on the integration path so their meaning is ambiguous. In order to have a well-defined propagator, we must specify some procedure to handle these singularities. For $y^0 < x^0$, the contour may be closed at infinity in the lower-half plane and we wish for only positive energies to contribute. This is accomplished by pushing the poles $k^0 = |\,\vec{k}\,|$ into the lower-half plane and the poles $k^0 = -|\,\vec{k}\,|$ into the upper-half plane. This physical[30] reasoning leads to the unambiguous propagators

$$\Delta^{a\,b}(k) = \frac{i\delta^{a\,b}}{k^2+i\varepsilon} \tag{1.44}$$

$$S(p) = \frac{i(\not{p}+m)}{p^2-m^2+i\varepsilon} \tag{1.45}$$

$$D^{a\,b}_{\mu\nu}(q) = \frac{[-g_{\mu\nu}+(1-\xi)(q_\mu q_\nu/(q^2+i\varepsilon))]}{q^2+i\varepsilon}\, i\delta^{a\,b} \tag{1.46}$$

for ghosts, quarks, and gluons[31] of momentum k, p, and q, respectively.

In axial gauges, these arguments go through in exactly the same way. The quark propagator is unchanged and the gluon propagator becomes

$$D^{a\,b}_{\mu\nu}(q) = \frac{i}{q^2+i\varepsilon}\left(-g_{\mu\nu}+\frac{n_\mu q_\nu + n_\nu q_\mu}{n\cdot q + i\vec{n}\cdot\vec{q}\varepsilon} - \frac{n^2+\xi q^2}{(n\cdot q+i\vec{n}\cdot\vec{q}\varepsilon)^2}q_\mu q_\nu\right)\delta^{a\,b}\,, \tag{1.47}$$

where I have taken $n^0 > 0$. This prescription is motivated in exactly the same way as above,[32] but here we cannot maintain frame independence. This unpleasing turn of events is not

[29] The signs in this calculation are somewhat tricky. We must remember that the ghost and quark fields and sources are Grassmann variables, and as such *anticommute* rather than commute.

[30] Since the ghosts are not physical particles, the motivation for the use of this prescription is not clear in this context. However, it is essential for the ghost propagator to have the same form as the gluonic propagator if the ghosts are to do their job properly.

[31] Strictly speaking, the $+i\varepsilon$ in the second term of the gluon propagator is not required by physics any more than that in the ghost propagator. These prescriptions need only be consistent with each other to ensure a well-defined gauge theory. It is only a matter of aesthetic convention that we treat them in the same way as the physical propagators.

[32] This prescription is appropriate for ordinary time-ordered perturbation theory, in which one imposes quantization conditions on the fields and their conjugate momenta at equal times. In practice, one often uses the axial gauge with n^μ light-like ($n^2 = 0$) and quantizes the theory at equal $n\cdot x$. In this case, the proper

unexpected since our gauge condition has already broken Lorentz invariance. Of course, the terms in which we need this prescription are merely gauge artifacts and are not required to satisfy the constraints put on physical propagation. Since there are no ghosts in this gauge, there is nothing we must be consistent with. For this reason, the complicated prescription suggested by physical arguments is often ignored. This leads to singularities of a very special kind which are required by gauge invariance to cancel in physical quantities. If they are handled properly, this does indeed happen.[33] However, the special care one must take in the axial gauge can lead to tremendous difficulties. This is one reason the covariant gauges are often employed in spite of the complications due to ghosts.

Although these $i\varepsilon$'s are quite important in many aspects of quantum field theory, there are many other aspects in which they are not essential. For this reason, I will not write them explicitly unless they are expected to contribute. However, it is understood that they are always there.

1.3 Feynman Rules and Perturbation Theory

The free theory is nice, but it cannot be expected to teach us the true nature of the strong interactions. At some point, we must allow our fields to interact with one another. Aside from the mathematical difficulties associated with defining the path integral, everything we have done up to now is exact. In order to go further, we must make some approximations.

As it stands, the generating functional is expressed as a series of derivatives acting on a product of exponentials. Since we cannot calculate the infinite number of derivatives necessary to take the full exponential into account, let us assume for the moment that the coupling g is sufficiently small to justify an expansion :

$$e^{i\mathcal{S}_{int}[\frac{1}{i}\frac{\delta}{\delta\mathcal{J}}]} \sim 1 + i\mathcal{S}_{int}\left[\frac{1}{i}\frac{\delta}{\delta\mathcal{J}}\right] \ . \tag{1.49}$$

prescription is given by

$$\frac{1}{n\cdot q} \to \frac{n^*\cdot q}{(n^*\cdot q)(n\cdot q)+i\varepsilon} \ , \tag{1.48}$$

where $n^{*\mu}$ satisfies $n^*\cdot n = 1$. This rule for handling the singularity, due to Mandelstam and Leibbrandt [39], ensures that only those particles with positive 'light-cone energy', $n\cdot q$, will propagate forward in 'light-cone time', $n\cdot x$. We will see more of this prescription when we study the axial gauge in Section 1.6.

[33] The term 'handled properly' is in some sense defined by this statement.




Figure 1.1: The fundamental quark-gluon interaction is represented graphically by a Feynman diagram. The factor associated with the emission of a gluon with polarization μ and color index a is $-igt^a\gamma^\mu$, along with a δ-function to ensure momentum conservation.

Consider the correlation function[34]

$$\left\langle \mathrm{T}\,\overline{\psi}^i_\alpha(x_1)\psi^j_\beta(x_2)\mathcal{A}^a_\mu(x_3)\right\rangle_{vac} = -\frac{1}{i}\frac{\delta}{\delta\eta^\alpha_i(x_1)}\,\frac{1}{i}\frac{\delta}{\delta\overline{\eta}^\beta_j(x_2)}\,\frac{1}{i}\frac{\delta}{\delta j^\mu_a(x_3)}\mathcal{Z}[\overline{\eta},\eta,j]\Big|_{\overline{\eta}=\eta=j=0}\,. \tag{1.50}$$

In the free theory, this is zero since the external gluon has nowhere to propagate. The way that sources appear in the exponentials implies that there must be an even number of external gluon fields and the same number of ψ's as $\overline{\psi}$'s in order for a matrix element to be nonzero. Taking into account one iteration of the interaction, we see that our correlation function becomes

$$\begin{aligned}\left\langle \mathrm{T}\,\overline{\psi}^i_\alpha(x_1)\psi^j_\beta(x_2)\mathcal{A}^a_\mu(x_3)\right\rangle^{\mathrm{LO}}_{vac} &= \left(-\frac{1}{i}\right)\frac{\delta}{\delta\eta^\alpha_i(x_1)}\,\frac{1}{i}\frac{\delta}{\delta\overline{\eta}^\beta_j(x_2)}\,\frac{1}{i}\frac{\delta}{\delta j^\mu_a(x_3)}\\ &\times\ (-ig)\int d^4w\left(-\frac{1}{i}\right)\frac{\delta}{\delta\eta^k_\gamma(w)}\,\frac{1}{i}\frac{\delta}{\delta j^\nu_b(w)}\gamma^\nu_{\gamma\delta}(t^b)_{k\,l}\,\frac{1}{i}\frac{\delta}{\delta\overline{\eta}^l_\delta(w)}\\ &\times\ e^{-i\int d^4y_1\,d^4y_2\,\overline{\eta}(y_1)S(y_1,y_2)\eta(y_2)}\\ &\times\ e^{-i/2\int d^4z_1\,d^4z_2\,j^\mu(z_1)D_{\mu\nu}(z_1,z_2)j^\nu(z_2)}\,|_{\overline{\eta}=\eta=j=0}\ .\end{aligned} \tag{1.51}$$

Since the sources are set to zero at the end of the calculation, all points must be coupled through propagators. There are only two ways to do this, one of which is zero since the generators of $SU(3)$ are traceless. The other gives

$$\left\langle \mathrm{T}\,\overline{\psi}^i_\alpha(x_1)\psi^j_\beta(x_2)\mathcal{A}^a_\mu(x_3)\right\rangle^{\mathrm{LO}}_{vac} = \int d^4w\,S^{j\,k}_{\beta\gamma}(x_2,w)\left(-ig\gamma^\nu_{\gamma\delta}(t^b)_{k\,l}\right)S^{l\,i}_{\delta\alpha}(w,x_1)D^{b\,a}_{\nu\mu}(w,x_3)\,. \tag{1.52}$$

[34] i and j refer to quark color while α and β refer to spin. Here and in the following, I will ignore the difference between the fundamental and anti-fundamental representations of $SU(3)$. There are simply too many indices floating around here for us to be particular about placement. One can always trace through the derivation to resolve any confusion.




Figure 1.2: This diagram presents a prime example of the reason symmetry factors are sometimes necessary to avoid overcounting. The Feynman rule for the triple gluon vertex counts each of these contributions as separate. In reality, however, only (a) and (b) represent different processes. In order to obtain the correct result using Feynman rules, one must include a factor of 1/2.

This expression consists of three propagators, each from an external point to the common point w. We would like to split it up into different pieces - one for each propagation and one for the interaction itself. However, all of these pieces are at present entangled by the integration over w. In momentum space, the separation is clear :

$$\int d^4x_1 d^4x_2 d^4x_3 e^{-ip_1\cdot x_1} e^{ip_2\cdot x_2} e^{ip_3\cdot x_3} \left\langle \mathrm{T}\,\overline{\psi}^i_\alpha(x_1)\psi^j_\beta(x_2)\mathcal{A}^a_\mu(x_3)\right\rangle^{\mathrm{LO}}_{vac}$$
$$= S^{j\,k}_{\beta\gamma}(p_2)[-ig(t^b)_{k\,l}\gamma^\nu_{\gamma\delta}(2\pi)^4\delta(p_2+p_3-p_1)]S^{l\,i}_{\delta\alpha}(p_1)D^{b\,a}_{\nu\mu}(p_3)\ . \qquad (1.53)$$

The factor associated with the interaction is just $-igt^b\gamma^\nu$ along with a δ-function enforcing energy-momentum conservation. This interaction is represented graphically in Figure 1.1. The straight lines represent quark propagation and the squiggly ones represent gluon propagation.[35] Diagrams such as this one are called Feynman diagrams and are extremely useful for organizing perturbation theory.

The above method can be used to obtain a set of rules, called 'Feynman rules', for all of the terms in the interaction lagrangian. One simply considers a correlation with an external field

[35] On these diagrams, ghost propagation is represented by a dashed line.

for each particle species in the interaction. The rules for all of the interactions in QCD with covariant gauge fixing are given in Appendix A.4. Feynman rules for axial gauge calculations are identical except for the gluon propagator, (1.47), and the fact that there are no ghosts. With these rules, one can in principle calculate any process to any order in perturbation theory:

1) One draws a propagator for each external field in the correlation and connects them in every way possible according to the rules.

2) Disconnected diagrams are discarded, as they will cancel with the normalization.

3) Each interaction vertex comes with an associated δ-function guaranteeing momentum conservation.

4) Unconstrained momenta are integrated over, like every other unconstrained quantum number in the diagram. The number of such integrations is called the number of loops.

5) The anticommuting nature of Grassmann fields generates an extra minus sign for each loop associated with these fields.

6) Certain diagrams come with extra factors because of their high degree of symmetry. These 'symmetry factors' arise because of an inadequacy in our Feynman rules. In the derivation of these rules we assume that the external fields are all distinct. While this is certainly the case if the fields are truly external, it need not be true for internal fields. Consider, for example, the diagrams of Figure 1.2. Our rules count each separate contraction as a distinct contribution to the diagram. However, there are really only two distinct contributions; Figs. 1.2a and d are identical, as are Figs. 1.2b and c. Our rules have overcounted the contribution of this diagram by a factor of two. In general, symmetrical diagrams must be divided by the number of ways one can interchange propagator lines without changing the diagram.

All of these rules arise naturally from the path integral. One can always resort to the tedious method above to ensure the correct symmetry factor or minus sign.

1.4 Renormalization in Covariant Gauges

At this point, it seems like we have all the ingredients necessary for a calculation in QCD. Let's test this hypothesis by applying our procedure to the quark propagator, $\langle \mathrm{T}\, \overline{\psi}(y)\psi(x)\rangle$. This object receives several kinds of corrections, as shown in Fig. 1.3. The diagram in Fig. 1.3a has two disconnected pieces. The gluon loop is not associated with the external sources

and therefore cancels with the denominator in Eq.(1.31). Such disconnected pieces are called vacuum bubbles and represent shifts in the vacuum energy. For our purposes, they can be completely ignored.[36] Diagrams like the one in Fig. 1.3b are called one-particle-reducible (1PR) since they can be broken into two disconnected pieces by cutting only one propagator. These diagrams are special because the momentum of the 'one-particle' propagator is fixed completely by the conservation law. This means that the two parts of this diagram are not coupled by a loop integration and they may be evaluated separately. Hence one can consider only one-particle-irreducible (1PI) diagrams, such as those in Fig. 1.3c, and string them together to get the effect of the 1PR diagrams.

Calling the 1PI quark self-energy $-i\Sigma$, we see that the full correction is[37]

$$\begin{aligned} S(p) &= \frac{i}{\not p - m} + \frac{i}{\not p - m}\,(-i\Sigma(p))\,\frac{i}{\not p - m} + \cdots \\ &= \frac{i}{\not p - m}\left(\frac{1}{1-\Sigma(p)(\not p - m)^{-1}}\right)\,, \end{aligned} \tag{1.55}$$

where I have summed the geometric series. In covariant gauge, Lorentz invariance tells us that Σ can depend only on p^μ. Since the strong interactions respect parity and there is only one vector in the problem, we may express Σ in terms of two scalar functions of p^2

$$\Sigma(p) = A(p^2) + B(p^2)\ \not p\,. \tag{1.56}$$

The explicit breaking of Lorentz invariance in the axial gauges destroys the validity of this expression. The special techniques necessary to handle this and the other problems with renormalization in these gauges will be discussed in Section 1.6.

Substituting this expression for Σ into the fully re-summed propagator, we obtain

$$iS^{-1}(p) = (1 - B(p^2))(\not p - m) - A(p^2) - mB(p^2)\,. \tag{1.57}$$

Unless $A(p^2)$ and $B(p^2)$ are rather trivial functions near $p^2 = m^2$, the corrected propagator no longer has a pole at this position and the residue at the shifted pole is no longer i. The

[36] Although they will not be useful for us, these diagrams do contribute to the effective action of the field theory. Analysis of this object gives insight into the vacuum structure of a theory.

[37] Here, we resort to the standard notation

$$\frac{i}{\not p - m} \tag{1.54}$$

for the bare (uncorrected) quark propagator rather than that found above. This is understood to mean $i(\not p - m)^{-1}$. The $i\varepsilon$ will be ignored for a while, but it is always understood to be present.




Figure 1.3: Three different kinds of contributions to the quark self-energy. The gluon loop in (a) is uncoupled to the sources and so contributes only to the vacuum energy. The reducible graph in (b) does not contain any new information; its contributions can be re-summed with a knowledge of irreducible diagrams, like that in (c).

normalization of the field is a convention, but the fact that it has changed is disconcerting. Although arbitrary, this normalization should be consistent. The resolution of this apparent difficulty rests with the fact that we cannot observe the fields present in our lagrangian. Turning off the interaction so that we can measure this quantity m in the lagrangian is not an option.[38] Hence we are forced to let go of any idea we have that the parameter m present in the lagrangian is a physical observable and that the field ψ is the quark field we see in experiment. These quantities are called 'bare' or 'unrenormalized' objects and require corrections before we can relate them to experiment.

We have the corrected propagator in terms of A and B, so we can calculate the shift in the mass and the new normalization of the quark field. The new pole appears at a renormalized mass $m_{\rm R}$ such that

$$\left(1 - B(m_{\rm R}^2)\right)(m_{\rm R} - m) - A(m_{\rm R}^2) - mB(m_{\rm R}^2) = 0\,, \tag{1.58}$$

and the residue at this pole is given by

$$\begin{aligned} Z_{\rm F} &= \frac{1}{i}{\rm Res}[S(p), m_{\rm R}] \\ &= \frac{1}{1 - B(m_{\rm R}^2) - 2m_{\rm R}\partial\left(A(p^2) + mB(p^2)\right)/\partial p^2\Big|_{p^2=m_{\rm R}^2}}\,. \end{aligned} \tag{1.59}$$

[38] Of course, in QCD seeing the objects under consideration *at all* is not an option. As mentioned in the introduction, defining quark 'mass' is itself difficult. However, we will ignore these difficulties for the discussion at hand as they are nonperturbative problems which do not affect our calculations in perturbation theory. For the rest of this chapter, we can ignore confinement completely. The penalty we pay for such arrogance will be discussed in the next chapter.

Since we have no information on the values of these bare parameters, it is not useful for us to express predictions in terms of them. We would prefer to know the variation of observables with physical quantities. Removing unphysical parameters in favor of physical quantities is called *renormalization*. It turns out that every term in our lagrangian requires renormalization.

The idea of renormalization is not new. Solid state physics has been using this process for many years. Electrons in the conduction band of a metal behave as though their masses are different since the actual excitations observed involve the background lattice. The idea here is the same : we are observing excitations on top of a background vacuum. The vacuum in quantum field theory is a complicated sea of quantum fluctuations and cannot be treated as trivial. The only difference between the solid state example and renormalization in a quantum field theory is that electrons in a metal can be removed from the metal and observed free of any background lattice. They can never be removed from their electromagnetic field fluctuations and observed in their bare form. This is the underlying reason bare quantities have no physical meaning.

There are two main ways to view renormalization. One way is to simply calculate everything in terms of bare parameters and substitute known expressions for the physical quantities at the end. This is known as unrenormalized perturbation theory. The unpleasing thing about unrenormalized perturbation theory is the fact that once the calculation is done, one still must do the substitution - something which is certainly not always trivial. The other main way to renormalize involves a systematic replacement of bare parameters with renormalized ones at each order in perturbation theory. This renormalized version has the advantage that unrenormalized quantities never appear in our expressions. Although it will lead to more diagrams, it is generally preferred to the procrastinator's method.

Before we can begin the task of renormalization, we must carefully define our quantities. We saw above that the normalization of the fermion field changes so that the residue of the pole becomes $iZ_{\rm F}$. Consistency of convention requires the residue for the physical field to be i. This can be obtained by defining[39] $\psi = Z_{\rm F}^{-1/2}\psi_b$ since

$$\int d^4x\, e^{ip\cdot x} \left\langle \mathrm{T}\, \psi(x)\overline{\psi}(0)\right\rangle_{vac} \qquad = Z_{\rm F}^{-1} \int d^4x\, e^{ip\cdot x} \left\langle \mathrm{T}\, \psi_b(x)\overline{\psi}_b(0)\right\rangle_{vac}$$

[39] From this point on, I will use the subscript b for bare parameters and leave renormalized parameters with no subscript.

$$= i\left[\not{p} - m - M(p)(\not{p} - m)^2\right]^{-1} \ , \tag{1.60}$$

where $M(p)$ is defined to take into account the rest of the p^2 variation in A and B. $M(p)$ will have nontrivial Dirac structure, but be regular as $\not{p} \to m$. The normalization of the two-point correlator is governed by the normalization of the kinetic term in the lagrangian, so one might think of this renormalization constant as coming from the operator $\overline{\psi}\, i\partial\!\!\!/ \psi$. It is also associated with the field itself. If we were dealing with a nonlocal field theory, this constant is the only renormalization we would need since nonlocal products of fields are renormalized simply by products of $Z_{\rm F}$. Local field theories are different because the structure of a local composite operator is quite different than that of a nonlocal product. This will become quite explicit in our discussion of the coupling.

The gluon and ghost fields also require a field-strength renormalization. We define $\mathcal{A} = Z_{\rm A}^{-1/2}\mathcal{A}_b$ and $c = Z_{\rm G}^{-1/2}c_b$.[40] There is also a relationship between the renormalized mass and the bare mass. Since the mass term is the only piece of our lagrangian which couples quarks of different chirality, a theory in which m_b=0 is fundamentally different from a theory in which $m_b \neq 0$.[41] A theory of massless fermions cannot generate a fermion mass. For this reason, the renormalized mass must be proportional to the bare mass. We write $m = Z_{\rm m}^{-1}m_b$. The renormalization of the coupling must also be multiplicative since a theory with zero bare coupling is free and certainly will not generate a coupling. Hence we define $g = Z_{\rm g}^{-1}g_b$. We will see below that even the gauge parameter ξ acquires multiplicative renormalization, $\xi = Z_\xi^{-1}\xi_b$.

How does all of this renormalization change our Feynman rules? Obviously, it doesn't change them at all in the unrenormalized version of perturbation theory since all the action happens after the calculation. However, in order to calculate quantities in renormalized perturbation theory, we will have to make some modifications. We begin by making the trivial

[40] Although the ghost and anti-ghost field renormalizations need not be the same since they are unrelated fields, they will always appear together in our applications. Hence for our purposes only the product of their field strengths has any meaning.

[41] Theories with small fermion masses can be approximated by massless theories. The effects of the small breaking of this chiral symmetry induced by nonzero fermion masses can be taken into account in perturbation theory. This is the idea behind Chiral Perturbation Theory, the theory of pions and nucleons mentioned in the introduction.




Figure 1.4: The one-loop correction to the quark self-energy.

observation that

$$\begin{aligned}
\mathcal{L}[\Phi_b, m_b, g_b, \xi_b] &= \mathcal{L}[\Phi, m, g, \xi] \\
&\quad + \sum_f (Z_{\rm F}^f - 1)\overline{\psi}_f \, i \not{\partial} \, \psi_f \\
&\quad - \sum_f m_f (Z_{\rm m}^f Z_{\rm F}^f - 1)\overline{\psi}_f \psi_f \\
&\quad - \sum_f g(Z_{\rm g} Z_{\rm F} Z_{\rm A}^{1/2} - 1)\overline{\psi}_f \not{\mathcal{A}} \psi_f \\
&\quad - \frac{1}{4}\left(Z_{\rm A} - 1\right)(\partial_\mu \mathcal{A}_\nu^a - \partial_\nu \mathcal{A}_\mu^a)(\partial^\mu \mathcal{A}^{a\,\nu} - \partial^\nu \mathcal{A}^{a\,\mu}) \\
&\quad - (Z_{\rm A} Z_\xi^{-1} - 1)\frac{(\partial \cdot \mathcal{A}^a)^2}{2\xi} \\
&\quad + \frac{1}{2} g (Z_{\rm g} Z_{\rm A}^{3/2} - 1) f^{a\,b\,c} \mathcal{A}_\mu^a \mathcal{A}_\nu^b \partial^\mu \mathcal{A}^{c\,\nu} \\
&\quad - \frac{1}{4} g^2 (Z_{\rm g}^2 Z_{\rm A}^2 - 1) f^{a\,b\,c} f^{a\,d\,e} \mathcal{A}^{b\,\mu} \mathcal{A}^{c\,\nu} \mathcal{A}_\mu^d \mathcal{A}_\nu^e \\
&\quad - (Z_{\rm G} - 1)\overline{c}^a \partial^2 c^a \\
&\quad + g(Z_{\rm g} Z_{\rm G} Z_{\rm A}^{1/2} - 1) f^{a\,b\,c} \overline{c}^a \partial^\mu \mathcal{A}_\mu^b c^c \quad .
\end{aligned} \tag{1.61}$$

This is nothing more than a simple rearrangement of terms after substitution. The Feynman rules are the same for the first line of this expression since the lagrangian is the same. All of the other terms are intended to be taken into account in perturbation theory. These terms are all proportional to g since the free theory has $Z = 1$ for all renormalization constants. Hence each of these terms will simply give us another interaction vertex to iterate in our Feynman diagrams. This is the tragedy of renormalized perturbation theory. The price of systematically replacing each bare parameter with the physical couplings is eight new interaction vertices. This is twice as many as we started with! The extra terms are called counterterms because they are constructed to cancel certain contributions which arise in our calculation. For the remainder of this section we will be concerned with calculating their coefficients at leading order in perturbation theory.

Let us begin by actually calculating $A(p^2)$ and $B(p^2)$. There is only one diagram which contributes to these quantities at next-to-leading order (see Fig. 1.4). Using the Feynman rules in Appendix A.4, we find the amplitude for this diagram to be[42]

$$-i\Sigma^{(1-\text{loop})}(p) = \int \frac{d^4k}{(2\pi)^4}\,(-igt^a\gamma^\mu)\left(\frac{i}{\not p - \not k - m_b}\right)(-igt^b\gamma^\nu)D^{a\,b}_{\mu\nu}(k)\;, \tag{1.62}$$

or[43]

$$\Sigma^{(1-\text{loop})}(p) = ig^2C_F\gamma^\mu\int\frac{d^4k}{(2\pi)^2}\frac{(\not p - \not k + m_b)[-g_{\mu\nu}+(1-\xi)(k_\mu k_\nu/k^2)]}{k^2[(p-k)^2-m_b^2]}\gamma^\nu\;. \tag{1.63}$$

A closer look reveals this integral to be divergent. For large k, it behaves as $\int dk/k$.[44] This tells us that extremely large momenta are making contributions to the process. The fact that these asymptotically large momenta contribute to this process irrespective of the size of the characteristic scales in the problem, p^μ and m_b, could imply a deep problem with the theory. It is divergences such as this one which turned people away from quantum field theory in the end of the 1960's. We need not give up so quickly, though. We have just seen that this quantity means nothing by itself. Only properly normalized physical matrix elements expressed in terms of physical quantities are observable. Hence in a strict sense it is unimportant that this quantity diverges.[45] It implies only that the bare parameters must also diverge.

While the above discussion is fine as a theoretical argument, we still have not obtained expressions for the quantities A and B (or $Z_{\rm F}$ and $Z_{\rm m}$). In some sense, we can write $A = 1 + g^2\infty_A^{(2)} + \mathcal{O}(g^4\infty_A^{(4)})$ with a similar expression for B. However, these expressions are not very useful and certainly bring doubts about the convergence of our perturbative expansion.

[42]Remember that the external propagators are not included in the definition of Σ!

[43]The casimir $C_F\delta^i_j = (t^at^a)^i_j$ is discussed in Appendix C.

[44]Since the vector index must be carried by the only vector in the problem, p^μ, the linear divergence $\int dk$ is thrown away by symmetry.

[45] In fact, from a theoretical point of view we might *expect* it to diverge. If it didn't, the bare parameters of the theory would be well-defined finite objects. However, we can think of a bare parameters as the way the particle looks extremely close-up, i.e. at asymptotically high energy. Since there are always corrections at a higher scale than we are observing at, we might not be surprised that there is an infinity behind them all. Looking at renormalization in this way, it also seems quite natural that the divergences from asymptotically large loop momentum combine with the divergent bare parameters representing the way the particles look at asymptotically large momentum transfer to give sensible finite results for physical momentum transfer.

In order to proceed in a constructive way, we must make sense of these divergences. We have already argued that the fact that they diverge is unimportant, but now we need to find a way to express *how* they diverge. The important thing here is that we cannot manipulate infinite expressions.[46] It makes no sense at all to write a relation like $\infty_1 - \infty_2 = 3$. However, it is perfectly correct and meaningful (mathematically) to say $\lim_{x\to\infty}(x + 3 \quad - x) = 3$. What we need to make sense of our theory is a regulator - something to make the divergences finite until we take a certain limit, which we will choose to postpone until the end of the calculation, so that we can manipulate them. As we will see, this necessarily introduces a scale which will act as the separation scale between physics we calculate explicitly and physics contained in our renormalized couplings.[47] Physical processes cannot depend on this arbitrary scale of separation, but intermediate parameters can and will gain dependence on it.

How can we regularize the theory? One simple answer is to merely introduce a high-momentum cutoff in our integrals. Instead of integrating to infinity, we could integrate only to Λ. This procedure has been used before in several theories. However, in gauge theories the price we pay for such savage butchery is high. If we use this regulator, our theory is no longer renormalizable. In unrenormalizable theories, divergences appear which cannot be re-absorbed into the constants we have introduced. For example, if the one-loop quark self-energy in Fig. 1.4 contained a divergence proportional to p^2, we would require a counterterm of the form $\overline{\psi}\partial^2\psi$. In the absence of such a term, the theory is not complete. Unrenormalizable theories need an infinite number of terms in order to be complete. Unless there is a good perturbative expansion which can be truncated, an infinite number of experiments need to be done to fix the renormalized coefficients of each term in the lagrangian. This is far from practical, especially since the theory we began with has only n_f+1 parameters - the quark masses and the coupling, g. The price of the simple cutoff regulator is just too high for us to pay.

[46] There are several schemes which attempt to avoid regularization entirely, defining counterterms during the calculation as sums of divergent integrals in such a way that the renormalized quantities are well-defined (see [40]). It is not clear that the manipulations used to define these schemes are applicable to arbitrary order or for an arbitrary process. At any rate, the scheme introduced below is easier to use and avoids these problems entirely.

[47] This should be obvious from Footnote 45. We cannot say that large momentum quantities combine to give a moderate momentum result without referring to a moderate momentum.

The reason this cutoff procedure exacts such compensation is that renormalizability is not an easy thing to come by - especially in a theory with vector bosons like this one. In fact, the only renormalizable theories involving spin-1 fields are gauge theories. The gauge invariance of these theories plays such a crucial role in the proof [48] that it no longer works if the method of regulation breaks the invariance.[49] Our simple cutoff idea butchers the theory with no respect for either gauge invariance or Lorentz invariance. In fact, there are only a few renormalization schemes which preserve the gauge invariance of our theory. The one which has been proven to be the most useful in many applications, including this one, is called Dimensional Regularization (DR). This scheme was introduced by 't Hooft and Veltman in 1973 expressly because it does not violate gauge invariance. Many of the technical details of DR are spelled out in Appendix E. The basic idea is that the integral in Eq.(1.63) is only divergent because we live in 4 dimensions. If we lived in 3 dimensions, for example, it would be perfectly finite as $k \to \infty$. Furthermore, the theory itself is expected to be well-defined in *any* number of dimensions and its amplitudes are expected to be *meromorphic* functions of the spacetime dimension. These assertions, along with a set of rules that are outlined in Appendix E, allow us to calculate matrix elements in an arbitrary number of dimensions, d. Our divergences will show up as poles in $d-4$. It is only at the end of the calculation that we evaluate these (physical) matrix elements at the spacetime dimension $d = 4$.

The scale we must introduce along with the regularization procedure is mysteriously absent. This is a good thing since the explicit appearance of scales is notorious for breaking gauge invariance. However, as argued above, there must be a scale introduced somewhere. It turns out that the scale in DR is hidden in the coupling constant. To see this, let us analyze QCD in d dimensions.

The mass-dimensions of quantum fields are obtained from the requirement that the action $\mathcal{S} = \int d^d x\, \mathcal{L}$ is dimensionless, implying that $\mathcal{L}$ has mass dimension d. Since derivatives have

[48] This proof is the work that won the Nobel Prize in physics for 't Hooft and Veltman mentioned in the introduction. The relevant references are [10].

[49] It can be shown that certain amounts of violence can be done to the gauge invariance before the theory actually becomes unrenormalizable. What is required is that the gauge invariance can be restored by adding a finite number of new counterterms to the lagrangian. Even so, a procedure in which one need not add any terms to the lagrangian (one which does not break the symmetry) is desirable.

mass dimension 1, the kinetic terms in the lagrangian fix the dimension of the fields : $(d-2)/2$ for bosons and $(d-1)/2$ for fermions. Applying this result to the rest of the lagrangian, we see that the coupling constant g has mass dimension $(4-d)/2$. We would rather work with a dimensionless coupling, so we introduce a scale, μ, to carry the dimension required. The bare coupling must also be given dimension, μ_0. Going back to the bare lagrangian and introducing Z-factors again, we see that μ_0 goes along for the ride; g must be replaced with $g\mu_0^{(4-d)/2}$ everywhere in the renormalized lagrangian. The reason that μ_0 appears rather than μ stems from the fact that Z_g relates the dimensionless couplings g and g_b rather than the dimensionful coefficients $g\mu^{(4-d)/2}$ and $g_b\mu_0^{(4-d)/2}$. We will see how μ_0 is replaced by μ below.

In principle, we are really only interested in $d = 4$, where g is dimensionless and the μ (and μ_0) dependence drops out. However, a consistent theory in d dimensions requires the inclusion of μ. Hence all of our final results should be independent of μ, but its introduction is required by our regulator. A study of the μ dependence of certain intermediate results leads to a deep understanding of renormalization and quantum field theory itself.

Armed with a regularization procedure, we are finally ready to calculate our Z-factors explicitly. Returning to the quark self-energy, this time in d dimensions, we write[50]

$$\Sigma^{(1-\text{loop})}(p) = ig^2\mu_0^{d-4}C_F\int\frac{d^dk}{(2\pi)^d}\left[\frac{(d-2)(\not{p}-\not{k})-dm_b}{k^2[(p-k)^2-m_b^2]}\right.$$
$$\left.+(1-\xi)\frac{(p^2-m_b^2)\not{k}-k^2(\not{p}-m_b)}{(k^2)^2[(p-k)^2-m_b^2]}\right]. \tag{1.64}$$

Using the integration formulae in Appendix A.3, we find[51]

$$A^{(1-\text{loop})}(p^2) = \frac{\alpha_sC_F}{4\pi}\Gamma\left(\frac{\epsilon}{2}\right)\left(\frac{m_b^2}{4\pi\mu_0^2}\right)^{-\epsilon/2}$$
$$\times\, m_b[d-(1-\xi)]\int_0^1 dx\, x^{-\epsilon/2}\left[1+(1-x)\left(-\frac{p^2}{m_b^2}\right)\right]^{-\epsilon/2}$$
$$B^{(1-\text{loop})}(p^2) = \frac{\alpha_sC_F}{4\pi}\Gamma\left(\frac{\epsilon}{2}\right)\left(\frac{m_b^2}{4\pi\mu_0^2}\right)^{-\epsilon/2}$$

[50] To obtain this result, we have used the fact that any term in which the quark propagator $(p-k)^2-m_b^2$ cancels is zero since it does not have any scale dependence. This argument is made in considerably more detail in Appendix E. The rules for d-dimensional γ matrix algebra can be found in Appendix A.1.

[51] Since g appears almost exclusively in the combination $g^2/4\pi$, I have introduced $\alpha_s \equiv g^2/4\pi$. This is the true expansion parameter of the strong interaction.

$$\times \left\{ (2-d)\int_0^1 dx\,(1-x)x^{-\epsilon/2}\left[1+(1-x)\left(-\frac{p^2}{m_b^2}\right)\right]^{-\epsilon/2} \right.$$
$$+(1-\xi)\int_0^1 dx\,x^{-\epsilon/2}\left[1+(1-x)\left(-\frac{p^2}{m^2}\right)\right]^{-\epsilon/2} \tag{1.65}$$
$$\left. +(1-\xi)\,\frac{\epsilon}{2}\,\frac{p^2-m_b^2}{m_b^2}\int_0^1 dx\,x^{-\epsilon/2}(1-x)\left[1+(1-x)\left(-\frac{p^2}{m_b^2}\right)\right]^{-1-\epsilon/2}\right\}\ ,$$

where I have introduced $\epsilon = 4-d > 0$ for convenience. At this point, it seems that finding expressions for $Z_{\rm F}$ and $Z_{\rm m}$ is merely a matter of substitution. In fact, since we only work to a certain order in perturbation theory, this process is easier than it appears at first. Looking at Eq.(1.58), we see that m is in fact defined implicitly. This could present problems - especially in view of the involved form of $A(p^2)$ and $B(p^2)$. At this order, however, we are not concerned with such complications. Since $m-m_b$ is necessarily of order g^2, we can completely ignore the factor $B(m^2)$ multiplying it in Eq.(1.58). Furthermore, the difference between evaluating $A(p^2)$ and $B(p^2)$ at m^2 and m_b^2 is also higher order.[52] These observations lead to the trivial relationship

$$m-m_b = A^{(1-\rm loop)}(m_b^2) + mB^{(1-\rm loop)}(m_b^2) + \mathcal{O}(\alpha_s^2)\,, \tag{1.66}$$

or[53]

$$Z_{\rm m} = 1 - \frac{\alpha_s C_F}{4\pi}\Gamma\left(\frac{\epsilon}{2}\right)\left(\frac{m^2}{4\pi\mu_0^2}\right)^{-\epsilon/2}\frac{3-\epsilon}{2-\epsilon} + \mathcal{O}(\alpha_s^2)\ . \tag{1.67}$$

We run into some difficulty when we attempt to calculate $Z_{\rm F}$. One of the derivatives we wish to take diverges like $\int_0^1 dx\,x^{-1-\epsilon/2}$. This is not a problem for $\epsilon < 0$, but in order to regulate the ultraviolet divergences we require $\epsilon > 0$. The origin of this problem lies in the fact that the quantities A and B are not analytic for ϵ strictly positive. This divergence comes from the region of small k in the integral[54] so it is of a different kind than those we have seen. The fact that it appears at all is in fact quite disheartening since it signals nonanalytic

[52] This is true if and only if $A(p^2)$ and $B(p^2)$ are analytic functions in a region of momentum space which includes both m^2 and m_b^2. We have already assumed that this is the case in defining a 'residue' of the 'pole' at $p^2 = m^2$ since these concepts do not exist unless our denominator is analytic in this region.

[53] Here, I use the fact that the last term in our expression for $B(p^2)$ does not contribute at $\mathcal{O}(\epsilon)$ as $p^2 \to m_b^2$. Due to the singularity in the integral, there will be contributions at higher orders in the ϵ expansion.

[54] This can be seen from the fact that the integrand

$$\sim \frac{1}{k^2[(p-k)^2-m_b^2]}\ , \tag{1.68}$$

behavior of A and B near m_b^2.[55] This has the potential to make the theoretical constructs of the last few pages collapse. Fortunately, there is hope for a solution. The fundamental principle of dimensional regularization is the assumption that the amplitudes in quantum field theory are *meromorphic* functions of spacetime dimension. Obviously, $\Sigma(p)$ is not a meromorphic function of ϵ in its present form since we must put restrictions on the sign of ϵ in order to make sense of it. However, this form can be interpreted as a representation of the function in which its meromorphic nature is not manifest.[56] If we assume that this is the case, we are free to calculate our divergent contribution in the region $\epsilon < 0$ and continue the result to $\epsilon > 0$ (or any other part of the complex ϵ plane). Since this divergence comes from the infrared region of momentum space, it is sometimes useful to keep it separate from the ultraviolet divergences. This is done via subscript. Our result for $Z_{\rm F}$ is

$$\begin{aligned} Z_{\rm F} &= 1 - \frac{\alpha_s C_F}{4\pi}\left(\frac{m^2}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[\Gamma\left(\frac{\epsilon_{\rm UV}}{2}\right)\frac{1+\epsilon}{1-\epsilon}\,\xi \right. \\ &\qquad \left. + \frac{4}{\epsilon_{\rm IR}}\Gamma\left(1+\frac{\epsilon}{2}\right)\right] + \mathcal{O}(\alpha_s^2) \ . \end{aligned} \tag{1.70}$$

The *scheme* we have used to define our renormalized parameters above is called the *onshell* scheme since it defines renormalized quantities at the point where fields are on their mass shells, i.e. $p^2 = m^2$. This is not required by the idea of renormalization. Only the *external* fields must be normalized properly, and the mass at the pole will be the physical mass regardless of how we choose to define m. This fact is exemplified by the existence of unrenor-

so if we differentiate with respect to p^2 and evaluate at $p^2 = m_b^2$, we get the expression

$$\sim \frac{1}{k^2(k^2 - 2p\cdot k)^2} \ . \tag{1.69}$$

This integrand does not cause a problem for large k, but can certainly lead to a divergence at small k.

[55] Strictly speaking, this is not necessarily a problem since we really only need analyticity near the physical pole m^2. However, we are attempting to calculate $m - m_b$ as a power series in α_s. Functions which are analytic at m but not at m_b do not have smooth limits as $\alpha_s \to 0$, hence cannot possess a Taylor expansion about $\alpha_s = 0$. Since Taylor expansions are the only things we know how to calculate in quantum field theory, we are foiled again. At this point, we must either admit defeat and go home or find some way to understand these divergences without compromising the analyticity of A and B near either m^2 or m_b^2.

[56] For example, the function $\sum_{n=0}^{\infty} z^n$ is not a meromorphic function in the complex plane as presented since it does not converge for $|z| \geq 1$. However, summing the series in the region where it converges, we find that it is a perfectly finite function of z for all $z \neq 1$.

malized perturbation theory, where we choose not to renormalize at all. At the end of the day, we simply eliminate the 'bare' parameters in favor of some arbitrarily defined 'renormalized' quantities which will be measured in experiment. In two different renormalization schemes the definitions of the renormalized parameters and the expressions for physical quantities in terms of these parameters will be different, but the combination is guaranteed to be the same. This arbitrariness allows us to make the choice which will make our calculations the simplest. The only residue of the onshell requirements which is left is the normalization of the *external* fields, which *must* be done in the onshell scheme. However, since the fundamental fields of QCD do not appear as external fields, even this condition is unimportant for our purposes.[57]

One might say that the easiest thing to do is simply not renormalize. The problem with this is that all of our parameters are divergent. One cannot measure divergent quantities, so in this case one cannot extract the parameters of the theory from experiment. The next best thing is to do the least amount of work so that our renormalized parameters are finite. This scheme is called *Minimal Subtraction* (MS). In MS, we define our Z-factors in such a way that the counterterms subtract all of the *ultraviolet* divergences in our diagrams at each order in perturbation theory. It must be emphasized here that infrared divergences have nothing to do with renormalization. As mentioned above, the bare parameters of our theory are related to the way things look at asymptotically large probing energies. Hence they have only to do with ultraviolet divergences. Infrared divergences must be handled in an entirely different way, as will be discussed at length in the next chapter. For now, we notice that the infrared divergences in this calculation occur only when we put the fermion on its mass shell. Away from the point $p^2 = m^2$, the amplitudes are infrared finite. This is a general phenomenon - offshell matrix elements are always infrared finite[58] since infrared divergences arise only in singular regions of momentum space. Since we no longer wish to calculate in the onshell scheme, there is no reason to consider this singular portion of phase space.

The counterterms which appear at this order in the quark correlator shift $A(p^2)$ and $B(p^2)$

[57] In QED one must be careful to perform this task before comparing with experiment. Otherwise, inconsistencies will appear between different experiments. We will also see that this step is necessary for calculating 'physical' matrix elements involving quarks and gluons in the next chapter.

[58] This statement is not true for certain gauge choices. In these cases, the infrared divergences come from singular regions of 'gauge choice space'. An explicit example of this phenomenon is given in Section 1.6.

to

$$\begin{aligned} A(p^2) &\rightarrow A(p^2) + m(Z_{\rm m}Z_{\rm F} - 1) \ , \\ B(p^2) &\rightarrow B(p^2) - (Z_{\rm F} - 1) \ . \end{aligned} \tag{1.71}$$

Looking at our expressions for A and B, we see that in the MS scheme

$$\begin{aligned} Z_{\rm F}^{\rm MS} &= 1 - \frac{\alpha_s^{\rm MS} C_F}{4\pi} \frac{2}{\epsilon}(\xi) \ , \\ Z_{\rm m}^{\rm MS} &= 1 - \frac{\alpha_s^{\rm MS} C_F}{4\pi} \frac{2}{\epsilon}(3) \ , \end{aligned} \tag{1.72}$$

where we have explicitly shown the dependence on scheme. The leftover contributions to the self-energy are straightforward to calculate:

$$\begin{aligned} \tilde{A}^{\rm MS}(p^2) &= -\frac{\alpha_s^{\rm MS} C_F}{4\pi} m \left\{ (3+\xi) \left[\log \frac{m^2}{\mu_0^2} + \gamma_E - \log(4\pi) \right.\right. \\ &\qquad \left.\left. + \left(1 - \frac{m^2}{p^2}\right) \log\left(1 - \frac{p^2}{m^2}\right) \right] - 2\,(2+\xi) \right\} + \mathcal{O}(\alpha_s^2) \ , \qquad (1.73) \\ \tilde{B}^{\rm MS}(p^2) &= \frac{\alpha_s^{\rm MS} C_F}{4\pi} \xi \left[\log \frac{m^2}{\mu_0^2} + \gamma_E - \log(4\pi) \right. \\ &\qquad \left. + \left(1 - \frac{m^4}{p^4}\right) \log\left(1 - \frac{p^2}{m^2}\right) - 1 - \frac{m^2}{p^2} \right] + \mathcal{O}(\alpha_s^2) \ . \qquad (1.74) \end{aligned}$$

These expressions are somewhat messy. The Euler-Masceroni constant γ_E comes from the expansion of the Γ-function and the $\log 4\pi$ comes from the d-dimensional phase space factor $(4\pi)^d$. A general ℓ-loop graph will contain ℓ powers of the coupling and the phase space, so divergences will in general be expressed as[59]

$$\Gamma^\ell \left(\frac{\epsilon}{2}\right) \left(\frac{\Lambda^2}{4\pi\mu^2}\right)^{-\ell\epsilon/2} \sim \left(\frac{2}{\epsilon}\right)^\ell \left(\frac{\Lambda^2 e^{\gamma_E}}{4\pi\mu^2}\right)^{-\ell\epsilon/2} \ , \tag{1.75}$$

where I have used the truncated expansion $\log\ \Gamma(1+\epsilon) \sim -\gamma_E \epsilon$ and the fact that $n\Gamma(n) = \Gamma(n+1)$. Λ is some mass scale in the process, i.e. $\Lambda \sim m$. Since these factors will always appear with the divergences, it is convenient to include them in the Z-factors. Furthermore, we are still plagued with the bare scale μ_0. A fully renormalized quantity should not depend on such an object. Z-factors relate renormalized and unrenormalized quantities, so it is natural

[59] Here, I have written the form of the most UV-divergent part of an ℓ-loop diagram. There will, of course, be less divergent pieces. The γ_E's and 4π's in these terms will also arrange themselves in a similar way, although this is more difficult to see.

to absorb the μ_0 dependence into Z. Doing this requires the introduction of a scale, which we choose as μ. Hence our divergence is written

$$\left[\left(\frac{2}{\epsilon}\right)^{\ell}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\ell\epsilon/2}\right]\left(\frac{\Lambda^2 e^{\gamma_E}}{4\pi\mu^2}\right)^{-\ell\epsilon/2} \quad . \tag{1.76}$$

The quantity in brackets is associated with the divergence (the transition from bare to renormalized quantities), and so is seen as a part of Z. The leftover piece depends only on physical parameters and the renormalized scale, and is free to enter renormalized amplitudes.

The scheme described above is called *Modified Minimal Subtraction* ($\overline{\rm MS}$) and will be used throughout the rest of this dissertation. In $\overline{\rm MS}$, $Z_{\rm F}$ and $Z_{\rm m}$ are written

$$\begin{aligned} Z_{\rm F}^{\overline{\rm MS}} &= 1 - \frac{\alpha_s^{\overline{\rm MS}}(\mu^2)C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(\xi) + \mathcal{O}(\alpha_s^2) \ , \\ Z_{\rm m}^{\overline{\rm MS}} &= 1 - \frac{\alpha_s^{\overline{\rm MS}}(\mu^2)C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(3) + \mathcal{O}(\alpha_s^2) \ , \end{aligned} \tag{1.77}$$

where we have explicitly shown the dependence of α_s on μ^2. It is obvious that this object will depend on μ^2 since Z_g depends on μ^2 and g_b does not. All renormalized quantities will depend on μ. For example, we see that

$$\begin{aligned} \mu\frac{d\,m^{\overline{\rm MS}}}{d\mu} &= -\frac{1}{Z_{\rm m}^{\overline{\rm MS}}}\mu\frac{d\,Z_{\rm m}^{\overline{\rm MS}}}{d\mu}m^{\overline{\rm MS}} \\ &= -3\frac{\alpha_s^{\overline{\rm MS}}(\mu^2)C_F}{2\pi}m^{\overline{\rm MS}} + \mathcal{O}(\alpha_s^2) \ . \end{aligned} \tag{1.78}$$

This means that QCD runs quark masses down as the probing scale is increased. This object is in fact not the physical pole mass of the quark (even in the absence of confinement). The physical pole mass, as defined in the onshell scheme, must be independent of μ since it is an observable. We note here that it is natural for these quantities to depend on scale since the scale decides which corrections are to be included in renormalized quantities and which are to be included explicitly in the amplitudes.

The renormalized quantities $\tilde{A}$ and $\tilde{B}$ can also be calculated,

$$\begin{aligned} \tilde{A}^{\overline{\rm MS}}(p^2) &= -\frac{\alpha_s^{\overline{\rm MS}}(\mu^2)C_F}{4\pi}m\left\{(3+\xi)\left[\log\frac{m^2}{\mu^2}\right.\right. \\ &\quad \left.\left. + \left(1-\frac{m^2}{p^2}\right)\log\left(1-\frac{p^2}{m^2}\right)\right] - 2\,(2+\xi)\right\} + \mathcal{O}(\alpha_s^2) \ , \end{aligned} \tag{1.79}$$

$$\begin{aligned} \tilde{B}^{\overline{\rm MS}}(p^2) &= \frac{\alpha_s^{\overline{\rm MS}}(\mu^2)C_F}{4\pi}\xi\left[\log\frac{m^2}{\mu^2}\right. \\ &\quad \left. + \left(1-\frac{m^4}{p^4}\right)\log\left(1-\frac{p^2}{m^2}\right) - 1 - \frac{m^2}{p^2}\right] + \mathcal{O}(\alpha_s^2) \ , \end{aligned} \tag{1.80}$$



Figure 1.5: The one-loop correction to the ghost self-energy.

and are somewhat cleaner in this scheme. Note the branch cut in our amplitude at $p^2 = m^2$. This cut leads to the infrared divergence in the onshell scheme. It is related to the possibility that the intermediate states in our calculation are onshell and free to propagate to infinity (in the absence of confinement, of course). The condition for this possibility is $(p-k)^2 = m^2$ and $k^2 = 0$, while $(p-k)^0$ and k^0 are positive. Working in the rest frame of $p-k$, one can show that these conditions imply $p^2 = m^2 + 2mk^0 \geq m^2$. Having learned our lesson from venturing into the onshell scheme, we choose to work far away from this kinematical region and take $p^2 < 0$.

We still have one point which must be addressed. What should be used for μ? If we wanted to calculate to all orders, this choice would be arbitrary since no physical quantities actually depend on μ. However, we have no intention of calculating *any* physical quantity to all orders and no hope of doing so in a finite amount of time (at least, not at this stage of the game). With this in mind, our choice of μ should be dictated by the size of the coefficient of α_s. For $-p^2 \ll m^2$, the leftover terms in A and B are all small except possibly the μ-dependent logarithm. Choosing $\mu^2 = m^2$ eliminates this problem. However, for $-p^2 \gg m^2$ this choice would lead to a term $\sim \log(-p^2/m^2)$ in the amplitude, rendering the perturbative expansion useless. In this region, it is preferable to choose $\mu^2 = -p^2$. Hence we see that for the one-loop quark self-energy the natural choice of μ^2 is the larger of $-p^2$ and m^2. Of course, we have no guarantee that these choices will not lead to disaster at higher orders in the expansion. The best choice of μ is obviously the one which makes *all* corrections as small as possible.[60]

The ghost propagator is renormalized in exactly the same way. The relevant diagram (Fig.

[60] In fact, one way to estimate the contributions of higher-order effects is to study the dependence of physical quantities on μ^2. In regions where the dependence is slight, our corrections should be small.

1.5) is almost identical to the quark self-energy. We obtain[61]

$$Z_{\rm G} = 1 + \frac{\alpha_s(\mu^2) C_A}{16\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} (3-\xi) + \mathcal{O}(\alpha_s^2) \ , \tag{1.81}$$

where C_A is defined in Appendix C.

The three diagrams shown in Fig. 1.6 contribute to the field strength renormalization of the gluon. Denoting the sum of the amputated 1PI gluon self-energy insertions as $-i\Pi^{\mu\nu}$, the quark loop (Figure 1.6a) contributes[62]

$$\begin{aligned} -i\Pi_{\rm F}^{\mu\nu}(q) &= -i\frac{2\alpha_s(\mu^2) T_{\rm F}}{\pi} \Gamma\left(\frac{\epsilon}{2}\right) \sum_f \left(\frac{m_f^2}{4\pi\mu_0^2} \right)^{-\epsilon/2} \\ &\times \int_0^1 dx\, x(1-x) \left[1 + x(1-x) \left(\frac{-q^2}{m^2} \right) \right]^{-\epsilon/2} \left(q^2 g^{\mu\nu} - q^\mu q^\nu \right) \ . \end{aligned} \tag{1.82}$$

The transverse structure $q^2 g^{\mu\nu} - q^\mu q^\nu$ appears as a consequence of current conservation, $q_\mu \Pi^{\mu\nu}(q) = 0$. This property of QCD is essential to its renormalizability [10, 40]; we will see in Appendix D.2 that it persists to all orders. The gluon and ghost loop contributions (Figs. 1.6b and c) do not satisfy this condition separately; their contributions have the UV divergences

$$\begin{aligned} -i(\Pi_{\rm A}^{\mu\nu})_{\rm UV}(q) &= i\frac{\alpha_s(\mu^2) C_A}{32\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \\ &\times \left[\left(\frac{53}{3} - 4\xi \right) \left(q^2 g^{\mu\nu} - q^\mu q^\nu \right) - \left(q^2 g^{\mu\nu} + q^\mu q^\nu \right) \right] \end{aligned} \tag{1.83}$$

$$\begin{aligned} -i(\Pi_{\rm G}^{\mu\nu})_{\rm UV}(q) &= i\frac{\alpha_s(\mu^2) C_A}{32\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\frac{\epsilon}{2}} \\ &\times \left[-\frac{1}{3} \left(q^2 g^{\mu\nu} - q^\mu q^\nu \right) + \left(q^2 g^{\mu\nu} + q^\mu q^\nu \right) \right] \end{aligned} \tag{1.84}$$

for the gluons and ghosts, respectively. Note that although each contribution separately violates current conservation, the sum does not. This is the reason the ghosts needed to be introduced - they exist only to cancel the unphysical gluonic degrees of freedom which appear in covariant gauges.

$\Pi^{\mu\nu}$ receives two counterterm contributions. The field strength contributes $-i(Z_{\rm A} - 1)(q^2 g^{\mu\nu} - q^\mu q^\nu)$, while the gauge parameter gives $i(Z_{\rm A} Z_\xi^{-1} - 1) q^\mu q^\nu / \xi$. Since all of our

[61] From here on, I will drop the superscript $\overline{\rm MS}$ on Z-factors and the like. All results will be given in this scheme unless otherwise specified.

[62] The $SU(3)$ generators are normalized by the relation $T_{\rm F} \delta^{a\,b} = {\rm Tr}\,[t^a t^b]$.




Figure 1.6: The three contributions to the gluon field-strength renormalization at one-loop order in covariant QCD.

divergences are proportional to the transverse structure, we can simply take $Z_{\rm A} = Z_\xi$ at this order. In fact, the counterterm relevant to the gauge parameter leads to nonconservation of our current. This is not surprising since this term came from our gauge fixing, so certainly cannot be expected to be gauge invariant. However, it can be shown that even the gauge fixed theory will conserve current [41], which implies $Z_\xi = Z_{\rm A}$ to all orders. We can see this in another way by returning to unrenormalized perturbation theory for a moment. Writing the full 1PI gluon contribution as

$$-i\Pi^{a\,b}_{\mu\nu}(q^2) = -i\Pi(q^2)(q^\mu q^\nu - q^2 g^{\mu\nu})\delta^{a\,b} \tag{1.85}$$

and re-summing the 1PR contributions, we obtain the full gluon propagator

$$D^{a\,b}_{\mu\nu}(q) = \frac{i\left[-g_{\mu\nu} + \left(1-(1-\Pi(q^2))\xi_b\right)(q_\mu q_\nu/q^2)\right]}{(1-\Pi(q^2))q^2}\delta^{a\,b}\,. \tag{1.86}$$

It is now obvious that the gluon fields and the gauge parameter are renormalized in the same way to all orders. This is a direct consequence of our assumption (1.85) that the 1PI gluon contribution is transverse.

Defining the other counterterm in such a way that it cancels the divergent contributions to Π, we find

$$Z_{\rm A} = 1 + \frac{\alpha_s(\mu^2)}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[\frac{1}{2}\left(\frac{13}{3}-\xi\right)C_A - \frac{2n_f}{3}(2T_F)\right]\quad, \tag{1.87}$$

where n_f is the number of flavors.

The leftover finite contribution to Π has a serious problem coming from the quark loop (Figure 1.6a). As in the quark self-energy diagram analyzed above, this expression contains logarithms involving the quark masses, the external momentum scale, and the renormalization scale. Once again, these logarithms become large unless we choose the renormalization scale

near the larger of the two physical scales. However, we cannot just arbitrarily choose μ^2 to satisfy this one particular diagram; there are two other diagrams which also contribute. These other diagrams demand a renormalization scale comparable to the external momentum (q^2). If any of the quarks in our theory have masses much larger than this scale, Figure 1.6a will lead to large finite corrections to the amplitude.

From a physical point of view, this is nonsense - extremely heavy quarks should decouple from low energy phenomena. The problem lies in our renormalization procedure itself. We are choosing counterterms to cancel *all* ultraviolet divergences, regardless of the scale at which they become important. This is not exactly what we want. The idea of renormalization is to choose a scale and calculate explicitly all physics below that scale, while systematically grouping the physics above it into constants which renormalize the fields and parameters of the theory. Since the heavy quark loop has support only for loop momenta near or above its mass scale, choosing a small renormalization scale should make its contribution to the renormalized amplitudes vanish. What we see is quite the contrary : the contribution of the quark loop diagram to the renormalized amplitudes is divergent as the quark mass goes to infinity. Since this diagram has also contributed to the infinite constant Z_{A}, and the strict limit $m_f \to \infty$ gives no contribution to the amplitude[63], we can interpret the divergence in the renormalized amplitude as an attempt to cancel the divergence in Z_{A}. This procedure leads us to believe that we should simply ignore the effects of very heavy quarks; the associated physics will be grouped into renormalization constants with the rest of the high-energy behavior of the theory. On the other hand, we cannot expect all of the physics associated with a quark to be represented by a simple constant. If it could be, QCD would certainly not have such rich structure. The complete dynamics associated with a heavy quark must be represented instead by a series of local operators which give the heavy quark effects the freedom to depend on external momenta. This is the hallmark of an unrenormalizable theory. Expanding the integrand of the quark loop contribution in a power series in $1/m_f$, we see that each term is more divergent than the last.[64] This implies that we will need an infinite number of counterterms to cancel them,

[63]This can be seen explicitly from the expression for the diagram; one takes m_f to ∞ *before* doing the integral.

[64]Of course, this is because our expansion does not converge for large loop momenta. These new divergences did not come out of thin air.

another indication that the effective theory is unrenormalizable.

The explicit form of this theory can be obtained by simply doing the path integral for the heavy quark fields. The result, up to a trivial constant factor which cancels with the vacuum fluctuations, is

$$\det\left(1-\frac{i\not{D}}{m_f}\right) \quad . \tag{1.88}$$

This determinant can be exponentiated via the identity

$$\log\det M = \mathrm{Tr}\log M \quad , \tag{1.89}$$

valid for any nonsingular matrix M, and added to the lagrangian of the theory. The expansion of the logarithm gives the infinite series of local operators required to encode the effects of the heavy quark.

However simple this expansion looks, it is certainly not clear what the trace means in this context. For this reason, one usually considers instead the equivalent[65] determinant

$$\det\left(1-(i\not{\partial}-m_f)^{-1}g\not{A}\right) \quad . \tag{1.90}$$

Here, the trace of a particular term in the expansion of the logarithm is represented as a heavy quark loop with the appropriate number of external gluon fields.[66] The contribution from each loop is calculated according to the above rules, with the gluon fields treated as external sources. At this point, there is no difference between the present calculation and the one that led to (1.82). The difference comes in our *interpretation* of the ultraviolet divergences.[67] These divergences have the form

$$\frac{2}{\epsilon}\left(\frac{m_f^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \quad . \tag{1.91}$$

This expression represents a battle between two different limits. If $\epsilon\log(m_f/\mu)\ll 1$, we have an ultraviolet divergence which is handled as before by renormalizing the gluon field strength. On the other hand, if $\epsilon\log(m_f/\mu)\gg 1$, this term approaches zero. Since we *cannot* take ϵ

[65] again, up to a trivial constant

[66] One simply identifies the factor $(i\not{\partial}-m_f)^{-1}$ with the free propagator for the heavy quark.

[67] Loops with five or more gluon fields are ultraviolet finite. These contributions are not as subtle as those that presently concern us.

strictly to zero in expressions of this form, the limit $m_f \to \infty$ annihilates them.[68] Contributions which are ultraviolet finite do not require ϵ to regulate them. Here, we can take ϵ strictly to zero. Since m_f is not *strictly* infinity, the factor

$$\left(\frac{m_f^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \to 1 \ . \tag{1.92}$$

Grouping the heavy quark effects into a series of local operators is now straightforward. We simply calculate the contribution from each *nonlocal* term in the expansion of (1.90), throw the ultraviolet divergence (if any) away, and expand the remainder in a series in $1/m_f$. Each term in the expansion is then replaced by an effective operator and added to the lagrangian. As an example, we calculate explicitly the two-gluon contribution :

$$\begin{aligned}
i\int d^4x\, \mathcal{O}_2(x) &= -\frac{g^2}{2}\int d^4x\, d^4y\, \mathrm{Tr}\left[\mathcal{M}^{-1}_{quark}(x,y)\not{\mathcal{A}}(y)\mathcal{M}^{-1}_{quark}(y,x)\not{\mathcal{A}}(x)\right] \\
&= -\frac{g^2 T_F}{2}\int \frac{d^4q}{(2\pi)^4}\tilde{\mathcal{A}}^a_\mu(q)\tilde{\mathcal{A}}^a_\nu(-q) \\
&\qquad \times \int \frac{d^dk}{(2\pi)^d}\frac{\mathrm{Tr}\left[(\not{k}+\not{q}+m_f)\gamma^\mu(\not{k}+m_f)\gamma^\nu\right]}{[(k+q)^2-m_f^2](k^2-m_f^2)} \\
&\to -i\frac{\alpha_s T_F}{10\pi}\int \frac{d^4q}{(2\pi)^4}\tilde{\mathcal{A}}^a_\mu(q)\tilde{\mathcal{A}}^a_\nu(-q)\left(q^2 g^{\mu\nu}-q^\mu q^\nu\right)\left(\frac{q^2}{m_f^2}\right) \\
&\qquad \times \left[\frac{1}{3}+\frac{1}{28}\,\frac{q^2}{m_f^2}+\cdots\right] \\
&= -i\frac{\alpha_s T_F}{10\pi}\int d^4x \mathcal{A}^a_\mu(x)\left(g^{\mu\nu}\partial^2-\partial^\mu\partial^\nu\right)\left(\frac{\partial^2}{m_f^2}\right) \\
&\qquad \times \left[\frac{1}{3}-\frac{1}{28}\,\frac{\partial^2}{m_f^2}+\cdots\right]\mathcal{A}^a_\nu(x) \ .
\end{aligned} \tag{1.93}$$

The final form of the local operator can certainly be expressed in a nicer form. In particular, for an abelian theory it is entirely equivalent to the gauge-invariant combination

$$\mathcal{O}_2 = \frac{\alpha_s T_F}{20\pi}\mathcal{F}^{\mu\nu}\left(\frac{1}{3}-\frac{1}{28}\,\frac{\partial^2}{m_f^2}+\cdots\right)\frac{\partial^2}{m_f^2}\,\mathcal{F}_{\mu\nu} \ . \tag{1.94}$$

For non-abelian theories, it is certainly not clear whether or not the higher order terms in our expansion of (1.90) will conspire to form a gauge-invariant combination at the end of the day. On one hand, we know that the original lagrangian is gauge invariant so we would expect

[68] One needn't worry about this expression for light quarks. As we have seen above, the terms reorganize themselves in this case; μ^2 is compared with q^2 rather than m_f^2.




Figure 1.7: The two vertex correction diagrams at one-loop order in QCD.

gauge invariance of our effective lagrangian as well. On the other, the determinant (1.90) is not obviously gauge invariant due to our cancelation of the infinite constant

$$\det\left(i\,\partial\!\!\!/ - m_f\right) \quad . \tag{1.95}$$

In any case, we will consider these correction terms small enough to ignore completely. This leads us to the result (1.87) obtained previously for the gluon field strength renormalization, but with n_f interpreted as the number of *active* flavors.

Now, we need only calculate Z_g. This quantity appears in several counterterms. Unless our gauge symmetry is broken by quantum effects, the g's in our lagrangian must remain the same after renormalization. This can be checked explicitly by calculating Z_g in all four different ways. However, a general analysis of the ways a symmetry can be broken by quantum effects tells us that vector symmetries (symmetries that treat the left- and right-handed fermions in the same way) are preserved.[69] Trusting this, we calculate the one-loop corrections to the quark-gluon vertex shown in Figure 1.7. Since we wish to work in $\overline{\rm MS}$, only the ultraviolet divergent parts of the amplitudes are required to deduce Z_g. Diagram 1.7a. contains the divergence

$$\frac{\alpha_s}{4\pi}\left(C_F - \frac{C_A}{2}\right)\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(\xi) \quad , \tag{1.96}$$

while 1.7b. contains

$$\frac{\alpha_s}{4\pi}\left(C_A\right)\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\frac{3}{4}\left(1+\xi\right) \quad . \tag{1.97}$$

Here, we have written only the constant factor multiplying the leading structure, $-igt^a\gamma^\mu$. In this language, the counterterm contributes

$$Z_{\rm F} Z_{\rm A}^{1/2} Z_g - 1 \quad . \tag{1.98}$$

[69] An exception to this rule is conformal symmetry, which is broken *explicitly* by our regularization procedure.

Putting all of this together and requiring the sum to be ultraviolet finite gives

$$Z_g = 1 - \frac{\alpha_s(\mu^2)}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[\frac{11}{6}C_A - \frac{2T_F}{3}n_f\right] \ . \tag{1.99}$$

As before, n_f is the number of quark flavors whose mass is less than the probing scale. We notice at once that the gauge dependence has canceled. This is necessary because the gauge coupling is independent of gauge choice. It would certainly be strange if renormalization required this quantity to acquire gauge dependence. However, the cancelation is highly nontrivial. Both of the diagrams, as well as the field strength renormalizations, are gauge-dependent. The gauge dependence of these diagrams is not truly an artifact of the coupling itself, but rather is associated with the external fields.

1.5 The Beta Function - Asymptotic Freedom

As we have seen, the ideas behind renormalization make it necessary for renormalized parameters to depend on the scale of separation between moderate and ultraviolet momenta. Although true physical observables are in fact independent of this scale, any truncation of the coupling expansion will introduce dependence. Furthermore, our *effective* coupling really does depend on scale. This scale dependence governs the region of applicability of perturbation theory, as we will see below.

The variation of the QCD coupling with scale is summarized in the QCD β-function, defined via

$$\beta(\alpha_s) \equiv \mu^2\frac{d\alpha_s}{d\mu^2} \ . \tag{1.100}$$

Since this dependence comes only from Z_g, the β-function will not depend on any scales except μ^2. Furthermore, the fact that it is dimensionless implies that even this dependence comes exclusively from $\alpha_s(\mu^2)$.[70] Since $g = Z_g^{-1}g_b$ and g_b is independent of μ, we have the relation

$$\beta(\alpha_s) = -2\alpha_s\, Z_g^{-1}\mu^2\frac{dZ_g}{d\mu^2} \ . \tag{1.101}$$

The leading order result for Z_g obtained in the last section gives

$$\beta(\alpha_s) = -\alpha_s^2(\mu^2)\frac{1}{4\pi}\left(\frac{11}{3}C_A - \frac{4}{3}n_f T_F\right) + \mathcal{O}(\alpha_s^3) \ . \tag{1.102}$$

[70] Since the β-function is manifestly finite *when expressed in terms of the renormalized coupling*, factors of $(\mu/\mu_0)^{-\epsilon} \to 1$.

A look at the QCD lagrangian, Eq.(1.5), tells us that the gluon fields are properly normalized for $T_F = 1/2$.[71] Since $C_A = 3$ for $SU(3)$ with this normalization, the leading order β-function is negative in QCD with less than seventeen active flavors. This means that if we start with a small enough coupling to apply perturbation theory, the coupling will decrease as its scale increases. A theory whose β-function exhibits this behavior is called asymptotically free.

The analysis of the last section tells us that the natural choice of μ^2 is the probing scale relevant to our process. A different choice will lead to large coefficients of the expansion due to leftover logarithms. In this sense, the variation of α_s with scale is a physical phenomenon. Asymptotic freedom assures us that QCD behaves as a weakly coupled theory when probed at high energy.[72] On the other hand, our expression for the β-function also implies that the effective coupling grows we decrease the probing scale. Truncating the β-function at leading order and solving (1.100) gives

$$\alpha_s(\mu_2^2) = \frac{\alpha_s(\mu_1^2)}{1 + \beta_0 \alpha_s(\mu_1^2) \log\left(\mu_2^2/\mu_1^2\right)} \quad , \tag{1.103}$$

where I have introduced $\beta_0 \equiv (11 - 2n_f/3)/4\pi$. Were this exact, our coupling would diverge at the scale

$$\mu_2^2 = \mu_1^2 e^{-\alpha_s(\mu_1^2)/\beta_0} \quad . \tag{1.104}$$

The truncation of the perturbative expansion will loose its validity long before the coupling diverges, so this is certainly not a physical effect. However, it does imply that there exists a scale small enough that QCD is nonperturbative. Hence we see that asymptotic freedom is a two-edged sword - it implies both that perturbation theory will be applicable at high energy *and* that it will not be applicable at low energy. Since low energy is related to long distance phenomena, this result means that the interaction between two quarks will grow as they are separated until perturbation theory breaks down and their behavior becomes incalculable. One can use this as a heuristic motivation for confinement, although it certainly does not constitute a proof.

[71] Appropriately normalized real vector fields have the kinetic term $-1/2\ \partial^\mu \mathcal{A}^\nu \partial_\mu \mathcal{A}_\nu$.

[72] Assuming, of course, that there exists some energy at which the coupling is small enough to apply perturbation theory. This assumption has been thoroughly justified by experiment, as exemplified by the observation of scaling in DIS.

Although asymptotic freedom tells us that our coupling will formally vanish as the energy is increased, it does not tell us at what energy it will become small. Experimentally, α_s is measured at the Z-boson mass (91.19 GeV) as

$$\alpha_s(M_Z^2) = 0.119(2) \ . \tag{1.105}$$

While this certainly seems small enough to apply perturbation theory, higher order effects should be measurable. For this reason, a systematic treatment of the corrections to the leading β-function is desired.

In general, one writes

$$\beta(\alpha_s) = -\beta_0\alpha_s^2 - \beta_1\alpha_s^3 - \cdots \ . \tag{1.106}$$

Of course, our renormalized coupling will depend on the scheme we use to define it. The relationship between the renormalized couplings in two different schemes can be calculated in perturbation theory :

$$\tilde{\alpha}_s = \alpha_s + \gamma\alpha_s^2 + \delta\alpha_s^3 + \cdots \ . \tag{1.107}$$

To see how the β-function depends on scheme, we write[73]

$$\tilde{\beta} = \beta + 2\gamma\alpha_s\beta + 3\delta\alpha_s^2\beta + \cdots \ . \tag{1.108}$$

Expanding β in terms of α_s and using the relations

$$\begin{aligned} \alpha_s^2 &= \tilde{\alpha}_s^2 - 2\gamma\tilde{\alpha}_s^3 + (5\gamma^2 - 2\delta)\tilde{\alpha}_s^4 + \cdots \ , \\ \alpha_s^3 &= \tilde{\alpha}_s^3 - 3\gamma\tilde{\alpha}_s^4 + \cdots \ , \\ \alpha_s^4 &= \tilde{\alpha}_s^4 + \cdots \ , \end{aligned} \tag{1.109}$$

we find

$$\begin{aligned} \tilde{\beta}_0 &= \beta_0 \\ \tilde{\beta}_1 &= \beta_1 \\ \tilde{\beta}_2 &= \beta_2 - \gamma\beta_1 - (\gamma^2 - \delta)\beta_0 \ . \end{aligned} \tag{1.110}$$

[73] We assume here that the difference between the two schemes is not scale dependent, i.e. γ and δ do not depend on scale.

Hence the first two coefficients of the β-function are independent of scheme[74], but the rest will depend on the conventions used to define α_s.

Since these higher-order terms depend on scheme, they may be chosen in any way we like. A convenient choice *defines*

$$\beta_n \equiv \left(\frac{\beta_1}{\beta_0}\right)^n \beta_0 \quad , \tag{1.111}$$

allowing us to re-sum the β-function to all orders in α_s:[75]

$$\beta(\alpha_s) = -\frac{\beta_0 \alpha_s^2}{1 - \beta_1 \alpha_s / \beta_0} \quad . \tag{1.112}$$

This function is not Lipshitz continuous near $\alpha_s = \beta_0/\beta_1$, so we have no guarantee that a solution to (1.100) exists. In fact, an analysis of this equation reveals two distinct branches of solutions. The region $\alpha_s < \beta_0/\beta_1$ is completely disjoint from $\alpha_s > \beta_0/\beta_1$. In the latter case, the β-function is *not* aymptotically free. Since asymptotic freedom is observed in nature, only the former region is physically realized.[76] However, the solution to Eq.(1.100) in this region,

$$\beta_0 \log \frac{\mu_2^2}{\mu_1^2} = \frac{1}{\alpha_s(\mu_2^2)} - \frac{1}{\alpha_s(\mu_1^2)} + \frac{\beta_1}{\beta_0} \log \frac{\alpha_s(\mu_2^2)}{\alpha_s(\mu_1^2)} \quad , \tag{1.113}$$

does not exist for arbitrary μ_2^2. It is readily seen that one must take

$$\mu_2^2 \geq \left(\frac{\beta_0}{\alpha_s(\mu_1^2)\beta_1}\right)^{\beta_1/\beta_0^2} \exp\left[-\frac{1}{\beta_0}\left(\frac{1}{\alpha_s(\mu_1^2)} - \frac{\beta_1}{\beta_0}\right)\right] \mu_1^2 \tag{1.114}$$

[74] This is *not* true if γ and δ are scale dependent. In that case, only the first coefficient will be scheme independent.

[75] It may seem as though we have just obtained something for nothing. However, this manipulation is not a true simplification; it only plays one on TV. If we wish to take this scheme seriously, we must use it consistently in all of our calculations. One way to do this is to calculate an amplitude in a known scheme, say $\overline{\text{MS}}$, and re-express the amplitude in terms of the new coupling via (1.107). The constants γ and δ are obtained from the requirement (1.111) in conjunction with (1.110). Even after all of this manipulation, our amplitude will still be correct only up to the order at which it was calculated. On the other hand, this scheme is certainly not without its merits. The simple evolution of α_s is offers may indeed be worth the effort.

[76] Although the latter region is not physical in the sense that it is not asymptotically free, it certainly does exhibit the kind of behavior we expect for a theory with well-defined infrared behavior. Unless our coupling really does diverge in the infrared, the β-function must become positive as α_s becomes large. This, along with the physical requirement of Lipshitz continuity, implies the existence of an infrared fixed point. These topics are discussed to some extent in [42].

to obtain a sensible value for $\alpha_s(\mu_2^2)$.[77] The actual value of this bound on μ_2^2 depends on the number of active flavors and the value of μ_1^2,[78] but its existence implies that we must be more careful when attempting to work beyond perturbation theory. Despite appearances, our expressions are really only valid in the region of small α_s.

One can obtain β_1 from a two-loop calculation of any vertex in the theory. Such a calculation is extremely complicated, but certain general aspects of it can give us insight into the behavior of higher order effects. To begin with, assume

$$Z_g = \sum_\ell z_\ell \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\ell\epsilon/2} \left(\frac{\alpha_s(\mu^2)}{4\pi}\right)^\ell \quad . \tag{1.115}$$

We emphasize here that it is the *renormalized* coupling which appears in this expansion. Keeping all terms up to order α_s^3, we write

$$\begin{aligned} \beta(\alpha_s) &= \alpha_s \left[z_1 \epsilon \frac{\alpha_s}{4\pi} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} - 2 z_1 \frac{\beta}{4\pi} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \right. \\ &\qquad \left. + (2z_2 - z_1^2)\epsilon \left(\frac{\alpha_s}{4\pi}\right)^2 \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-2\epsilon/2} \right] \\ &= \alpha_s \left[z_1 \epsilon \frac{\alpha_s}{4\pi} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} + (2z_2 - 3z_1^2)\epsilon \left(\frac{\alpha_s}{4\pi}\right)^2 \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-2\epsilon/2} \right] \quad . \end{aligned} \tag{1.116}$$

Evidently, the coefficient we are after is

$$\beta_1 = \frac{3z_1^2 - 2z_2}{(4\pi)^2}\epsilon \quad . \tag{1.117}$$

Since z_1 is divergent like $1/\epsilon$, it looks as though the second coefficient of the β-function diverges in the same manner. This is unacceptable. The renormalizability of our theory guarantees a finite renormalized coupling. If g is finite, it does not have divergent variation with μ^2. The only way out of this situation is to require a complete cancelation of z_1^2, i.e. if $z_2 = a/\epsilon^2 + b/\epsilon$ then $a = 3(\epsilon z_1)^2/2$. This is in fact what happens. The fact that g is finite implies that the double poles in z_2 are not independent of z_1. This behavior is expected since double poles come from regions of momentum space where two loop-momenta

[77] Qualitatively, one can identify the two disjoint regions at the point $\alpha_s = \beta_0/\beta_1$. The phase plane portrait of (1.100) then reveals this value as a special kind of infrared fixed point. This analysis does not survive the transition to quantitative reasoning, however, since increasing the scale then gives ambiguous results.

[78] A standard treatment from the Z mass, changing n_f as quark masses are crossed, gives a lower bound of approximately 0.4 GeV^2.

are simultaneously large. Renormalizability assures us that all divergences obtained when one loop momentum gets large are taken care of by the one-loop counterterms, so these double poles are naturally related to products of the one-loop results. This is a general aspect of renormalizable quantum field theories - all higher-order ultraviolet divergences in an ℓ-loop diagram are related to the divergences in $n < \ell$-loop diagrams. Only the $1/\epsilon$ pole in a diagram gives new information. Stated compactly, all true ℓ-loop effects are given by the residue of the ℓ-loop amplitude at $\epsilon = 0$. This is a nice result, but difficult to exploit in practice. One usually needs to compute the entire divergence structure of a graph to extract the $1/\epsilon$ pole. However, it can be an extremely useful check of a calculation.

The result for β_1 can be found in the literature [43]. It is

$$\begin{aligned} \beta_1 &= \frac{1}{(4\pi)^2}\frac{2}{3}\left[17C_A^2 - 2n_f T_F(3C_F + 5C_A)\right] \\ &= \frac{1}{(4\pi)^2}\left(102 - \frac{38}{3}n_f\right) \quad . \end{aligned} \tag{1.118}$$

Since $\beta_1 > 0$ for $n_f < 9$, this coefficient only serves to hasten the asymptotic vanishing of the coupling.

Looking at the diagrams we used to arrive at this result, we see that the β-function for QED can be obtained by taking $C_A = 0$ and $T_F = C_F = 1$. In this theory, the β-function is *positive* in perturbation theory. We can understand this in terms of *screening* of electric charge. A charge sitting in the vacuum polarizes the virtual fluctuations. Since QED causes opposite charges to attract, this polarization results in a screening of the true electric charge. As we probe the charge at higher and higher energy, we effectively penetrate some of this shielding and get a look at the true charge. Hence, we would expect the QED coupling to increase with scale. The analogue of this argument in QCD results in anti-screening since the gluons themselves participate in the vacuum polarization. The effect of gluons is opposite that of quarks, so acts to increase the charge experienced at great distances. Although this argument and its extensions cannot rigorously prove that colored objects are confined, they can certainly go a long way towards giving us a qualitative understanding of this phenomenon.

1.6 Renormalization in the Axial Gauge - The Price of a Ghost-Free Theory

Now that we understand the need for renormalization in quantum field theories, we would like to see how it plays out in another gauge. As introduced in Section 1.2, the axial gauge seems rather simple. There are no ghosts to worry about and the freedom to choose n can in principle allow us to simplify calculations a great deal. The only apparent complication is the rather lengthy gluon propagator, Eq.(1.47).

This gauge belongs to a set of 'physical gauges' since contraction of the gauge propagator with its momentum gives

$$q^\mu D_{\mu\nu} = i\left[\frac{n_\nu}{n\cdot q} - \frac{q_\nu}{(n\cdot q)^2}\left(n^2+\xi q^2\right)\right] \quad , \tag{1.119}$$

which does not contain a propagator pole at $q^2 = 0$. Hence the unphysical degree of freedom associated with the longitudinal polarization state does not propagate in these gauges.[79] This is the reason there are no ghosts in axial gauges - the degree of freedom they live to cancel does not propagate here. It seems that axial gauge, rather than covariant, is the natural choice for gauge theories from this reasoning.

Once we begin to look at renormalization, however, this gauge changes its tune. The new tune is one of violent attacks on Lorentz invariance and unregulated divergences. To study these effects, let us consider a simpler situation than the general case. Apparently, taking $n^2 = \xi = 0$ simplifies the gluon propagator immensely. Since we intend to choose $\vec{n}$ as a special vector in our problem, we take it collinear to all external momenta in our problem. Note that this implies that our use of axial gauge requires collinear kinematics.[80] This space of collinear vectors is spanned by the basis $\{n^\mu = (1,0,0,-1)/2\Lambda, p^\mu = (1,0,0,1)\Lambda\}$, where Λ is an arbitrary scale reflecting the residual boost invariance along the 3-direction.[81] Our

[79] This is *not* the case in covariant gauges. There, contraction gives $-i\xi q_\nu/q^2$. Hence in covariant gauges unphysical degrees of freedom propagate unless one takes $\xi = 0$ (Landau's gauge).

[80] One can, of course, use the axial gauge in situations which are not collinear. However, the integrals one is faced with in these situations are a great deal more complicated than those we will encounter. The choice of axial gauge is usually made expressly for the reason that the kinematics of a process favor one direction over all others. Otherwise, there is no advantage to its use.

[81] This invariance is really an illusion since any specific choice for n fixes Λ.

collinear kinematics requirement is that all external momenta can be expressed as a linear combination of n and p, i.e. all particles move in the 3-direction.[82] Note that we have chosen our basis vectors such that $p^2 = n^2 = 0$ and $n \cdot p = 1$.

The goal of this section is to obtain the QCD β-function in the axial gauge. Along the way, we will see some major differences between this gauge choice and the covariant choice. As with the covariant gauges, our first task is to calculate the quark self-energy. For simplicity, we begin with massless quarks.[83] On general grounds, we can decompose the 1PI contribution as[84]

$$\Sigma(q) = A \not{q} + B\frac{q^2}{2n \cdot q} \not{n} \quad . \tag{1.120}$$

The appearance of $\not{n}$ is a direct consequence of the explicit Lorentz symmetry breaking induced by our axial gauge choice. This also leads to the fact that A and B are functions of q^2 *and* $n \cdot q$. Our choice $n^2 = 0$ suppresses their dependence on n^2. The factor $q^2/2n \cdot q$ multiplying B causes A and B to have the same dimensionality (since n^μ is like an inverse mass) and will prove convenient in the future.

Iterating the 1PI result to all orders to obtain the full 1PR propagator as in (1.55) gives

$$\frac{i}{\not{q}} \sum_n \left[(-i\Sigma) \frac{i}{\not{q}} \right]^n = \frac{i}{\not{q}} \left[1 - A - \frac{1}{2n \cdot q} B \not{n} \not{q} \right]^{-1} \quad . \tag{1.121}$$

One look at this expression tells us that things are not as simple as they were in the covariant gauges. We cannot renormalize the quark fields with a simple constant. The form of this full propagator implies that our renormalized fields must be related to the unrenormalized ones through a rotation in Dirac space; there is no other way to remove the $\not{n}$. Rewriting (1.121) as

$$\frac{1}{1-A} \left[1 + \frac{1}{2n \cdot q} \left(\sqrt{1 - B(1-A)^{-1}} - 1 \right) \not{q} \not{n} \right]^{-1}$$

[82] This is not as restrictive as it may seem. In two-particle systems, we can always transform into a frame in which the momenta are collinear. In this frame, we are free to quantize the theory and introduce n. However, once we have introduced n we are no longer free to change frames. This is a consequence of the axial gauges' violence to Lorentz symmetry.

[83] Since we are concerned here only with the ultraviolet behavior of the theory, this can be done without loss of generality. We will briefly consider massive quarks at the end of this section.

[84] We take the quark momentum as q here to avoid confusion with the basis vector p.

$$\times \frac{i}{\not{q}} \times \left[1 + \frac{1}{2n\cdot q}\left(\sqrt{1 - B(1-A)^{-1}} - 1\right) \not{n}\,\not{q}\right]^{-1} , \tag{1.122}$$

it is obvious that the renormalized fields are defined via

$$\psi = \sqrt{1-A}\left[1 + \frac{1}{2n\cdot q}\left(\sqrt{1 - B(1-A)^{-1}} - 1\right) \not{q}\,\not{n}\right]\psi_b \ . \tag{1.123}$$

This transformation is certainly not a constant multiplicative factor. Its nontrivial dependence on $n\cdot q$ implies that the manipulations used in Section 1.4 to define renormalized perturbation theory will lead to an infinite series of local operators, each of which renormalizes in its own special way. Hence our theory appears unrenormalizable.

What happened? In covariant gauges, QCD is a perfectly reasonable renormalizable field theory. Since the gauge choice we make is in principle arbitrary, there should be no real difference between the two choices; we certainly should not have the situation that QCD is renormalizable with one choice but not the other! However, we must be much more careful when discussing renormalizability. Taken by itself with no gauge-fixing and no regularization procedure, QCD is a nonsensical field theory. It is plagued by infinities of many different kinds and presents no way for us to extract sensible predictions from its amplitudes. Only after we decide on a regularization and gauge-fixing procedure will QCD mean something. For this reason, we certainly cannot expect QCD to be renormalizable for any and every choice of ultraviolet cut-off and gauge-fixing function. Renormalizability is as much a property of these choices as it is of the lagrangian. As mentioned above, QCD is not renormalizable even in covariant gauges if the ultraviolet cut-off breaks Lorentz invariance. In the present case, this symmetry is broken explicitly by the gauge-fixing procedure. However, since here it is only the gauge choice that breaks the symmetry, we can handle these difficulties and define a sensible quantum field theory in the axial gauge.[85]

To see how this is done, recall that we are not forced to use renormalized perturbation theory. In Section 1.4 we saw that we could also use unrenormalized perturbation theory. The results are the same as long as we express the final result in terms of renormalized quantities. In particular, all *external* fields in the amplitude must be renormalized. Writing the full quark

[85] This is valid as long as we choose an ultraviolet regulator which does not further mutilate either Lorentz symmetry or gauge invariance. We will employ dimensional regularization for this purpose.

propagator as

$$V^{-1}\frac{i}{\not q}\gamma^0\left(V^{-1}\right)^\dagger\gamma^0 \quad , \tag{1.124}$$

with

$$V \equiv \sqrt{1-A}\left[1+\frac{1}{2n\cdot q}\left(\sqrt{1-B(1-A)^{-1}}-1\right)\not q\not n\right] \quad , \tag{1.125}$$

we see that a properly normalized amputated Greens function is obtained by simply removing $V^{-1}i/\not q$ from the full propagator of all outgoing quark lines and $i/\not q\ \gamma^0(V^{-1})^\dagger\gamma^0$ from the full propagator of all incoming quark lines. All remaining divergences[86] are attributed to the operators themselves.

Now that we understand the basic idea behind renormalization in the axial gauge, let us calculate the all-important quantity V. Since A and B start at $\mathcal{O}(\alpha_s)$, a one-loop calculation sees only their leading powers in V :

$$V \sim \left(1-\frac{1}{2}A-\frac{1}{4n\cdot q}B\not q\not n\right) \quad . \tag{1.126}$$

The calculation of A and B is complicated by the presence of $1/k\cdot n$ in the gluon propagator. This term renders the integration techniques of Appendix A.3 impotent since it is not quadratic in loop momenta.[87] Alternatively, using our special collinear kinematics, we can decompose the d-dimensional integration as[88]

$$d^dk = d(n\cdot k)d(p\cdot k)d^{d-2}k_\perp \quad . \tag{1.128}$$

Since the denominator contains two powers of $p\cdot k$, this part of the integration may be done by contour[89] if no powers of $p\cdot k$ appear in the numerator. Powers that do appear may be

[86] after a similar manipulation is performed for all external gluon lines, of course

[87] A variation of the Feynman parameter approach in Appendix A.3 based on the fact that

$$\frac{1}{AB}=\int_0^\infty\frac{dz}{[A+zB]^2} \quad . \tag{1.127}$$

can be employed in this situation. Obviously, generalizations of this relation can be used to combine any number of normal propagators and 'light-cone' denominators $1/k\cdot n$. However, the introduction of a new Feynman parameter here leads to complications we would rather avoid. The techniques introduced below are much better suited for our purpose.

[88] Our choice of p and n gives no change in measure.

[89] The application of contour integration in this context is spelled out more clearly in Sections 2.2 and 3.3.

removed via the replacement

$$2p\cdot k = \frac{n\cdot(q-k)}{(n\cdot q)^2}q^2 \quad , \tag{1.129}$$

which is valid in the numerator.[90] In this manner, we obtain the result

$$\begin{aligned}\Sigma \;\; = \;\; & -\frac{\alpha_s C_F}{2\pi}\Gamma\left(\frac{\epsilon}{2}\right)\left(\frac{-q^2}{4\pi\mu_0^2}\right)^{-\epsilon/2}\int_0^x \frac{dy}{x}\left[\frac{y}{x}\left(1-\frac{y}{x}\right)\right]^{-\epsilon/2} \\ & \times\left\{\left(1-\frac{\epsilon}{2}\right)\left(1-\frac{y}{x}\right)\not{q}+\frac{2y}{x-y}\frac{q^2}{2x}\not{n}\right\} \quad ,\end{aligned} \tag{1.131}$$

where $x \equiv n\cdot q$. The ultraviolet divergence is contained in the factor $\Gamma(\epsilon/2)$ multiplying our expression.

In analogy with Section 1.4, we would like to include only the ultraviolet divergent part of Σ in the definition of V. This procedure leads to an extra divergence of the form[91]

$$\int_0^x \frac{dy}{x}\frac{2x}{x-y} \tag{1.132}$$

in the second term. Tracing it back through the calculation, we see that our new divergence comes directly from the factor $1/k\cdot n$ in the gluon propagator. This behavior is reminiscent of the divergence we encountered in Section 1.4 when we were working in the onshell scheme. There, the divergence appeared because of our choice of external momenta. Once we decided that we did not wish to renormalize in the onshell scheme, it no longer posed a problem for us. On the other hand, this divergence is present regardless of which external states we choose. Deforming n^2 away from zero, we see that instead of a divergence we obtain terms like $\log^2(n^2q^2/(q\cdot n)^2)$ in the *finite* part of our amplitude. The part of our amplitude that generates these light-cone singularities does not even *contribute* to the ultraviolet divergence for n^2 finite.

---

[90] In deriving this result, I have used

$$\int d(n\cdot k)d(p\cdot k)f(n\cdot q, n\cdot k)\frac{d^{\,d-2}k_\perp}{(2\pi)^d}\frac{1}{k^2} = \int d(n\cdot k)d(p\cdot k)f(n\cdot q, n\cdot k)\frac{d^{\,d-2}k_\perp}{(2\pi)^d}\frac{1}{(k-q)^2} = 0 \tag{1.130}$$

by absence of scale. This is true regardless of the form of f since both $n\cdot q$ and $n\cdot k$ are dimensionless. The validity of this replacement also rests on the fact that q has no transverse component. This is one place where complications due to general kinematics rear their ugly heads.

[91] For finite ϵ, this integral is quite finite. Looking at Eq.(1.131), we see that the divergence appears as $\Gamma(-\epsilon/2)$. However, consistency requires that only *ultraviolet* divergences be included in V. A well-defined separation between infrared and ultraviolet physics can be obtained only if we use different techniques to regularize the two types of divergences.

To gain insight into the meaning of these divergences, it is necessary to take a closer look at the calculation. A characteristic integral that leads to divergences of this sort is

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(n \cdot k - \omega)\, k^2\, (q-k)^2} \ . \tag{1.133}$$

A simple power-counting argument implies immediately that this integral is not ultraviolet divergent. However, performing the k^- integral by contour, we immediately arrive at the opposite conclusion. This fact is extremely disconcerting. Renormalization proofs rely heavily on power-counting, so the ultraviolet divergence contained in this one integral has the power to destroy the renormalizability of our theory. Actually performing this integral away from the singular points in the complex plane of ω reveals that it not only diverges, but produces a divergence which is dependent on *logarithms* of the external momenta. Since $\log(x)$ is not analytic at $x = 0$, these divergences cannot even be absorbed by local operators. If we intend to take this theory seriously, our lagrangian requires counterterm operators of the form $\overline{\psi}\ \log(in \cdot \partial)\, \psi$. Obviously, this is quite unacceptable.

In order to understand why the integral (1.133) does not follow the rules of power-counting, we must remember how those rules are derived. Since the square of a Euclidean vector is positive-definite, power-counting arguments are always valid in Euclidean space. In order to use Euclidean arguments on a Minkowskii integral, our integrand must allow us to perform a rotation through $\pi/2$ radians in the complex energy plane. The causal prescription for the physical propagator poles introduced in Section 1.2 satisfies this requirement.[92] We have not explicitly chosen a prescription for the pole in the gluon propagator, but our manipulations imply that our prescription does not depend on any of the other coordinates. Since $n \cdot k$ is linear in k^0, our poles occur either always above or always below the axis. No prescription of this form admits Wick rotations, so we cannot apply power-counting techniques.

As mentioned in Section 1.2, there is a causal prescription for our new poles. Since we intend to perform our integrals in light-cone coordinates, we prefer the Mandelstam-Liebbrandt (ML) prescription [39]

$$\frac{1}{n \cdot k} \to \frac{p \cdot k}{(p \cdot k)(n \cdot k) + i\varepsilon} = \frac{2p \cdot k}{k^2 + k_\perp^2 + i\varepsilon} \tag{1.134}$$

[92] This prescription forces negative-energy poles to be above the real axis and positive-energy poles to be below. Hence the axis can be rotated freely in the ultraviolet region without encountering any poles.

over the one involving $\vec{k}\cdot\vec{n}$. The form of this prescription makes our earlier mathematical indiscretion apparant. We were considering contour integration in the k^- plane, while the poles in the k^+ plane are the ones that truly dictate the analytic behavior of our integral. The fact that we ignored the analytic structure of this pole before *induced* the non-causal principle value (PV) prescription[93]

$$\frac{1}{n\cdot k}\to\lim_{\delta\to 0}\frac{1}{2}\left(\frac{1}{n\cdot k+i\delta}+\frac{1}{n\cdot k-i\delta}\right)=\lim_{\delta\to 0}\frac{n\cdot k}{(n\cdot k)^2+\delta^2}\quad . \tag{1.135}$$

While this prescription certainly regularizes the light-cone divergence, it is not at all obvious that it will lead to a renormalizable quantum field theory.

To illustrate the difference between these two prescriptions, it is instructive to see how our calculation works in each. Since we already have an expression for the quark self-energy with the principle value prescription, we begin with this choice. Defining

$$I_0\equiv\lim_{\delta\to 0}\int_0^1\frac{xdx}{x^2+\delta^2}\sim\int_0^1\frac{dx}{x}\quad , \tag{1.136}$$

we can extract the ultraviolet divergent parts of Eq.(1.131) :

$$\begin{aligned}
[A(x)]_{\rm UV} &= -\frac{\alpha_s C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\\
[B(x)]_{\rm UV} &= -\frac{\alpha_s C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(4I_0+4\log x-4)\quad .
\end{aligned} \tag{1.137}$$

As promised, a dependence on the log of $n\cdot q$ has been generated.

In the gluon sector, the situation is analogous. The fact that the axial gauge is 'physical' implies that the gluon self-energy must satisfy current conservation without any help from ghosts. The most general form for the gluon 1PI self-energy with our choice of kinematics is

$$\begin{aligned}
\Pi^{\mu\nu}(q) &= \Pi_1\left(q^\mu q^\nu-q^2 g^{\mu\nu}\right)\\
&\quad+\Pi_2\left(q-\frac{q^2}{q\cdot n}n\right)^\mu\left(q-\frac{q^2}{q\cdot n}n\right)^\nu\quad .
\end{aligned} \tag{1.138}$$

The full gluon propagator,

$$D^{\mu\nu}_{a\,b}(q)=\frac{\delta^{\,a\,b}}{1-\Pi_1}\qquad\left[\delta^\mu_\alpha+\left(\sqrt{1+\frac{\Pi_2}{1-\Pi_1-\Pi_2}}-1\right)\frac{1}{n\cdot q}n^\mu q_\alpha\right]$$

[93] I should mention here that we intend to use this prescription in the most naieve way possible. The subtleties of defining products and powers of this distribution are considered in [44].

$$\times \frac{id^{\alpha\beta}(q)}{q^2} \times \left[\delta^\nu_\beta + \left(\sqrt{1+\frac{\Pi_2}{1-\Pi_1-\Pi_2}}-1\right)\frac{1}{n\cdot q}q_\beta n^\nu\right] \quad , \tag{1.139}$$

where $d^{\mu\nu}(q) \equiv -g^{\mu\nu} + (q^\mu n^\nu + q^\nu n^\mu)/q\cdot n$, is obtained by summing the 1PR contributions. Inverting the matrix on either side, we see that the renormalized gluon field is defined via

$$\mathcal{A}^\mu = \sqrt{1-\Pi_1}\left[\delta^\mu_\alpha + \left(\sqrt{1-\frac{\Pi_2}{1-\Pi_1}}-1\right)\frac{1}{n\cdot q}n^\mu q_\alpha\right]\mathcal{A}^\alpha_b \quad . \tag{1.140}$$

At the one-loop level, only two diagrams contribute to Π. The quark loop is identical to its covariant gauge counterpart and does not depend on our prescription for the light-cone singularity. The other diagram, the gluon loop, can be evaluated using the technique outlined above. The full result contains the UV divergence

$$\begin{aligned}
[\Pi_1(x)]_{\rm UV} &= \frac{\alpha_s C_A}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(\frac{11}{3}-\frac{4n_f T_F}{3C_A}-4I_0-4\log x\right)\\
[\Pi_2(x)]_{\rm UV} &= \frac{\alpha_s C_A}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(-4+4I_0+4\log x) \quad .
\end{aligned} \tag{1.141}$$

The three-point function that defines the renormalized coupling receives one-loop corrections from the same diagrams we considered in covariant gauge. The integrals are a little more complicated than in the self-energies, but can be taken readily if we are interested only in the ultraviolet divergent part. Let us consider a longitudinally polarized gluon. These amplitudes are obtained by contracting the gluon polarization index with n.[94] The quark correction (Figure 1.7a) has the divergence

$$(-ig_b t^a \not{n})\frac{\alpha_s}{4\pi}\left(C_F-\frac{C_A}{2}\right)\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(4I_0+2\log x_i+2\log x_f-3) \quad , \tag{1.142}$$

while the gluon correction (Figure 1.7b) contains

$$(-ig_b t^a \not{n})\frac{\alpha_s C_A}{8\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(1+\frac{x_i+x_f}{x_f}\log\frac{x_i}{x_g}+\frac{x_i+x_f}{x_i}\log\frac{x_f}{x_g}\right) \quad . \tag{1.143}$$

Here, x_i, x_f, and x_g are the light-cone momenta $n\cdot q$ for the initial quark, final quark, and gluon, respectively.

[94] For example, the leading three-point function for polarization α is $-igt^a\gamma^\alpha$. For longitudinal polarization, we write $-igt^a\not{n}$.

The self-energy corrections are performed by tacking the full propagators (1.122) and (1.139) on to the leading order diagram, and removing the middle propagator and the external Z_F-matrix to renormalize the external states. The quark,

$$(-ig_b t^a \not{n}) \frac{\alpha_s C_F}{4\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} (3 - 4I_0 - 2\log x_i - 2\log x_f) \quad , \tag{1.144}$$

and gluon,

$$(-ig_b t^a \not{n}) \frac{\alpha_s C_A}{8\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \left(\frac{11}{3} - 4 - \frac{4n_f T_F}{3C_A} \right) \quad , \tag{1.145}$$

combine with the vertex corrections to give the full one-loop vertex function,

$$\begin{aligned} &(-ig_b t^a \not{n}) \left[1 + \frac{\alpha_s}{4\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \left(\frac{11}{6} C_A - \frac{2n_f T_F}{3} \right.\right. \\ &\left.\left. + \frac{C_A}{2} \left(\frac{x_i - x_f}{x_f} \log x_i + \frac{x_f - x_i}{x_i} \log x_f - \frac{(x_i + x_f)^2}{x_i x_f} \log x_g - 4I_0 \right) \right) \right] \quad . \end{aligned} \tag{1.146}$$

If we simply ignore the dependence on x_i, x_f, and x_g for the moment, this expression implies that the effective coupling at scale μ is given by[95]

$$g(\mu^2) \equiv g_b \left[1 + \frac{\alpha_s}{4\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \left(\frac{11}{6} C_A - \frac{2n_f T_F}{3} \right) \right] \quad , \tag{1.147}$$

which gives the β-function found in the last section.

The fact that the logarithms do not cancel implies that this gauge choice with the non-causal prescription for the light-cone singularity is truly not renormalizable, as predicted.[96] However, the specific form (1.135) of the prescription is not actually used until the integrals are

[95] Note the difference between renormalized and unrenormalized perturbation theory. In renormalized perturbation theory, we would have determined Z_g from a subtraction of ultraviolet divergences. Here, we evaluate the full contribution without a thought given to renormalization (except for the external states). The effective coupling is determined by the requirement that our matrix element be finite when expressed in terms of it.

[96] In the literature, one can find a calculation of the β-function in this gauge which does not contain these difficulties [45]. Expressions for the momentum-dependent 'Z-factors' from various diagrams are given, with the result that the unsightly terms cancel in the renormalized coupling. Their calculation makes no mention of the gluon polarization, and the wavefunction renormalization factors they employ are appropriate for the product $\overline{\psi}\ \not{n}\mathcal{A}_\perp\psi$. Since this combination does not appear in the lagrangian, it is unclear how to interpret their result. In addition, several papers present calculations of the QCD splitting functions using PV [46]. This prescription closely resembles that used for eikonal propagation introduced in Section 2.3, so it may indeed be appropriate for this purpose. However, if a renormalization procedure that produces sensible results in *all* cases exists for this prescription, it is certainly not straightforward.

performed. If we choose instead to allow the light-cone denominators to act only *algebraically*, and simply ignore any terms of the form $c/k\cdot n$, with c independent of k, the embarrassing terms never appear. Unfortunately, since the integrals in each diagram contain different limits, this 'prescription' is hard to quantify mathematically. In addition, it is entirely unclear how this procedure would be implemented at higher orders.

The situation for transversely polarized gluons is analogous. Here, even the Dirac structure is nontrivial. The quark

$$\left(-ig_b t^a \gamma^\perp\right) \frac{\alpha_s}{4\pi}\left(C_F - \frac{C_A}{2}\right)\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(-1 - \frac{x_f}{x_g}\log\frac{x_i}{x_f}\ \not{n}\ \not{p} + \frac{x_i}{x_g}\log\frac{x_f}{x_i}\ \not{p}\ \not{n}\right) \tag{1.148}$$

and gluon

$$\left(-ig_b t^a \gamma^\perp\right) \frac{\alpha_s C_A}{8\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(-1 + 4I_0 + 4\log x_g - \log\frac{x_i}{x_f}\ \not{n}\ \not{p} - \log\frac{x_f}{x_i}\ \not{p}\ \not{n}\right) \tag{1.149}$$

vertex diagrams combine with the wavefunction renormalization factors to generate the full vertex

$$\begin{aligned}(-igt^a\gamma^\perp)\frac{\alpha_s}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \quad & \left[\frac{11}{6}C_A - \frac{2}{3}n_f T_F + \frac{C_A}{2}\left(\frac{1}{x_g}\log\frac{x_i}{x_f}\ \not{n}\ \not{p}_i - \frac{1}{x_g}\log\frac{x_f}{x_i}\ \not{p}_f\ \not{n}\right)\right. \\ & + C_F\left(\frac{1}{x_g}\log x_f\ \not{n}\ \not{p}_i - \frac{1}{x_g}\log x_i\ \not{p}_f\ \not{n}\right) \\ & \left. - C_F\left((I_0 - \frac{x_i}{x_g}\log x_f)\ \not{p}\ \not{n} + (I_0 + \frac{x_f}{x_i}\log x_i)\ \not{n}\ \not{p}\right)\right]\ .\end{aligned} \tag{1.150}$$

Once again, we see that the correct one-loop β-function is obtained if the offensive contributions are simply ignored.

Before moving on to the Mandelstam-Liebbrandt prescription, it is instructive for us to study the way in which the offensive terms pollute our final results. In the longitudinal direction, (1.146) shows that the unsightly logarithms confine themselves to the C_A-part of the coupling. Tacking external states[97] onto (1.150) causes all but the C_F-part of these terms to vanish. This fact is intimately related to the gauge choice we have made and its implications on the physical states of our theory.

[97] This procedure is somewhat delicate since truly onshell external states will be annihilated by $\not{n}\ \not{p}$ whether or not it is accompanied by x_i. The ambiguity comes directly from our choice of collinear kinematics. To resolve it, we could choose to add a small transverse momentum component to each of our external legs; since this pole prescription is flawed in any case, we will not bother.

Quantizing our theory at equal values of $n \cdot x \equiv x^+$,[98] rather than at equal times, we see that the QCD equations of motion[99] in the gauge $n \cdot \mathcal{A} = 0$ reveal a severe pathology with our formalism : they do *not* dictate the x^+ ('time') evolution of either $\mathcal{A}^-$ or $\psi_- \equiv 1/2\ \not{n}\not{p}\psi$. This means that these field operators do *not* propagate freely in this gauge. Only the transverse gluon field components and the physical 'good' quark field $\psi_+ \equiv 1/2\ \not{p}\not{n}\psi$ can be directly interpreted in the perturbative theory we have derived. The other components must be re-expressed in terms of the freely-propagating ones through their equations of motion before they can be considered in our formalism. In light of this observation, it is not surprising that the longitudinal contribution, which is associated with $\mathcal{A}^-$, is polluted by C_A-type gluon effects and the transverse contribution, which contains ψ_-, is polluted by C_F-type quark effects. In some sense, the fact that these field components do not propagate freely makes our renormalization of the external states insufficient to remove all associated divergences. Notice that the residual divergence structures depend on *all three* momenta, so the renormalization of these 'bad' field components actually depends on the diagram they are embedded in, further evidence of their composite nature.

Now that we have seen the consequences of a noncausal prescription, let us see what happens when we use one that is physically motivated. The most important practical difference between the Mandelstam-Liebbrandt prescription (1.134) and the PV prescription is that ML introduces another vector to the problem. This effectively adds a new scale and invalidates arguments such as that in Footnote 90. However, this added complication is more than compensated by the fact that ML preserves power counting.[100] Integrals of the form

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{n \cdot k (k-q)^2} \tag{1.152}$$

contribute to ultraviolet divergences, but those of the form (1.133) do not. This leads to a

[98] See Appendix D.1.

[99] See Appendix D.2

[100] I should mention here that the power-counting techniques of covariant calculations must be slightly modified to obtain the correct results. Each of our d-dimensions can conceivably generate an ultraviolet divergence, so we must power count each dimension separately. The integral

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(q-k)^2} \frac{1}{(n \cdot k)(n \cdot (k-\ell))(n \cdot (q-k))} \tag{1.151}$$

is divergent even in spite of the fact that its overall degree of divergence is negative.

tremendous simplification. In particular, the integrals which generate the offensive contributions in the PV prescription are automatically thrown away by ML.

The calculation is straightforward. One simplifies the expression of each diagram, taking care to retain only those terms which contribute to the UV divergence, and performs the k^+ integration by contour. The transverse integrations are now trivial, and one is left with a simple one-dimensional integral for k^-. With the same conventions as above, the self-energy diagrams contribute

$$\begin{aligned}
[A(x)]_{\rm UV} &= -\frac{\alpha_s C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(1) \\
[B(x)]_{\rm UV} &= -\frac{\alpha_s C_F}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(-4) \quad ; \qquad (1.153) \\
[\Pi_1(x)]_{\rm UV} &= \frac{\alpha_s C_A}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(\frac{11}{3}-\frac{4n_f T_F}{3C_A}\right) \\
[\Pi_2(x)]_{\rm UV} &= \frac{\alpha_s C_A}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}(-4) \quad , \qquad (1.154)
\end{aligned}$$

which is certainly quite a bit cleaner than the PV results! It is interesting to note that the two prescriptions agree if we ignore the logarithm-type terms. This behavior is also seen in the three-point diagrams, Figs. 1.7(a) and (b) in the longitudinal,

$$(-ig_b t^a \not{n})\,\frac{\alpha_s}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(C_F-\frac{C_A}{2}\right)(-3) \quad , \qquad (1.155)$$

$$(-ig_b t^a \not{n})\,\frac{\alpha_s}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(\qquad\frac{C_A}{2}\right)(\ \ 1) \quad , \qquad (1.156)$$

and transverse,

$$\left(-ig_b t^a \gamma^\perp\right)\frac{\alpha_s}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(C_F-\frac{C_A}{2}\right)(-1) \quad , \qquad (1.157)$$

$$\left(-ig_b t^a \gamma^\perp\right)\frac{\alpha_s}{4\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left(\qquad\frac{C_A}{2}\right)(-1) \quad , \qquad (1.158)$$

sectors, respectively. From these results, we easily arrive at the correct β-function, (1.102), pollutant-free.

In order to understand why ML was able to circumvent the problems associated with renormalizing the external composite 'bad' fields and generate the correct β-function, we appeal to some of the arguments in the beginning of this section. Since the violence to Lorentz symmetry (which leads directly to the fact that different components of the fields behave in different

ways) occurs only in the choice of gauge, it is unphysical. Its effects will cancel in the calculation of gauge-invariant quantities, giving the same result for all well-defined gauge choices. As we have seen, the contribution to the three-point functionfrom each diagram depends on the gluon polarization. The PV prescription does not exhibit a cancelation of this dependence, leading to a directionally-dependent 'renormalized coupling' in direct violation of the gauge symmetry. In ML, the observed cancelation allows us to define a sensible, directionally-independent renormalized coupling. This is direct evidence that the causal prescription does not do further damage to our theory and invalidate our gauge principle. Here, the explicit breaking of Lorentz symmetry is indeed artificial and constrained to cancel in a sum over all contributions.

The ML prescription contains one loose end that should be taken care of before we conclude this section. When calculating the self-energy diagrams above, we implicitly assumed that they have the same form in both prescriptions. In light of (1.123) and (1.140), this assumption still leads to renormalization 'constants' which depend on the field momenta. Since we intend to take ML seriously, we cannot be satisfied with this behavior. Looking at these expressions again, we see that for the collinear kinematics we used the replacement $q^\mu/n\cdot q \to p^\mu$ is valid. As ML introduces p^μ *independently* of the external momenta, dependence on p^μ is perfectly acceptable.[101] To test the validity of this replacement unambiguously, it is necessary to consider transverse momenta. We do this explicitly in the quark sector.

Consider the complete 1PI quark self-energy in ML. Its most general form[102] is

$$\Sigma(q) = A\, \not{q} + B\, p\cdot q\, \not{n} + C\, n\cdot q\, \not{p} + D m_b \ . \tag{1.159}$$

For completeness, we work with massive quarks. Iterating this result as above, one obtains the full propagator

$$\frac{i}{\not{q} - m_b - \Sigma} \ . \tag{1.160}$$

We would like to decompose this into a form reminiscent of (1.124), but with the added requirement that the matrix V be independent of q. Working through this, we arrive at the

[101] Note that this is *not* the case for PV. There, dependence on the external momenta is unavoidable - even in the absence of the logarithms.

[102] Since QCD conserves parity and the self-energy is hermitian in the sense that $\Sigma^\dagger = \gamma^0\Sigma\gamma^0$, terms involving products of two or more γ-matrices either vanish or reduce to these structures.

full propagator

$$\frac{1-A-B}{(1-A)^2}\left[1-\frac{B}{2(1-A)}\;\not{p}\,\not{n}\right]^{-1}\frac{i}{\not{q}-m}\left[1-\frac{B}{2(1-A)}\;\not{n}\,\not{p}\right]^{-1}\quad , \tag{1.161}$$

where the renormalized mass is given by

$$m=\frac{1+D}{1-A}\,m_b\;\;, \tag{1.162}$$

along with the consistency relation

$$C(1-A-B)+B(1-A)=0\;\;. \tag{1.163}$$

This implies that our renormalized fields are given by

$$\psi=\frac{1-A}{\sqrt{1-A-B}}\left[1-\frac{B}{2(1-A)}\;\not{p}\,\not{n}\right]\psi_b\;\;. \tag{1.164}$$

If A, B, C, and D are all independent of q and satisfy (1.163), we are free to define proper renormalized fields in the spirit of Section 1.4 and arrive at a sensible, consistent theory. At one-loop order, we can calculate these quantities explicity; with the common overall factor

$$\frac{\alpha_s C_F}{4\pi}\,\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\quad , \tag{1.165}$$

we have

$$A = 1 \tag{1.166}$$

$$B = 2 \tag{1.167}$$

$$C = -2 \tag{1.168}$$

$$D = 2\;\;. \tag{1.169}$$

Note that (1.163) is satisfied to the order at which we work and the correct gauge-invariant dependence of $m(\mu)$ on μ,

$$\mu\frac{d}{d\mu}m(\mu)=-3\frac{\alpha_s C_F}{2\pi}\,m(\mu)\;\;, \tag{1.170}$$

is reproduced.

In the collinear limit, n^μ, k^μ, and p^μ are no longer linearly independent. Here, we can calculate only the combined quantities $A_0 \equiv A + C$ and $B_0 \equiv B - C$. Fortunately, our

consistency relation (1.163) allows us to extract A, B, and C from these two:

$$A = 1 - \sqrt{1 - (2A_0 + B_0) + A_0(A_0 + B_0)} \ , \tag{1.171}$$

$$B = \frac{1 - A_0 - B_0}{2 - A_0 - B_0} B_0 \tag{1.172}$$

$$C = -\frac{1 - A}{2 - A_0 - B_0} B_0 \ . \tag{1.173}$$

Using these expressions, we see that (1.123) is just a re-expression of (1.164) in terms of A_0 and B_0. In this way, we can calculate the correct renormalization constants using collinear kinematics.

In the gluon sector of the theory, things are similar. Here, since there is no mass, we have no need to calculate explicitly the analogues of A, B, and C. The collinear expressions Π_1 and Π_2 presented above are quite sufficient for all of our calculations.

It is interesting to note that the entire β-function is contained in the factor Π_1 of the gluon self-energy. In our calculation of the three-point function, all of the other contributions conspire to cancel each other. This is a general property of the axial gauges at one-loop order, and provides an explicit example of how changes in gauge can shift contributions from graph to graph without changing the final result.

In this section, we explored a non-covariant gauge choice. We have seen how subtle differences in prescription can make or break renormalizability in a gauge which explicitly violates Lorentz symmetry. If we had broken the spacetime symmetry more explicitly, summing all contributions would not have been enough to restore its former health. This is what happens when we regularize divergences by a momentum cut-off, for example. There, special counterterms must be added to the lagrangian to forcibly restore the symmetry order by order in perturbation theory.

The axial gauge's problems are not insurmountable. As we have shown, the theory possesses well-defined matrix elements which produce the correct behavior if implemented properly. Though some of its aspects are more involved than those of its covariant brother, this gauge is not without its charms. Among other things, it restores a certain degree of physical intuition to QCD. This gauge also represents tremendous simplifications in some applications, most notably those involving extremely high energy processes. For these reasons, it is often employed despite its difficulties.

Chapter 2

Deep-Inelastic Scattering

In the last chapter, we saw that asymptotic freedom assures us QCD perturbation theory is applicable to processes whose natural scales are large compared to the typical energy at which the coupling gets large. Unfortunately, confinement implies that *all* physical processes are inextricably linked to the low energy phenomena that lead to bound states. In order to calculate even the simplest of observables, we must first understand the complicated QCD bound states that appear in our experiments. Since perturbation theory cannot be applied to these states, the technology developed in the last chapter is useless for this purpose. In the absence of nonperturbative techniques for solving interacting quantum field theories, we must throw our hands in the air and give up the task of calculating *any* strong interaction cross-section from scratch.

Although we cannot calculate truly physical quantities completely in QCD, there is hope that we can use experimental results to learn about these nonperturbative quantities. First advanced by Feynman in 1969 [27], this idea is based on a separation of scales in certain processes. If the high- and low-energy contributions to a process can systematically be separated, one can calculate the high-energy physics in perturbation theory and use experimental data to *extract* the nonperturbative physics. These nonperturbative quantities can then be used not only to predict the outcome of other similar experiments, but also to teach us about the structure of QCD's bound states.

The purpose of this chapter is to describe the scale-separation process in the context of inclusive deep-inelastic scattering (DIS). The formalism developed here can be applied to a number of physical processes, and represents the main way QCD is used to predict the outcome

of experiments and give us insight into hadronic structure.

The process of DIS, along with the relevant structure functions and kinematics, is described in Section 2.1. It is here that we will discuss what is measured in the experiment and how the measurements are related to hadronic physics.

In Section 2.2, a qualitative argument is used to motivate the separation of scales. We use this argument to perform perturbative calculations of the hard scattering amplitude and find that it contains infrared divergences. These divergences signal soft nonperturbative physics that cannot be calculated reliably within perturbation theory.

Section 2.3 explains that these divergences represent contributions from nonperturbative distribution functions. We carefully define these distributions and show that our amplitudes are perfectly finite when expressed in terms of them. A measurement of the cross-section, coupled with knowledge of the perturbative coefficients, will now allow us to extract these nonperturbative quantities. Some of the structural information contained in these distributions is discussed, as well as certain consistency relations they are required to satisfy.

The analysis of Section 2.4 is devoted to infrared divergences and their relation to parton distribution functions. It allows us to show that the soft, nonperturbative contributions to inclusive DIS can be attributed exclusively to the parton distribution functions of Section 2.3 to *all* orders of QCD perturbation theory. Such a statement is called a *factorization theorem*. The tools we will use to derive this result were originally developed by G. Sterman [47], S. Libby [48], and J. Collins [49] using the results of L. Landau [50].

Since the perturbative physics in our process is related only to the microscopic degrees of freedom of QCD, quarks and gluons, all target dependence is isolated in the nonperturbative distribution functions. Section 2.5 shows how to use this fact to arrive at an operator relation between the nonlocal product of two electromagnetic currents and an infinite series of local partonic operators. Such local operator product expansions (OPE's) were first considered by Wilson in 1969 [29]. Since then, they have been used extensively to study the analytic structure of amplitudes and characterize scattering mechanisms. Using the results of Section 2.3, we derive the next-to-leading order coefficients of the OPE and discuss the interpretation of the local operators.

Section 2.6 contains a summary and some concluding remarks about inclusive DIS.

2.1 Kinematics

Deep-inelastic scattering (DIS) is a process in which a lepton is scattered off of a hadron at very high energy. The final state of this process will be a backscattered lepton and a complicated hadronic state which we denote as $|X\rangle$. We consider an electron scattering via the exchange of a highly virtual photon off of a proton, but the theoretical tools developed here also apply to weak exchanges and processes involving other hadronic states. At leading order in the electromagnetic coupling, the amplitude for this process is given by

$$\left\langle k's', X \left| \mathrm{T}\left((-i) \int d^4x \, J_e^\mu(x) A_\mu(x) (-i) \int d^4y \, J_q^\nu(y) A_\nu(y) \right) \right| ks, PS \right\rangle \ , \tag{2.1}$$

where I have denoted the electron's initial (final) momentum and spin as $k(k')$ and $s(s')$, respectively, and the proton's momentum and spin by P and S, respectively. The currents $J_e^\mu = e\,\overline{\psi}_e \gamma^\mu \psi_e$ and $J_q^\mu = \sum_f e_f \overline{\psi}_f \gamma^\mu \psi_f$ are the usual electron and quark electromagnetic currents, respectively. Recognizing that there are no photons in either external state[1] and that the two currents are fundamentally different allows us to separate the matrix elements and arrive finally at the amplitude[2]

$$\frac{i}{(k'-k)^2} \left\langle k's' \left| J_e^\mu(0) \right| ks \right\rangle \left\langle X \left| J_q^\nu(0) \right| PS \right\rangle g_{\mu\nu} (2\pi)^4 \delta(p_X + k' - P - k) \ , \tag{2.2}$$

where p_X represents the 4-momentum of the state $|X\rangle$ and I have employed Feynman's gauge ($\xi = 1$).

This is an extremely complicated object which is different for each possible final hadronic state. General processes such as this one are difficult to pin down theoretically. However, a great deal of information is contained in the *inclusive* cross-section,

$$\begin{aligned} \sigma \propto \quad & \frac{1}{q^4} \left\langle k's' \left| J_e^\alpha(0) \right| ks \right\rangle \left\langle ks \left| J_e^\mu(0) \right| k's' \right\rangle g_{\mu\nu} g_{\alpha\beta} \\ & \times \sum_X \left\langle PS \left| J_q^\nu(0) \right| X \right\rangle \left\langle X \left| J_q^\beta(0) \right| PS \right\rangle (2\pi)^4 \delta^{(4)}(P + q - p_X) \ , \end{aligned} \tag{2.3}$$

for DIS. Here, I have defined $q \equiv k - k'$. Such total cross-sections are much cleaner theoretically and make the relationship between microscopic and macroscopic degrees of freedom more obvious.

[1] at leading order in the electromagnetic coupling

[2] I have used the operator identity $\mathcal{O}(x) = e^{i\hat{P}\cdot x} \mathcal{O}(0) e^{-i\hat{P}\cdot x}$, where $\hat{P}$ is the momentum operator introduced in Appendix B.2.

Since the physics outside the brackets is quite well understood (it can be done within the framework of QED), we ignore it for the time being and focus on the hadronic object

$$\begin{aligned} W^{\mu\nu} &\equiv \frac{1}{4\pi}\sum_X \left\langle PS\left|J_q^\mu(0)\right|X\right\rangle\left\langle X\left|J_q^\nu(0)\right|PS\right\rangle(2\pi)^4\delta(P+q-p_X) \\ &= \frac{1}{4\pi}\int d^4z\, e^{iq\cdot z}\left\langle PS\left|J^\mu(z)J^\nu(0)\right|PS\right\rangle \quad . \end{aligned} \tag{2.4}$$

In deriving the final form I have used the fact that $\sum_X |X\rangle\langle X| = 1$ for any complete set of orthonormal states. Since $W^{\mu\nu}$ is concerned only with quark electromagnetic currents, I have dropped the subscript q. The factor $1/4\pi$ is a convention for future convenience.

Let us explore some of the properties of $W^{\mu\nu}$. Hermiticity of the currents implies that $(W^{\mu\nu})^* = W^{\nu\mu}$. $W^{\mu\nu}$ also conserves parity since all of our interactions and currents do. Furthermore, electromagnetic current conservation implies $q_\mu W^{\mu\nu} = q_\nu W^{\mu\nu} = 0$. These properties allow us to deduce the tensor structure[3]

$$\begin{aligned} W^{\mu\nu} = & \left(\frac{q^\mu q^\nu}{q^2} - g^{\mu\nu}\right)F_1(\nu,q^2) \\ & + \left(P-\frac{\nu}{q^2}q\right)^\mu\left(P-\frac{\nu}{q^2}q\right)^\nu\frac{1}{\nu}F_2(\nu,q^2) \\ & - i\epsilon^{\mu\nu\alpha\beta}q_\alpha S_\beta\frac{2M}{\nu}G_1(\nu,q^2) \\ & - i\epsilon^{\mu\nu\alpha\beta}q_\alpha\left(\nu S - q\cdot S\, P\right)_\beta\frac{2M}{\nu^2}G_2(\nu,q^2) \end{aligned} \tag{2.5}$$

for W, where $\nu \equiv q\cdot P$ and M is the proton mass. The totally antisymmetric Levi-Cevita tensor, $\epsilon^{\mu\nu\alpha\beta}$, is introduced with the convention $\epsilon^{0123} = +1$ in Appendix B.1. The real scalar functions $F_{1,2}$ and $G_{1,2}$ are called *structure functions* of the proton. Our task is to relate these functions to the nonperturbative wavefunction of the proton. Our three external vectors define six invariants. Since $P^2 = M^2$, $S^2 = -1/4$, and $S\cdot P$=0, we are left with two scalars and one pseudoscalar. The pseudoscalar $q\cdot S$ can contribute only linearly because its appearance is dictated by the Dirac structure of the matrix element. In four dimensions, one can reduce any Dirac structure to terms independent of or linear in S^μ. Parity invariance forbids the latter from appearing anywhere except in the tensor structures multiplying $G_{1,2}$. This leaves us with the two invariants q^2 and ν.

[3]The factors of 2 adorning G_1 and G_2 compensate for the implied factor of 1/2 in the spin vector S^μ. Note that this decomposition does not coincide with that found most often in the literature. These structure functions are proportional to those found in more standard treatments.

Our analysis up to this point has been quite general. The decomposition (2.5) is valid for all[4] values of q^2 and ν. However, in order to make contact with the fundamental degrees of freedom of perturbative QCD, we must specialize to a specific region. In the last chapter, we saw that QCD becomes nonperturbative at low energies. This means that the familiar quarks and gluons of the QCD lagrangian are no longer the relevant excitations. At these low energy scales, we must consider collective excitations of quarks and gluons ('quasi-partons') rather than the degrees of freedom introduced in the last chapter. This behavior is reminiscent of solid state physics, where strong interactions between electrons and the lattice require us to consider 'quasi-electron' excitations rather than the familiar electron-ion excitations we see in 'free' electrodynamics. After some reasonable assumptions, the lagrangian for these effective degrees of freedom can be derived directly from the fundamental lagrangian. It can easily be seen that the quasi-electrons interact weakly and are therefore the relevant degrees of freedom.[5] Presumably, something similar is happening in our case. However, since we are dealing with an ultra-relativistic quantum field theory rather than non-relativistic quantum mechanics, we can no longer smoothly make the transition from fundamental to effective degrees of freedom.[6] This leaves us with two options. We can either write down a theory which is not[7] motivated by the fundamental theory, or stick to regions in which the effective degrees of freedom are the quarks and gluons of QCD. There are many models and theories which opt for the former, among them the nuclear shell model, Yukawa's pion theory, the constituent quark model, and chiral perturbation theory. Here, we attempt to learn something about QCD itself by taking the latter path.

To this end, we will consider DIS in the *Bjorken* limit, formally defined as $Q^2, \nu \to \infty$ while

[4] Strictly speaking, we must choose physical values for q^2 and ν. We will see below that this restricts $q^2 \leq 0$ and $\nu \geq 0$. However, our structure functions can be analytically continued to almost every region of the complex plane of q^2 and ν.

[5] Renormalization can also be viewed in this way : the fundamental pointlike fields in our lagrangian interact strongly with vacuum fluctuations. The more appropriate degrees of freedom are collective excitations of the fundamental fields and vacuum fluctuations, the renormalized fields.

[6] This is merely another signal of our incompetence. Certainly, a transition can be made; we just don't know where to begin.

[7] directly

$x_B \equiv Q^2/2\nu$ remains constant. Here,[8]

$$Q^2 \equiv -q^2 = 2k^0(k')^0(1-\cos\theta) = 4k^0(k')^0\sin^2(\theta/2) \geq 0 \ , \tag{2.6}$$

with θ the angle between the initial and final electron spatial momenta. This special kinematic regime admits quarks and gluons as appropriate degrees of freedom since they interact weakly at high energy. In this limit we can ignore the proton mass and all natural scales associated with QCD. A straightforward application of dimensional analysis[9] reveals that W, and therefore all of our structure functions, is dimensionless. In the Bjorken limit, the only relevant scales in our problem are Q^2 and ν. Hence our structure functions should depend only on their ratio, x_B. While this is strictly true in the formal limit, for any finite Q^2 one will see violations of this scaling law coming from the dependence of the QCD coupling on scale. Since QCD is asymptotically free, these violations will get smaller and smaller as Q^2 is increased. This is one of the main reasons QCD was chosen as the underlying theory of the strong interactions, as detailed in the Introduction.

The kinematics of the Bjorken limit allow us to write the vectors q and P in a very simple way. Since we have only two dynamical vectors, we can always choose a frame in which their spatial momenta are parallel to the 3-axis. In this frame, we can expand q and P in terms of the two basis vectors $p^\mu = \Lambda(1,0,0,1)$ and $n^\mu = (1,0,0,-1)/2\Lambda$. Λ is an arbitrary scale reflecting our freedom to boost along the 3-axis. Note that we have chosen basis vectors which satisfy $p^2 = n^2 = 0$ and $n \cdot p = 1$. We will call the component of a vector which is parallel to p the '+' component and that which is parallel to n the '−' component. This implies

$$k^0 = \Lambda k^+ + k^-/2\Lambda \ ; \qquad\qquad k^3 = \Lambda k^+ - k^-/2\Lambda \ , \tag{2.7}$$

for any vector k. Our vectors have the expansion

$$\begin{aligned} P &= p + \frac{M^2}{2}n \\ q &= -\zeta p + \frac{Q^2}{2\zeta}n \ , \end{aligned} \tag{2.8}$$

[8] ignoring the electron mass

[9] Our states are normalized via $\langle \vec{p}\,' \,|\vec{p}\rangle = 2p^0(2\pi)^3\delta(\vec{p}\,' - \vec{p})$, as in Appendix D.1, so each state contributes mass dimension −1.

where

$$\zeta \equiv \frac{Q^2}{2x_B M^2}\left(-1+\sqrt{1+\frac{4x_B^2M^2}{Q^2}}\right) \to x_B \tag{2.9}$$

as $Q^2 \to \infty$. Since the $-$ component of P cannot form large scalars, we will ignore it whenever we can and simply write $P = p$.

Due to the role played by the light cone in special relativity, truly massless fields are not associated with spin vectors. The analogous degree of freedom for these fields is the helicity, h.[10] As shown in Appendix B.2, the replacement $MS \to hP$ is valid as $M \to 0$. While this identification gets rid of the unsightly factors of M multiplying the spin-dependent structure functions, the tensor structure associated with G_2 vanishes identically. In this term, we are certainly not free to take the proton mass to zero! As a consequence of this, we will see that the structure function G_2 is suppressed in the Bjorken limit.

We are now ready to discuss some of the contributions to DIS from various structure functions in the Bjorken limit. The lepton tensor we ignored above,

$$L^{\mu\nu} \equiv \langle k's' \left| J_e^\mu(0) \right| ks \rangle \langle ks \left| J_e^\nu(0) \right| k's' \rangle \quad , \tag{2.10}$$

can be calculated at leading order in QED. For massless electrons, only helicity eigenstates make sense. Since these do not interact with each other, the relation

$$u_\alpha(k,\lambda)\overline{u}_\beta(k,\lambda') = \frac{1}{2}\delta_{\lambda\lambda'}\left[(1+2\lambda\gamma_5)\ \not{k}\right]_{\alpha\beta} \quad , \tag{2.11}$$

where $\lambda, \lambda' = \pm 1/2$ label helicities, holds. Substitution gives

$$L^{\mu\nu} = 2e^2\left(k^\mu k'^\nu + k^\nu k'^\mu - g^{\mu\nu}k\cdot k'\right) - 4ie^2\epsilon^{\mu\nu\alpha\beta}q_\alpha(\lambda k)_\beta \quad . \tag{2.12}$$

As required by electromagnetic gauge invariance, L conserves current if the electrons are both onshell. This makes the contraction $W^{\mu\nu}L_{\nu\mu}$ especially painless. We obtain

$$\begin{aligned} W^{\mu\nu}L_{\nu\mu} \quad = \quad & 2e^2Q^2F_1(\nu,Q^2) \\ & +2e^2\left[2P\cdot k'\left(1+\frac{P\cdot k'}{\nu}\right) - x_BM^2\right]F_2(\nu,Q^2) \\ & +2e^2\lambda\frac{MS\cdot(k+k')}{\nu}Q^2G_1(\nu,Q^2) \\ & +4e^2\lambda\frac{MS\cdot(\nu k' - P\cdot k'q)}{\nu^2}Q^2G_2(\nu,Q^2) \quad . \end{aligned} \tag{2.13}$$

[10] I use the convention $h = \pm 1/2$ rather than ± 1.

If the initial and final electrons do not have transverse momentum components, the kinematics of the process require $k = q^- n$ and $k' = -q^+ p$. In this case, the quantity

$$2P \cdot k' = -M^2 q^+ = x_B M^2 + \mathcal{O}(M^4/Q^2) \tag{2.14}$$

causes the contribution of F_2 to be suppressed by M^4/Q^4 relative to that of F_1. Writing

$$MS^\mu = s^+ p^\mu + M^2 s^- n^\mu + MS_\perp^\mu \ , \tag{2.15}$$

on dimensional grounds, we see that G_1 contributes at the same order as F_1 and G_2 is suppressed by (M^2/Q^2). Processes with appreciable transverse momentum see the much smaller suppression $(k_\perp^2/Q^2)$ of F_2 and allow the transverse components of S^μ to contribute as $(MS_\perp \cdot k_\perp/Q^2)$ to $G_{1,2}$, but do not alter the leading behavior. We will discuss these contributions in Chapter 4. For now, however, we ignore the subleading structure functions and focus on F_1 and G_1.

2.2 Deep-Inelastic Scattering on Partonic Targets

The Bjorken limit was not introduced merely as a way to get rid of two structure functions (although this aspect is not without its charms). Rather, it was alleged to help us make contact with the fundamental degrees of freedom in QCD. To understand how this happens, let us take a moment to think about the physical process at hand. Our proton is a conglomerate of partons interacting strongly with each other. The process we are interested in concerns the scattering of an intruder off this complicated mess.

Fortunately, some general arguments allow us to simplify our task. Causality implies that our partonic interactions do not happen instantaneously. Consider, for example, two partons under the influence of the nonperturbative force that binds them together. Since they feel this strong restoring force, it stands to reason that they are nearing the end of their leash. Their separation distance, d, should be of the same order as the nucleon size and is related to the energy scale, $\Lambda_{\rm QCD}$, at which QCD becomes nonperturbative :

$$d \sim 1/\Lambda_{\rm QCD} \ . \tag{2.16}$$

Since they are separated by this distance, the interaction requires a time of order $1/\Lambda_{\rm QCD}$ in which to occur. On the other hand, our virtual photon is living on borrowed time. The form

of its propagator implies that it can only exist for a time of order M/Q^2 in the rest frame of the nucleon.[11] If $Q^2/M \gg \Lambda_{\rm QCD}$, the partons simply do not have time to scatter nonperturbatively during the intruder's visit. Consequently, the intruder sees a frozen non-dynamic collection of partons rather than the complicated interacting soup we do not understand. By the time the partons can re-scatter, the process we are interested in is over and done with. These final-state interactions cannot affect the total cross-section for our process[12] and hence are unimportant to our treatment.

Since the partons do not interact with each other during the scattering, we can treat each parton as though it were alone. In this way, the complete scattering tensor $W^{\mu\nu}$ becomes a weighted average over 'partonic scattering tensors' :

$$W_T^{\mu\nu} = \sum_a f_{a/T} w_a^{\mu\nu} \quad . \tag{2.18}$$

Here, $f_{a/T}$ is the probability of finding a parton of type a in our target T and $w_a^{\mu\nu}$ is the scattering tensor for a partonic 'target' of type a. The sum extends over all parton species, spin, and momentum. I emphasize here that this decomposition can be done only in cases where the scattering is *incoherent*, i.e. the partons do not interact with each other during the scattering.[13] Measurements of the running of α_s and the size of various hadrons reveals the strong interaction scale to be of order 300 MeV; as long as we have Q of order $2-3$ GeV

[11] Its propagator is of the form

$$\int \frac{d^4x}{x^2}\, e^{iq\cdot x} \quad . \tag{2.17}$$

For large $q\cdot x$, small variations in x cause wild oscillations in the integrand which effectively damp contributions. Hence dominant contributions come only from regions where *both* $q\cdot x$ *and* x^2 are small. Since q^μ is not light-like (far from it, in fact), this implies x^μ is small. How small is dictated by the components of q^μ in the frame we are using. For example, in the rest frame of the proton, the time alotted to our virtual photon is on the order of $1/q^0 \sim M/Q^2$.

[12] They can, however, affect an *exclusive* cross-section like $\gamma^* p \to n\, 2\pi^+\, \pi^-\, \pi^0$. This process could've begun as $\gamma^* p \to p\, \pi^0$, with re-scattering effects producing the observed final state long after the virtual photon scattering. This is one of the main simplifications provided by *inclusive* DIS.

[13] We have shown that for $Q^2/M \gg \Lambda_{\rm QCD}$ partons do not interact over long distances during the scattering. Short distance perturbative interactions certainly can take place during the scattering. However, the exchange of vast amounts of energy between partons within a nucleon is suppressed simply because there is not a lot of ambient energy in the medium. These kinds of parton correlation effects are suppressed by powers of Q. We will discuss a certain class of them in the last chapter.

or more, the scattering can reasonably be considered incoherent. Some of the approximations of the last section require somewhat larger Q since they ignore the nucleon mass (roughly 1 GeV). Ideally, inclusive DIS is considered at energies of order 10 GeV or larger.

Equation (2.18) is a mathematical expression of the physical separation of scales in our process. It states that the full amplitude can be *factorized* into a product of two parts. The functions $f_{a/T}$ describing the probability of finding various partons within our target, called *parton distribution functions*, are nonperturbative in nature and cannot be calculated using the techniques considered here. These objects are related to soft, low-energy physics and do not depend on the specific scattering process. On the other hand, the partonic scattering tensors $w_a^{\mu\nu}$ are expressed entirely in terms of the fundamental degrees of freedom in QCD and can in principle be calculated in perturbation theory. They are associated with the hard, high-energy physics of the scattering and do not depend on the target we consider. This separation between hard and soft physics, called *factorization*, is indispensable to modern QCD. Many recent results in perturbative QCD involve identifying and isolating the relevant distributions analogous to $f_{a/T}$ for a certain process and calculating their coefficients. The remainder of this section is devoted to a calculation of DIS on partonic targets. We shall have a lot more to say about parton distributions in the next section.

At present, perturbation theory is not readily applicable to the calculation of W because it does not involve a time-ordered product. We can remedy this in the following roundabout way. Consider the tensor

$$\int d^4z\, e^{iq\cdot z} \left\langle PS \left| J^\nu(0) J^\mu(z) \right| PS \right\rangle \quad . \tag{2.19}$$

Introducing a complete set of intermediate states and shifting the argument of J^μ to the exponent, we see that it is equal to

$$\sum_X \left\langle PS \left| J^\nu(0) \right| X \right\rangle \left\langle X \left| J^\mu(0) \right| PS \right\rangle (2\pi)^4 \delta\left(q + p_X - P\right) \quad . \tag{2.20}$$

The sum extends over all physical intermediate states with energy-momentum $p_X = P - q$. Since we intend to work in the region[14] $(q^0)_{rf} \gg (P^0)_{rf} = M$, there are no physical states

[14]The subscript rf reads 'in the rest frame of the proton'. Since the zero component of a four vector is not a scalar, we must specify a frame to give it meaning.

which satisfy the δ-function. Hence we can write

$$W^{\mu\nu} \sim \frac{1}{4\pi} \int d^4z\, e^{iq\cdot z} \left\langle PS \left| [J^\mu(z), J^\nu(0)] \right| PS \right\rangle \quad , \tag{2.21}$$

where the '$\sim$' is meant to remind us that this identification is only valid in the physical region of scattering, i.e. $(q^0)_{rf} > 0$.[15] We still do not have a time-ordered product, but can use the same trick on the tensor

$$T^{\mu\nu} \equiv i \int d^4z\, e^{iq\cdot z} \left\langle PS \left| \mathrm{T} J^\mu(z) J^\nu(0) \right| PS \right\rangle \tag{2.22}$$

to obtain

$$T^{\mu\nu} \sim i \int d^4z\, \Theta(z^0)\, e^{iq\cdot z} \left\langle PS \left| [J^\mu(z), J^\nu(0)] \right| PS \right\rangle \quad . \tag{2.23}$$

The step-function, $\Theta(z^0)$, comes from the time-ordered product. The complex conjugate of $T^{\mu\nu}$,

$$(T^{\mu\nu})^* \sim -i \int d^4z\, \Theta(-z^0)\, e^{iq\cdot z} \left\langle PS \left| [J^\nu(z), J^\mu(0)] \right| PS \right\rangle \quad , \tag{2.24}$$

differs from $T^{\nu\mu}$ only in the argument of the Θ-function. Writing

$$\Theta(z^0) = \frac{1}{2}[1 + \mathrm{sign}(z^0)] \quad , \tag{2.25}$$

where $\mathrm{sign}(x)$ returns the sign of its argument, it becomes obvious that

$$\Im m\, T^{(\mu\nu)} \sim 2\pi W^{(\mu\nu)} \tag{2.26}$$

for the symmetric parts and

$$\Re e\, T^{[\mu\nu]} \sim 2\pi i W^{[\mu\nu]} \tag{2.27}$$

for the antisymmetric parts. Decomposing $T^{\mu\nu}$ as above,

$$\begin{aligned} T^{\mu\nu} = \quad & \left(\frac{q^\mu q^\nu}{q^2} - g^{\mu\nu} \right) T_1(\nu, q^2) \\ & + \left(P - \frac{\nu}{q^2} q \right)^\mu \left(P - \frac{\nu}{q^2} q \right)^\nu \frac{1}{\nu} T_2(\nu, q^2) \\ & - i\epsilon^{\mu\nu\alpha\beta} q_\alpha S_\beta \frac{2M}{\nu} S_1(\nu, q^2) \\ & - i\epsilon^{\mu\nu\alpha\beta} q_\alpha \left(\nu S - q \cdot S\, P \right)_\beta \frac{2M}{\nu^2} S_2(\nu, q^2) \ , \end{aligned} \tag{2.28}$$

[15] Since baryon number is conserved in all interactions we will consider, the state $|X\rangle$ must include at least one baryon. Hence its energy must be at least the proton mass.



Figure 2.1: The two leading-order diagrams for virtual photon-quark scattering. The crossing symmetry allows us to obtain one from the other.

we see that the structure functions associated with W are just $1/2\pi$ times the imaginary part of those associated with T. This relation is only true in the physical region, but we are free to calculate our structure functions in any way we wish. Since T involves a time-ordered product, it is of exactly the form we need to apply perturbation theory.

Let us consider scattering on a massless target of momentum xp. The leading order amplitude involves a simple contraction of the quark currents with each other and the external states. The two relevant diagrams are shown in Figure 2.1. We need only consider one of them explicitly because the other can be obtained from crossing symmetry, $q \to -q$ and $\mu \to \nu$. This is an exact symmetry of T, independent of the QCD dynamics, as can be seen directly from its definition. Figure 2.1a is trivial to evaluate. It has the value[16]

$$\frac{e_q^2}{2\nu} \frac{1}{x_B - x - i\varepsilon} \overline{u}(h, xp)\gamma^\mu(x\not{p} + \not{q})\gamma^\nu u(h, xp) \quad , \tag{2.29}$$

where $\overline{u}(h, xp)$ and $u(h, xp)$ are the Dirac wavefunctions of a free quark of momentum xp and helicity h introduced in Appendix B.3, and e_q is the charge of a quark of flavor q.[17] As above, the Dirac product can be simplified by writing

$$u_\alpha(h, xp)\overline{u}_\beta(h, xp) = \frac{1}{2}[(1 + 2h\gamma_5)x\not{p}]_{\alpha\beta} \quad , \tag{2.30}$$

which is valid for massless spinors. In this way, the amplitude reduces to the simple Dirac

[16] We have kept our definition of $\nu = p \cdot q$ rather than changing to $xp \cdot q$ since in the end we will be concerned with the tensor structures relevant to the true target of momentum P rather than this fictitious quark target.

[17] Here, I resort to the standard notation q for quark flavor rather than f.

trace

$$\frac{e_q^2}{2\nu}\,\frac{x}{x_B - x - i\varepsilon}\,\frac{1}{2}\mathrm{Tr}\left[\gamma^\mu(x\not{p}+\not{q})\gamma^\nu(1+2h\gamma_5)\,\not{p}\right] \quad . \tag{2.31}$$

The '1' contributes only to the symmetric part of $T^{\mu\nu}$, while γ_5 contributes only to the anti-symmetric part. Concentrating on the symmetric part for the moment, we find

$$\begin{aligned} T_q^{(\mu\nu)} &= xe_q^2\left(\frac{1}{x_B - x - i\varepsilon} - \frac{1}{x_B + x - i\varepsilon}\right)\left(\frac{q^\mu q^\nu}{q^2} - g^{\mu\nu}\right) \\ &+ 2x^2 e_q^2\left(\frac{1}{x_B - x - i\varepsilon} + \frac{1}{x_B + x - i\varepsilon}\right)\frac{1}{\nu}\left(p - \frac{\nu}{q^2}q\right)^\mu\left(p - \frac{\nu}{q^2}q\right)^\nu \quad , \end{aligned} \tag{2.32}$$

where the crossing term, $q \to -q$ and $\mu \leftrightarrow \nu$, has been taken into account.[18] The structure functions relevant to our experiment are now easy to compute :[19]

$$\begin{aligned} {}_qF_1(x,\nu,Q^2) &= xe_q^2\frac{1}{2}\delta(x_B - x) \\ {}_qF_2(x,\nu,Q^2) &= x^2e_q^2\delta(x_B - x) = 2x_B\ {}_qF_1(x,x_B) \quad . \end{aligned} \tag{2.33}$$

Here, we see explicitly that the structure functions scale; they depend only on the ratio x_B between of the two large scales ν and Q^2.[20] Along with the Callan-Gross relation, ${}_qF_2 = 2x_B\ {}_qF_1$, scaling is a general property of this kind of scattering process. This behavior can be shown quite generally via the current algebra techniques of Gell-Mann [25]. The fact that it is broken by quantum effects was one of the main motivations of D. Gross' search for a proof that quantum field theory cannot describe experiment. His subsequent discovery of asymptotic freedom turned his argument on its head and led to the realization that only a theory like QCD can be consistent with experiment.

The antisymmetric part of T is given by the γ_5 term in the trace. The result is

$$T_q^{[\mu\nu]} = -i\epsilon^{\mu\nu\alpha\beta}q_\alpha(hp)_\beta\frac{2xe_q^2}{\nu}\left(\frac{1}{x_B - x - i\varepsilon} + \frac{1}{x_B + x + i\varepsilon}\right) \quad . \tag{2.34}$$

This implies

$${}_qG_1(\nu,Q^2) = xe_q^2\frac{1}{2}\delta(x_B - x) = {}_qF_1(\nu,Q^2) \quad . \tag{2.35}$$

[18] Note that the crossing is not exactly $\mu \leftrightarrow \nu$ and $q \to -q$ since the $+i\varepsilon$ must also change sign. This fact is related to the analytic properties of $T^{\mu\nu}$ as a function of x_B.

[19] The $+i\varepsilon$'s introduced in Section 1.2 play a pivotal role here.

[20] The dependence of T_q on the + component of the quark momentum is viewed as dependence on the type of parton.

Once again, we see scaling.

The presence of $\delta(x_B - x)$ in our structure functions is easy to understand from the standpoint of 'physical' quark scattering. At leading order, the only intermediate state accessible to the quark-photon system is an excited quark. If this state is to be physical, the quark must be onshell. This implies $2x\nu = Q^2$, or $x = x_B$.

Our amplitude requires corrections due to virtual gluon exchange, as shown in Figure 2.2. As presented, these diagrams are riddled with ultraviolet divergences. The discussion on renormalization in Section 1.4 tells us how to deal with these divergences. It is simplest to use the $\overline{\text{MS}}$ scheme in Feynman's gauge ($\xi = 1$). In renormalized perturbation theory, there are three relevant counterterms. The obvious insertion is $i(Z_F - 1)(x\not{p} + \not{q})$ for the intermediate quark propagator. The other two come from the electromagnetic sector of the theory.

Consider for a moment the quark-photon interaction term in QED. Its structure is identical to the quark-gluon interaction with t^a replaced by 1:

$$\mathcal{L}_{int}^{\text{QED}} \sim -Q_q e_b \overline{\psi}_b \not{A}_b \psi_b \quad , \tag{2.36}$$

where Q_q is the quark charge, e_b is the bare electromagnetic coupling,[21] and A_b is the bare photon field. Defining our Z-factors in the same way as for QCD, we see that this leads to the counterterm

$$-(Z_e Z_F Z_A^{1/2} - 1) Q_q e \, \overline{\psi} \not{A} \psi \tag{2.37}$$

in the QED lagrangian. Electromagnetic gauge invariance requires the running of the coupling to be the same for *all* electromagnetically charged particles. Hence the QCD ultraviolet divergence structure of the quark electromagnetic current must be identical to the divergence structure of the quark self-energy. This must be satisfied *diagram by diagram*, i.e. for each

[21] The *electromagnetic coupling* is a parameter which summarizes the strength of the interaction of *all* charged fields with the photon. It is analogous to the QCD coupling g. The *charge* of a field describes the way a field transforms under the gauge group. It is analogous to the **3** of $SU(3)$. For example, the *charge* of an up quark is $2/3$ since under an electromagnetic gauge transformation parameterized by α its phase rotates by the amount $2\alpha/3$. In contrast, the (renormalized) *electromagnetic coupling* $e \sim \sqrt{4\pi/137} \sim 3/10$ is the same for all electromagnetically charged fields. When the difference between these two objects is inessential, it is often ignored by introducing the 'quark electromagnetic charge' $e_q \equiv Q_q e$. However, one must always remember that they are *not* the same. In particular, the coupling e is just a number while the charge Q is a quantum mechanical operator describing the behavior of fields under a gauge transformation.

QCD diagram the sum of all possible electromagnetic current insertions must have an ultraviolet divergence which is identical to (the negative of) that of the corresponding self-energy diagram. The cancelation, being a consequence of *electromagnetic* gauge invariance, is completely independent of the QCD dynamics.

The full counterterm contribution to the amputated diagrams is taken into account simply by multiplying the leading order result by Z_F. While this is all that is necessary to cancel the ultraviolet divergences, it will not result in a properly normalized amplitude. Since we have not used the onshell renormalization scheme, the residue of our renormalized propagator is not i. While this fact is unimportant for intermediate quark propagation, our external quark wavefunctions must be correctly normalized.[22] This is done exactly as in the onshell scheme : we multiply the amputated Feynman diagrams by the full (1PR) propagators, (re)renormalize the external states by dividing by the square root of the propagator residue,[23] multiply by the inverse propagator, take the onshell limit, and attach external free wavefunctions. This procedure is known as Lehmann-Symanzik-Zimmermann (LSZ) reduction formula [51]. It is quite general and applies to all quantum field theories, regardless of whether or not they are renormalizable. Since the full propagators contain a power of the residue,[24] the end result is simply multiplying the amputated graphs by the square root of the residue for each physical external state.

To see this explicitly, let us examine the fully renormalized quark propagator in this scheme. The 1PI graphs sum to $-i\Sigma = -iB(p^2)\not{p}$, with a counterterm contribution $i(Z_F - 1)\not{p}$. The full propagator becomes

$$S(p) = \frac{i}{(Z_F - B(p^2))\ \not{p}} \quad . \tag{2.38}$$

The residue of this object at $\not{p} = 0$ is $[Z_F - B(0)]^{-1}$. Since $B(p^2)$ is dimensionless, it may be

[22] Since this calculation does not represent actual physical scattering, the difference between properly and improperly normalized quark fields can be absorbed into the distribution functions. However, if we wish to relate these functions to the effective high-energy degrees of freedom of QCD, we has better normalize them properly.

[23] The word 'residue' is used in this context to mean 'mathematical residue divided by i', or 'thing that should be 1'.

[24] Actually, since we intend to take the onshell limit after all of these manipulations, the full propagator is effectively *just* the pole contribution.

written[25]

$$B(p^2) = \sum_{n=1}^{\infty} \alpha_s^n B_n \left(\frac{p^2}{\mu_0^2} \right)^{-n\epsilon/2} \quad . \tag{2.39}$$

Since our renormalized amplitude is free of ultraviolet divergences, we are free to take $\epsilon < 0$.[26] In this case, it becomes obvious that $B(0) = 0$. This implies that the correct amplitude is obtained by multiplying the amputated result by $Z_F^{-1/2}$ for each external quark leg, or $1/Z_F$ in all. It must be emphasized that this factor is *not* ultraviolet divergent.[27] All of the ultraviolet divergences have been removed by renormalization. This factor contains the *infrared* divergence generated by taking the onshell limit. However, as long as we recognize that there are no ultraviolet divergences left in the problem, we may allow this Z_F to cancel the one from the counterterms. This means that after all is said and done all of the Z_F factors necessary for a properly normalized amplitude cancel. The true amplitude is simply the sum of the diagrams, without a thought given to counterterms or proper normalization. We must keep in mind, though, that all of the divergences we run into from now on are infrared and cannot simply be swept into renormalization constants.

In unrenormalized perturbation theory, the story is somewhat different. The procedure here is to calculate *all* diagrams, renormalize *external* states, and express the result in terms of renormalized quantities at the end of the day. The result of these manipulations is guaranteed to be an ultraviolet finite quantity. In addition, onshell external states require infrared (re)renormalization as discussed above. In our case, the electromagnetic currents do not require renormalization at this order in QED.[28] As mentioned above, all of their ultraviolet divergences cancel with intermediate self-energy diagrams. The external states, on the other hand, require an overall division by Z_F to cancel their ultraviolet divergences. The amplitude we are left with is guaranteed to be free of ultraviolet divergences. Once again, we (re)renormalize the onshell external states through multiplication by an infrared divergent Z_F. The end result is

[25] As before, the μ_0's appear to keep the coupling dimensionless.

[26] We *cannot* do this in the unrenormalized amplitude since ϵ is required positive to regulate the ultraviolet divergences. Only in an ultraviolet finite quantity can we take $\epsilon < 0$.

[27] This is why we never had to take it into account before. If we are concerned only with the ultraviolet behavior of an operator, it is best not to take the onshell limit so this kind of 'mixing of divergences' does not occur.

[28] At higher orders, the renormalization of these currents is reflected in the renormalized coupling.




Figure 2.2: Contributions to virtual photon-quark scattering at (a) leading and (b)-(e) next-to-leading order in the QCD coupling.

that all of our Z_F factors cancel; we need only calculate the amputated diagrams and attach onshell external wavefunctions.

The final step in the LSZ program involves taking the onshell limit of our amplitudes. As we have seen, this leads to infrared divergences in general. Although they seem like a terrible defect at first, the appearance of these divergences is quite natural. Their origins and treatment are discussed in the next section; for now, we simply acknowledge their existence and proceed.

There are several ways to do this calculation. One method is that used for the leading order amplitude. The unpolarized (symmetric) part of T is calculated by averaging over external helicities and the polarized (antisymmetric) part is obtained by taking half the difference of right- and left- handed helicities. In this way, both parts become a trace which can be simplified before performing the loop integration. Unfortunately, subtleties associated with the definition of γ_5 in d dimensions[29] make the polarized result ambiguous. Thus, in order to resolve this ambiguity, one must choose a consistent γ_5 scheme. The most natural thing to do is simply perform the loop integration over the full Dirac structure, leaving the introduction of γ_5 for later. This approach has the advantage of obtaining both the polarized and unpolarized amplitudes simultaneously. After the loop integration, the only vectors left over are external and may be taken to be in the first four dimensions. At this point, we are free to use the above

[29] see Appendix E.

method to extract the symmetric and antisymmetric parts explicitly.

The intermediate quark self-energy diagram, Figure 2.2b, can be done using the techniques of Appendix A.3. Ignoring terms of order ϵ, the result is

$$\frac{\alpha_s C_F}{4\pi}\left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[\frac{2}{\epsilon}+1-\log\frac{x_B-x-i\varepsilon}{x_B}\right]\frac{e_q^2}{2\nu}\,\frac{1}{x-x_B+i\varepsilon}\gamma^\mu(x\not{p}+\not{q})\gamma^\nu \\ +(\mu\leftrightarrow\nu, q\to -q)\ , \qquad (2.40)$$

where the external states have been omitted. Once again, we see the addition $(\mu\leftrightarrow\nu, q\to -q)$ from the photon crossing diagram. In the interest of space, this crossing contribution is implied in all of the following results unless indicated otherwise. As before, the $+i\varepsilon$ plays a fundamental role in the extraction of the imaginary part of the amplitude. Note that we can ignore it here simply by endowing x_B with a small negative imaginary part, $\tilde{x}_B \equiv x_B - i\varepsilon$. This behavior persists in all of our diagrams, provided that the crossing is obtained by $\tilde{x}_B \to -\tilde{x}_B$.[30] In the following, we drop the tildé on $\tilde{x}_B$ and simply remember that x_B has a small negative imaginary part.

The calculation of the remaining diagrams is greatly simplified by taking μ and ν in the transverse directions. A look at the tensor structures relevant to F_1 and G_1 tells us that this simplification is free of charge - none of the leading dynamics is affected by it. Since the external states in our calculation satisfy the free equation of motion, $\not{p}u(xp,h)=0$, the intermediate self-energy diagram may be rewritten for transverse $\mu\nu$ as

$$\frac{\alpha_s C_F}{4\pi}\left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[\frac{2}{\epsilon}+1-\log\frac{x_B-x}{x_B}\right]\frac{e_q^2}{2}\,\frac{1}{x-x_B}\,\gamma^\mu\not{n}\gamma^\nu\ . \qquad (2.41)$$

We will see that all of our amplitudes can be expressed in terms of this one simple Dirac structure and its hermitian conjugate.

An application of the formulae in Appendix A.3 to the integrals encountered in the remaining diagrams is possible, but extremely tedious. This method of covariant integration does not use the special kinematics of our problem. A superior method[31] breaks the inte-

[30] One can see this by examining each diagram separately, keeping the ε's explicitly. Alternatively, a study of the amplitude in the complex x_B-plane reveals T to be an analytic function of x_B with branch cuts along the real axis stemming from $\pm x$. The discontinuity relevant to our physical structure functions can then be obtained by approaching the positive real axis from underneath.

[31] This method is identical to that employed in Section 1.6. The presence of a nonzero scale complicates this process, so we discuss its application explicitly in this case.

gration into +, −, and ⊥ components : $d^d k = dk^+ dk^- d^{d-2}k_\perp$. The definitions of + and − ensure that the measure is invariant under this rotation. Defining $y \equiv k^+$, we may substitute $k^\mu = yp^\mu + k^- n^\mu + k_\perp^\mu$ into our expressions. The fact that $n^2 = p^2 = 0$ and none of our external momenta have transverse components leads to a tremendous simplification. In particular, the large k^- behavior of the integral is never worse than $\int dk^-/(k^-)^2$. This sets the stage for a contour integration in the k^--plane. Restoring the $+i\varepsilon$'s, we can analyze the pole structure of the integrand. Since k^- is always accompanied by a function of y, the position of its pole will drop from above the real axis ($y \ll 0$) to below the real axis ($y \gg 0$) as y makes its trek from $-\infty$ to $+\infty$. The precise position of the drop point depends on the pole in question, but for either very large y or very large $-y$ all of the poles appear either below or above. We are free to close the contour in the pole-free half of the plane in these regions, giving no contribution to the integral. For intermediate y, we may close the contour in either the upper- or the lower-half-plane; Cauchy's theorem assures us the result will be identical. The transverse dimensions now have an integral of the form[32]

$$\int \frac{d^{d-2}k_\perp}{(2\pi)^{d-2}} \frac{\alpha Q^2 + \beta k_\perp^2}{(k_\perp^2 + M^2)(k_\perp^2 + N^2)} \ , \tag{2.43}$$

where α and β are functions of x,x_B, and y, which can easily be done via the methods of Appendix E. The final integration can always be done with the formula

$$\int_0^1 dy\, y^{\alpha-1}(1-y)^{\beta-1} = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} \tag{2.44}$$

after a suitable change of variables.

This analysis is complicated by the existence of physical intermediate states for $x_B \geq x$. These states lead to branch cuts in our amplitude and generally get in the way. We can avoid them *during* the calculation by working in the unphysical region, $x_B \gg x$. At the end of the day, we are free to analytically continue x_B to the physical region of interest.

The two vertex diagrams (Figs. 2.2c and d) are related by hermitian conjugation[33] and $\mu\nu$

[32] We can always choose $N^2 = 0$ by refusing to pick up the Q^2-dependent pole. In this case, both integrals reduce to

$$\int \frac{d^{d-2}k_\perp}{(2\pi)^{d-2}} \frac{1}{k_\perp^2 + M^2} \ . \tag{2.42}$$

[33] This conjugation is to be understood as taking place only in the Dirac sector. The $-i\varepsilon$ associated with x_B

exchange, as can be seen directly from their expressions. This symmetry leaves us with only two diagrams to calculate explicitly. Including both contributions, the vertex gives

$$-\frac{\alpha_s C_F}{4\pi} e_q^2 \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \left\{ \frac{1}{x_B - x}\left(\frac{2}{\epsilon} + 4\right) + \left[\left(\frac{1}{x} + \frac{1}{x_B - x}\right)\left(\frac{4}{\epsilon} + 3 - \log\frac{x_B - x}{x_B}\right) - \frac{1}{x_B - x}\right]\log\frac{x_B - x}{x_B}\right\} \gamma^\mu \not{n} \gamma^\nu \quad . \tag{2.45}$$

The presence of a squared propagator in the final diagram (Fig. 2.2e) complicates matters slightly. However, since there are still only three poles, our method can easily be modified to handle this situation. We obtain

$$-\frac{\alpha_s C_F}{4\pi} e_q^2 \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \left\{ \frac{2}{\epsilon}\frac{1}{2x}\left(\gamma^\mu \not{n} \gamma^\nu + \gamma^\nu \not{n} \gamma^\mu\right) + \left[\left(\frac{2}{\epsilon} - \frac{1}{2}\log\frac{x_B - x}{x_B}\right)\gamma^\mu \not{n} \gamma^\nu + \frac{5}{2}\left(\gamma^\mu \not{n} \gamma^\nu - \gamma^\nu \not{n} \gamma^\mu\right)\right]\frac{x_B - x}{x^2}\log\frac{x_B - x}{x_B}\right\} \quad . \tag{2.46}$$

All of our diagrams contain infrared divergences[34] which do not cancel in the full amplitude,

$$-\frac{\alpha_s C_F}{8\pi} e_q^2 \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \left\{ \left(\frac{2}{\epsilon} + 3\right)\frac{3}{x_B - x}\gamma^\mu \not{n} \gamma^\nu + \left[\left(\frac{x_B + x}{x^2} + \frac{2}{x_B - x}\right)\left(\frac{4}{\epsilon} + 3 - \log\frac{x_B - x}{x_B}\right) - 3\left(\frac{x_B - x}{x^2} + \frac{1}{x_B - x}\right)\right]\log\frac{x_B - x}{x_B}\,\gamma^\mu \not{n} \gamma^\nu + \frac{2}{\epsilon}\,\frac{1}{x}\left(\gamma^\mu \not{n} \gamma^\nu + \gamma^\nu \not{n} \gamma^\mu\right) + 5\,\frac{x_B - x}{x^2}\log\frac{x_B - x}{x_B}\,\left(\gamma^\mu \not{n} \gamma^\nu - \gamma^\nu \not{n} \gamma^\mu\right)\right\} \quad , \tag{2.47}$$

as promised in the beginning of this chapter. At this point, we are free to project out the

does *not* change sign. Since the final state wavefunction comes with a γ^0, $\overline{u} \equiv u^\dagger \gamma^0$, the conjugation simply reverses the order of the γ-matrices.

[34]Strictly speaking, the self-energy diagram we calculated first does not contain any infrared divergences. It contains only ultraviolet divergences which are destined to cancel with counterterms. However, since we allowed the ultraviolet divergences in our counterterms to cancel the infrared divergences associated with the LSZ reduction formula, these ultraviolet divergences have been transmuted into infrared ones.



Figure 2.3: Leading diagrams for gluon-virtual photon scattering.

structure functions as above. We obtain

$$\begin{aligned} T_1^q(x,x_B) = -\frac{\alpha_s C_F}{4\pi} e_q^2 \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \Bigg\{ \left(\frac{2}{\epsilon}+3\right)\frac{3x}{x_B-x} + \frac{4}{\epsilon} \\ + \left[\left(\frac{x_B+x}{x}+\frac{2x}{x_B-x}\right)\left(\frac{4}{\epsilon}+3-\log\frac{x_B-x}{x_B}\right) \right. \\ \left. -3\left(\frac{x_B-x}{x}+\frac{x}{x_B-x}\right)\right]\log\frac{x_B-x}{x_B}\Bigg\} \\ + (x_B \to -x_B) \end{aligned} \tag{2.48}$$

for the symmetric part and

$$\begin{aligned} S_1^q(x,x_B) = -\frac{\alpha_s C_F}{4\pi} e_q^2 \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \Bigg\{ \left(\frac{2}{\epsilon}+3\right)\frac{3x}{x_B-x} \\ + \left[\left(\frac{x_B+x}{x}+\frac{2x}{x_B-x}\right)\left(\frac{4}{\epsilon}+3-\log\frac{x_B-x}{x_B}\right) \right. \\ \left. +7\,\frac{x_B-x}{x}-3\,\frac{x}{x_B-x}\right]\log\frac{x_B-x}{x_B}\Bigg\} \\ - (x_B \to -x_B) \end{aligned} \tag{2.49}$$

for the antisymmetric. In these expressions, the crossing contribution has been included explicitly.

At this order in QCD, virtual photons can also scatter off gluons in our target via the virtual quark loop diagrams in Figure 2.3. These diagrams must be ultraviolet finite since there are no counterterms at this order. The two gluons have transverse polarizations[35] i and

[35] As explained in Appendix B.2, massless particles are classified by their helicity. For vector objects, like the gluon, the form of the rotation generators implies that helicity eigenstates are obtained only for *transverse* polarization.

j. Once again, there are polarized and unpolarized contributions to the amplitudes. These can be obtained in exactly the same way as for the quarks, by performing the integration first and projecting out the symmetric and antisymmetric parts of the amplitude later. When averaging over the gluon polarizations in d dimensions, we divide by $d-2$ rather than 2 because this is the number of polarization states available to a gluon in d dimensions. The result is

$$T_1^g(x,x_B) = \frac{\alpha_s T_f}{2\pi}\left(\sum_q e_q^2\right)\left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left\{-\frac{4}{\epsilon}-2 + \left[\left(1+2x_B\frac{x_B-x}{x^2}\right)\left(\frac{4}{\epsilon}+4-\log\frac{x_B-x}{x_B}\right)-2\right]\log\frac{x_B-x}{x_B}\right\} + (x_B\to -x_B) \quad (2.50)$$

for the symmetric structure and[36]

$$S_1^g(x,x_B) = \frac{\alpha_s T_f}{2\pi}\left(\sum_q e_q^2\right)\left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \times\left\{\left[\left(1+2\frac{x_B-x}{x}\right)\left(\frac{4}{\epsilon}+4-\log\frac{x_B-x}{x_B}\right)-2\right]\log\frac{x_B-x}{x_B}\right\} - (x_B\to -x_B) \quad (2.51)$$

for the antisymmetric. Once again, we see infrared divergences. These divergences are intimately related to those in the quark sector, as we will see in the next section.

2.3 Infrared Divergences and Parton Distributions

As we have seen, DIS on onshell massless quark and gluon targets contains infrared divergences which do not cancel in 'physical' amplitudes. One might say, "Who cares? These amplitudes are not physical anyway, so it's not important that they are unavoidably divergent.". However, the physical arguments of last section indicate that these amplitudes contribute to physical scattering processes involving hadrons. The relation between physical and partonic amplitudes involves probability functions which are certainly not expected to diverge. Furthermore, since non-abelian nature of QCD contributes only trivially to the calculations of the last section, there is no way that our calculation can 'know' that the process isn't physical. In fact, this

[36] For the antisymmetric structure, the easiest way to get the result is by explicitly taking the photon and gluon indices in the 1- and 2-directions and interpreting the result as multiplied by $i\lambda\epsilon^{-+12}$.

problem also exists in QED where the process *is* physical. How, then, can we make sense of these divergences?

In QED, one argues that the quantity we have calculated is not really physical. If we were to consider an actual experiment, we would have no way to distinguish an electron of momentum k from an electron-photon *system* of momentum k in which the photon carried either a very small (soft) momentum or moved collinear to the electron. These two situations are considered separate in the calculation. If one wants to compare with experiment, one should add the cross-section for these processes to this one. It is from precisely these regions of momentum space that the infrared divergences come.[37]

Since infrared divergences come from regions of soft or collinear momentum, they are associated mainly with long time scales. The arguments of last section then imply that these regions are independent of the hard scattering. They correspond to processes in which, for example, a quark splits into another quark and a collinear gluon long before the scattering. In effect, it is only the second quark which participates in the hard scattering.[38] This fact is in direct conflict with the definition of $w_a^{\mu\nu}$. These quantities were *defined* as that part of the high-energy scattering which involves a parton of type a. The above discussion implies that the object we calculated contains a piece in which the parton relevant to the scattering is of a different type. Apparently, this new type of parton can be found inside our old parton with a certain probability. This is reminiscent of the discussion of last section. There, we considered the target as a loosely (over the relevant time scale) bound collection of partons and discovered that the scattering takes place with each parton incoherently. Here, we see that by probing at a large scale the partons themselves can break up into other partons. This suggests that we

[37] It is also precisely these regions of momentum space that one cannot properly take into account in QCD perturbation theory. The β-function of QCD implies that the coupling will *grow* as μ^2 is *decreased* until it no longer makes sense to use perturbation theory. Of course, at that point we don't know precisely what the behavior of the β-function is because we cannot calculate it. However, one thing is sure - in the regions associated with the infrared divergences of the last section the dynamics of QCD is nonperturbative. This fact will be addressed below.

[38] This analysis is complicated by diagrams like Fig. 2.2c. However, as we will see below, its inclusion does little more than ensure the gauge-invariance of the distributions $f_{a/T}$.

consider our partons as targets in the sense of Equation (2.18) :

$$T_a^{\mu\nu} = \sum_{a'} f_{a'/a}\, t_{a'}^{\mu\nu} \quad . \tag{2.52}$$

In the last section, we calculated the left-hand-side of this equation rather than the coefficients $t_a^{\mu\nu}$ on the right-hand-side. The nature of these coefficients implies that they *cannot* be calculated directly. Our procedure for perturbative calculations necessarily involves integration over all portions of momentum space, which will lead unavoidably to contributions from low-energy. The functions $f_{a'/a}$, on the other hand, involve no explicit momentum cutoff. It is merely expected that they do not receive corrections from asymptotically large momentum scales. Unfortunately, the regions in which they are expected to be generated are nonperturbative in nature. We have already included contributions from these regions in $T_a^{\mu\nu}$.

Although these contributions have been calculated incorrectly, the method we used can be extended *consistently* to incorrect perturbative calculations of other quantities, $f_{a'/a}$ in particular. Since the incorrect contributions are calculated in the same way, we can use Equation (2.52) to cancel them out and extract correct expressions for $t_a^{\mu\nu}$ order by order in perturbation theory. It is these quantities (rather, the imaginary parts of their form factors) which appear in the physical cross-section and will be free of divergences. A formal proof of this statement involves a complete classification of the regions which generate infrared divergences in DIS and an understanding of the relationship between these regions and the regions which generate the parton distributions. An outline of such a proof appears in the next section. For now, let us see explicitly how it works at one-loop order. Along the way, we will learn a great deal about the parton distributions.

In order to calculate parton distribution functions in perturbation theory, we must write them as matrix elements. The probability of finding a quark of flavor q' and momentum xp in a quark state of flavor q and momentum p is the same as the probability that I can *remove* a quark of flavor q' and momentum xp[39] from a quark state of flavor q and momentum p and

[39]The final expression we will use to define $f_{q'h'/qh}(x)$ actually receives contributions from all partons whose momentum k satisfies $n\cdot k = x$. The other components of k give subleading contributions to the hard scattering. These kinds of corrections are discussed in Chapter 4.

be left with a physical[40] state :

$$\begin{aligned} f_{q'h'/qh}(x) &\sim \sum_X \langle ph|\, \psi^\dagger_{q'}(0)\,|X\rangle\,\langle X\,|\psi_{q'}(0)|\,ph\rangle \delta(1-x-n\cdot p_X) \\ &\sim \int \frac{d\lambda}{2\pi} e^{-i\lambda x}\,\langle ph|\,\psi^\dagger_{q'}(\lambda n)\psi_{q'}(0)\,|ph\rangle \quad . \end{aligned} \tag{2.53}$$

This expression can be viewed as an impressionistic view of the distributions functions at best since its derivation completely ignores the requirement of gauge invariance and the presence of spinor indices on the quark fields. The latter complaint concerns dependence on h'.[41] Since we wish to consider both values, we will leave the spinor indices free for now, but multiply $\psi^\dagger$ by γ^0 to form a true Dirac index. Achieving gauge invariance is also quite simple. According to Appendix C, our distribution is in need of a gauge link between 0 and λn. The path the gauge link traverses should be along n^μ to minimize its dependence on λ. In this way, we arrive at the expression

$$\left(\mathcal{M}_{q'/q}\right)_{\alpha\beta}(x) = \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle ph \left| \overline{\psi}^{q'}_{\beta}(\lambda n)\, \mathcal{G}_n(\lambda n, 0)\, \psi^{q'}_{\alpha}(0) \right| ph \right\rangle \quad . \tag{2.54}$$

Although $\mathcal{M}$ is the minimal gauge invariant extension of (2.53), its interpretation is quite different. The first manipulation changes it from a probability to a density matrix. This allows us to consider both helicity states simultaneously, much as considering $f_{a'/a}$ as a function of x allows us to combine a continuously infinite number of probabilities into one probability *function*. Our density matrix can readily be split into two probability functions with the help of some projection operators. However, the explicit form of the gauge link

$$\mathcal{G}_n(\lambda n, 0) \equiv \mathrm{P}\left\{ e^{-ig\int_0^\lambda n\cdot\mathcal{A}(\zeta n)d\zeta} \right\} \tag{2.55}$$

reveals that the second manipulation has changed the *nature* of our simple distribution. Rather than the removal and subsequent replacement of a quark, $\mathcal{M}$ describes a correlation between quark and gluon fields. This complication is unavoidable in gauge theories because quarks do not make sense by themselves. The fact that a quark field transforms under a gauge transformation implies that 'the probability of finding a quark' is itself not a gauge invariant

[40] In calculations with partonic external states, the word 'physical' is used to mean physical in the absence of confinement. I.e. a transversely polarized onshell gluon is a 'physical' state while an offshell quark is not. Since ghosts are nothing but mathematical artifacts, they cannot be present in 'physical' states.

[41] Parity invariance implies that we can take $h = +1/2$ without loss of generality.

concept. However, once we choose a gauge this statement has meaning. In particular, the gauge restriction $n \cdot \mathcal{A} = 0$ makes our gauge link trivial. In this gauge, we are free to interpret $\mathcal{M}$ as a true quark density matrix. Unfortunately, the difficulties associated with this singular gauge encourage us to make a less controversial choice. Nevertheless, its existence has taught us that the gluons appearing in the gauge link are unphysical in the sense that they can be removed by a gauge rotation.[42] Since they are unphysical, we can view them as gauge artifacts and continue to interpret $\mathcal{M}$ as a quark density matrix.

The gluons fields in the gauge link are evaluated at space-time points between the two quark fields. These contributions are generated in DIS by diagrams like Fig. 2.2c in which a gluon interacts with the quark *during* the scattering. Although the arguments above indicate that there is not enough time for this to occur, the inclusion of this graph is required by gauge invariance. Since all of our arguments concerning factorization have depended critically on the idea that no low-energy QCD interactions can happen during the scattering, this process could potentially be very dangerous. The offensive contribution is of the form

$$\gamma^\mu \frac{\not{p}+\not{q}}{(p+q)^2}\gamma^\alpha \frac{y\not{p}+\not{q}}{(yp+q)^2}\gamma^\nu \tag{2.56}$$

in the collinear region, where the gluon has momentum $(1-y)p$. In the Bjorken limit, the leading contribution has Dirac structure $\gamma^\mu \not{n} \gamma^\alpha \not{n} \gamma^\nu$. This structure vanishes unless $\alpha = -$, so leading behavior is associated with the field $\mathcal{A}^+ = n \cdot \mathcal{A}$.[43] This implies that our troublesome graph does not contribute to the leading behavior in light-cone gauge. Furthermore, since the gluon momentum is proportional to p^μ, the leading region is equivalent to one in which the gluon field has been replaced by its momentum. Writing

$$(1-y)p_\alpha \gamma^\mu \frac{\not{p}+\not{q}}{(p+q)^2}\gamma^\alpha \frac{y\not{p}+\not{q}}{(yp+q)^2}\gamma^\nu \sim \gamma^\mu \frac{y\not{p}+\not{q}}{(yp+q)^2}\gamma^\nu - \gamma^\mu \frac{\not{p}+\not{q}}{(p+q)^2}\gamma^\nu \ , \tag{2.57}$$

it becomes obvious that this gluon insertion does not constitute a nontrivial contribution to the scattering in the infrared region. In reality, this graph simply represents a combination of scattering off of a quark of momentum yp (just the quark) and a quark of momentum p (coherent scattering off of the quark and the gluon combined). We will see in the next section

[42] This fact is critically dependent on the choice of path implicit in the definition of the gauge link. *Any* other choice will necessarily introduce physical gluons, destroying the desired interpretation of $\mathcal{M}$.

[43] Remember that γ^α came from the interaction $\not{\mathcal{A}} = \mathcal{A}^+\gamma^- + \mathcal{A}^-\gamma^+ - \vec{\mathcal{A}}_\perp \cdot \vec{\gamma}_\perp$.



Figure 2.4: Feynman rules for eikonal line propagation.

that this behavior generalizes to any number of gluons to all orders in α_s as a consequence of current conservation in QCD. Returning to the parton distribution, we see that the gluon fields in the gauge link are generated by graphs like these. They are required by gauge invariance, but the same principle also forbids them from having a nontrivial effect on the hard scattering. Nonetheless, they are present and must be taken into account in any nonsingular gauge.

Now that we have unambiguously defined what we mean by a parton distribution function, we would like to calculate it in perturbation theory. Once again, we find ourselves in an awkward situation because of the lack of a time-ordered product. The trick we used in the last section cannot be applied here because the term we would like to add is not strictly zero. More importantly, the gauge link will not permit this sleight-of-hand. However, we can massage this quantity into a more useful form by writing

$$\begin{aligned}
\Psi(\lambda n) &\equiv \mathcal{G}_n(\infty,\lambda n)\psi(\lambda n) \\
\mathcal{M}^{\alpha\beta}_{q'/q}(x) &= \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle ph \left| \overline{\Psi}^{\beta}_{q'}(\lambda n)\, \Psi^{\alpha}_{q'}(0) \right| ph \right\rangle \\
&= \int \frac{d\lambda}{2\pi} e^{-i\lambda(x+n\cdot p_X - 1)} \sum_X \left\langle ph \left| \overline{\Psi}^{\beta}_{q'}(0) \right| X \right\rangle \left\langle X \left| \Psi^{\alpha}_{q'}(0) \right| ph \right\rangle \quad .
\end{aligned} \tag{2.58}$$

The fields Ψ are gauge invariant in the sense that gauge transformations at infinity can be taken as trivial in perturbation theory.[44] Since the 0-component of n^μ is positive, the path-ordering of the exponential is equivalent to a time-ordering operation in the matrix element on the right. This exponential represents a coherent sum of gluon fields propagating from 0 to ∞. Coherent propagation of this kind is known as an *eikonal* line and plays an important role in proofs of factorization. The eikonal propagation itself is represented in Feynman diagrams as a double line. The Feynman rules for its propagation and interaction with gluon fields are shown in Figure 2.4.

[44] The nontrivial topology of our gauge group will cause problems here if we intend to do a calculation outside of perturbation theory.



Figure 2.5: Contributions to the 'quark distribution in a quark' at (a) leading and (b)-(d) next-to-leading order in QCD. The final state in (a) and (b) is the vacuum, while (c) and (d) represent gluonic external states.

The calculation of $\mathcal{M}$ is now straightforward : we simply draw all Feynman diagrams corresponding to the matrix element on the right for a given final state, sum their contributions, multiply by the complex conjugate[45], and sum over final states. For quark external states, the necessary diagrams up to order α_s are shown in Figure 2.5. At this order there are only two final states - the vacuum and the one gluon state. The sum over gluon final states involves an integration[46] over onshell positive-energy momenta and a sum over polarization states. This integration is greatly simplified by the δ-function constraint on the gluon's + momentum imposed by the λ integration. The polarization sum in Feynman's gauge is given by $-g_{\mu\nu}$.

The only subtlety left is that all corrections to this object seem to be exactly zero in dimensional regularization due to the lack of scale. As discussed in Appendix E, this zero represents a cancelation between infrared and ultraviolet divergences. However, we really wish to consider a *renormalized* matrix element defined at a certain scale μ^2 to be free of ultraviolet divergences.[47] This object is formally defined through its moments, which will be introduced in Section 2.5. The local operators they are related to can be renormalized with a Z-factor as with the quark-gluon current $\overline{\psi}\not{A}\psi$ in the last chapter. However, at this order we can renormalize simply by subtracting the ultraviolet divergence in a particular scheme.

To isolate the ultraviolet part of the divergence, we regularize the infrared by taking p^2

[45] To be explicit, I should mention that the complex conjugate must be multiplied by γ^0 from the right.

[46] in d dimensions

[47] Strictly speaking, we *cannot* take $p^2 \to 0$ until we renormalize our matrix elements because this procedure leads to *unregularized* infrared divergences. Until our distribution is ultraviolet finite, we need $\epsilon > 0$ to regularize ultraviolet divergences. In this region, $\lim_{p^2\to 0}(p^2)^{-\epsilon/2}$ is certainly *not* zero. However, once the ultraviolet divergences have been removed, we are free to take $\epsilon < 0$ and $p^2 \to 0$ without ambiguity.

vanishingly small but finite.[48] This has the effect of postponing the cancelation of infrared and ultraviolet divergences until the ultraviolet divergence can be identified and removed. In addition to the direct subtraction, division by Z_F is necessary to properly normalize the external quark states.[49] In the $\overline{\rm MS}$ scheme, the full result is[50]

$$\begin{aligned}\mathcal{M}_{q'/q}(x) = \Bigg\{ & \delta(1-x) - \frac{\alpha_s C_F}{2\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \\ & \times \left[\frac{1+x^2}{1-x} - \delta(1-x)\int_0^1 dy \frac{1+y^2}{1-y}\right] \\ & + \mathcal{O}(\alpha_s^2)\Bigg\} \delta_{qq'} u_q(p,h)\overline{u}_q(p,h) \quad . \end{aligned} \tag{2.59}$$

Since $u(p,h)\overline{u}(p,h) = (1+2h\gamma_5)\,\not{p}/2$, we can split the density matrix into two scalar pieces :

$$\left(\mathcal{M}_{q'h'/qh}\right)_{\alpha\beta} = \frac{1}{2} f_{q'/q}(x)\,(\not{p})_{\alpha\beta} + \tilde{f}_{q'/q}(x)\,(\gamma_5 h\not{p})_{\alpha\beta} \quad ; \tag{2.60}$$

$$f_{q'/q}(x) = \frac{1}{2}\int \frac{d\lambda}{2\pi} e^{-i\lambda x}\left\langle ph\left|\overline{\psi}^{q'}(\lambda n)\,\mathcal{G}_n(\lambda n,0)\not{n}\,\psi^{q'}(0)\right|ph\right\rangle \tag{2.61}$$

$$\begin{aligned} = \delta_{qq'}\Bigg\{ & \delta(1-x) - \frac{\alpha_s C_F}{2\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \\ & \times \left[\frac{1+x^2}{1-x} - \delta(1-x)\int_0^1 dy \frac{1+y^2}{1-y}\right] + \mathcal{O}(\alpha_s^2)\Bigg\} \quad ; \end{aligned} \tag{2.62}$$

$$\tilde{f}_{q'/q}(x) = \frac{1}{4h}\int \frac{d\lambda}{2\pi} e^{-i\lambda x}\left\langle ph\left|\overline{\psi}^{q'}(\lambda n)\,\mathcal{G}_n(\lambda n,0)\not{n}\,\psi^{q'}(0)\right|ph\right\rangle \tag{2.63}$$

$$\begin{aligned} = \delta_{qq'}\Bigg\{ & \delta(1-x) - \frac{\alpha_s C_F}{2\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2} \\ & \times \left[\frac{1+x^2}{1-x} - \delta(1-x)\int_0^1 dy \frac{1+y^2}{1-y}\right] + \mathcal{O}(\alpha_s^2)\Bigg\} \quad . \end{aligned} \tag{2.64}$$

As mentioned above, the helicity-dependent scalar distributions can be projected out of

[48] This amounts to neglecting it everywhere except in the denominators.

[49] In renormalized perturbation theory, this factor is required to renormalize the external states; in unrenormalized perturbation theory, it renormalizes the quark fields.

[50] The divergent integral here will be discussed below. For now, we just treat it with the love and care it requires as an ill-defined mathematical object.

$\mathcal{M}$. Taking the target helicity positive, we have[51]

$$\begin{aligned} f_{q'\pm/q+} &= \frac{1}{2}\,\mathrm{Tr}\,\left[\frac{1}{2}(1\pm\gamma_5)\mathcal{M}\;\not{n}\right] \\ &= \frac{1}{2}\left(f_{q'/q}(x)\pm\tilde{f}_{q'/q}(x)\right)\;\; ; \end{aligned} \tag{2.65}$$

$$f_{q'/q}(x) = f_{q'+/q+} + f_{q'-/q+}\;\; , \tag{2.66}$$

$$\tilde{f}_{q'/q}(x) = f_{q'+/q+} - f_{q'-/q+}\;\; . \tag{2.67}$$

Hence $f_{q'/q}(x)$ represents the probability that a quark of flavor q' and momentum xp will be found in a quark of flavor q and momentum p regardless of its polarization, and $\tilde{f}_{q'/q}(x)$ represents the *difference* between the same helicity distribution and the opposite helicity distribution. Our result above indicates that at one-loop order the probability of finding a quark with opposite helicity is zero. This is in agreement with the fact that QCD is a vector theory, and as such does not change the helicity of its constituents. However, higher order radiative corrections can generate opposite helicity fermions through loops. This simplification is only an artifact of the low order at which we work.

Since QCD dynamics are independent of flavor,[52] quarks can only be generated in flavor-singlet pairs. Every contribution to $f_{q'/q}(x)$ must be countered by an equal contribution to an antiquark distribution function $f_{\overline{q}'/q}(y)$. This suggests that the integral of the difference between quark and antiquark probability distributions for any given flavor cannot be altered by QCD dynamics. This is, in fact, true. The integral

$$\int_0^1 dx\left[f_{q/T}(x) - f_{\overline{q}/T}(x)\right] \equiv N_{q/T} \tag{2.68}$$

is completely independent of QCD dynamics and returns an integer, the *net* number of quarks of flavor q, for every target T. Of course, in order to have a rigorous statement, we must first define the antiquark distribution. Writing[53]

$$\left(\overline{\mathcal{M}}_{\overline{q}'/q}\right)_{\alpha\beta}(x) = \int\frac{d\lambda}{2\pi}e^{-i\lambda x}\left\langle ph\left|\psi_\alpha^{q'}(\lambda n)\,\overline{\psi}_\beta^{q'}(0)\,\mathcal{G}_n(0,\lambda n)\right|ph\right\rangle \tag{2.69}$$

[51] The $\not{n}$ here appears simply because it allows us to project out a scalar distribution involving single-helicity fields. Alternatively, one can simply multiply $\mathcal{M}$ by $(1\pm\gamma_5)/2$ on the left and work out the contributing functions.

[52] assuming, as always, that all quarks are massless

[53] Note the position of the gauge link. Its indices are contracted as before.

$$= \frac{1}{2} f_{\overline{q}'/q}(x) \, (\not{p})_{\alpha\beta} + \tilde{f}_{\overline{q}'/q}(x) \, (\gamma_5 \, h \not{p})_{\alpha\beta} \quad , \tag{2.70}$$

in analogy with Eqs.(2.54) and (2.60), we see that this density matrix contains the antiquark probability distributions

$$\begin{aligned} f_{\overline{q}'\pm/q+} &= \frac{1}{2} \, \mathrm{Tr} \, \left[\frac{1}{2}(1 \pm \gamma_5) \overline{\mathcal{M}} \, \not{n} \right] \\ &= \frac{1}{2} \left(f_{\overline{q}'/q}(x) \pm \tilde{f}_{\overline{q}'/q}(x) \right) \qquad (2.71) \\ f_{\overline{q}'/q}(x) &= f_{\overline{q}'+/q+} + f_{\overline{q}'-/q+} \qquad (2.72) \\ \tilde{f}_{\overline{q}'/q}(x) &= f_{\overline{q}'+/q+} - f_{\overline{q}'-/q+} \quad . \qquad (2.73) \end{aligned}$$

Here, the helicity label does not correspond to the helicity of the antiquark itself. Rather, it refers to the helicity of the quark related to the antiquark. For example, $f_{\overline{q}'+/q+}(x)$ represents the probability of finding an antiquark of flavor q', momentum xp, and *negative* helicity in a quark of flavor q with momentum p and positive helicity.

Since quarks and antiquarks are so closely related, it is customary to combine their contributions into the same distribution. In light of Equation (2.69), the relationship

$$\mathcal{M}_{\alpha\beta}(-x) \sim -\overline{\mathcal{M}}_{\alpha\beta}(x) \tag{2.74}$$

is valid in the singular axial gauge, $n \cdot \mathcal{A} = 0$, provided that we ignore the anticommutator.[54] Since we have not at this point attributed any meaning to the functions $f_{q'/q}(x)$ on the negative real axis, we can *define*

$$\begin{aligned} f_{q/T}(-x) &\equiv -f_{\overline{q}/T}(x) \\ \tilde{f}_{q/T}(-x) &\equiv -\tilde{f}_{\overline{q}/T}(x) \end{aligned} \tag{2.75}$$

for $x > 0$. This is not in any way a statement about the analytic properties of $f_{q'/q}(x)$.[55] It is merely a convention to simplify the appearance of our equations and acknowledge the similarity between quarks and antiquarks. Using this notation, Eq.(2.68) can be written in the compact form

$$\int_{-1}^{1} dx \, f_{q/T}(x) = N_{q/T} \quad . \tag{2.76}$$

[54] It can be shown that this term vanishes in a particular quantization scheme. See [52].

[55] The operators which are able to generate *connected* contributions are quite different in the two cases.

The vector nature of QCD allows one to construct the sum rule

$$\int_{-1}^{1} dx\, f_{q\pm/T}(x) = N^{\pm}_{q/T} \tag{2.77}$$

using arguments identical to those that led to (2.76). These new quantities can be used to define the *net* number of positive and negative helicity quarks of each flavor in the target T. According to (2.65), this implies

$$N^{+}_{q/T} + N^{-}_{q/T} = \int_{-1}^{1} dx\, f_{q/T}(x) = \quad N_{q/T} \tag{2.78}$$

$$N^{+}_{q/T} - N^{-}_{q/T} = \int_{-1}^{1} dx\, \tilde{f}_{q/T}(x) = \Delta N_{q/T} \quad . \tag{2.79}$$

As mentioned above, $N_{q/T}$ is conserved separately for each flavor. This fact follows from the result[56]

$$\partial_\mu \langle \mathrm{T}\left(\overline{\psi}_q(x)\gamma^\mu \psi_q(x)\right)\rangle = 0 \tag{2.80}$$

for physical matrix elements.

The analogous relation for $\Delta N_{q/T}$ is broken by quantum effects [53]. However, these effects are the same for each species of quark. This makes the difference $\Delta N_{q'/T} - \Delta N_{q/T}$ independent of QCD dynamics. Systematically, one can define the nonsinglet distributions

$$f^{i}_{NS/T}(x) \quad \equiv \quad \sum_{j=1}^{i} f_{j/T}(x) - i f_{i+1/T} \tag{2.81}$$

$$\tilde{f}^{i}_{NS/T}(x) \quad \equiv \quad \sum_{j=1}^{i} \tilde{f}_{j/T}(x) - i \tilde{f}_{i+1/T} \quad . \tag{2.82}$$

In a theory of n_f approximately massless quarks, the $n_f - 1$ nonsinglet distributions represent purely quark effects. These distributions are immune to the dynamical anomaly that alters $\Delta N_{q/T}$, so the objects

$$\Delta N^{i}_{NS/T} \equiv \int_{-1}^{1} dx\, \tilde{f}^{i}_{NS/T}(x) \tag{2.83}$$

are physical quantities which are conserved to all orders of perturbation theory.[57] The orthog-

[56] This identity is proved in Appendix D.2; it follows from QED gauge invariance.

[57] This still doesn't put them on the same level as the $N_{q/T}$. These quantities are conserved separately regardless of quark mass effects, while conservation of the nonsinglet ΔN's relies heavily on the assumption of massless quarks. Even $N_{q/T}$ is altered by the flavor-changing weak interactions. Only the singlet quantity $N_{S/T}$ corresponding to the total number of quarks in the target remains inviolate in the full theory.

onal degree of freedom is composed of the sum of all flavors,

$$f_{S/T}(x) \equiv \sum_{j=1}^{n_f} f_{j/T}(x) \tag{2.84}$$

$$\tilde{f}_{S/T}(x) \equiv \sum_{j=1}^{n_f} \tilde{f}_{j/T}(x) \ . \tag{2.85}$$

This singlet[58] distribution is not immune to the effects of the dynamical anomaly that alters ΔN. This implies $\Delta N_{S/T}$, which represents the difference between the number of positive and negative helicity quarks irrespective of flavor in the target, is dynamically dependent. This manifests itself in a dependence on renormalization scale. Since it is not loyal to any particular flavor, the effects of this distribution cannot be distinguished from those of the gluon field. As we will see, this fact is intimately related to the presence of infrared divergences in the gluon amplitude.

The soft physics relevant to gluon contributions to DIS comes in two different forms. On one hand, gluons can fluctuate into quark-antiquark pairs which then participate in the hard scattering. In this case, the fluctuation occurs long before the hard scattering and cannot truly be accounted for perturbatively. This leads to a term of the form $f_{q/g}\, t_q^{\mu\nu}$ in $T_g^{\mu\nu}$. The distributions relevant to this contribution are contained in Equation (2.54) and its anti-counterpart (2.69), with gluonic external states. Defining[59]

$$\mathcal{M}_{q/g\eta}(x) = \frac{1}{2} f_{q/g}(x)\ \not{p} + \frac{1}{2}\tilde{f}_{q/g}(x)\gamma_5(\eta \not{p}) \tag{2.86}$$

$$\overline{\mathcal{M}}_{\overline{q}/g\eta}(x) = \frac{1}{2} f_{\overline{q}/g}(x)\ \not{p} + \frac{1}{2}\tilde{f}_{\overline{q}/g}(x)\gamma_5(\eta \not{p}) \ , \tag{2.87}$$

where $\eta = \pm 1$ is the gluon helicity, we have

$$f_{q/g}(x) = -\frac{\alpha_s T_F}{2\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[x^2 + (1-x)^2\right] \tag{2.88}$$

$$\tilde{f}_{q/g}(x) = -\frac{\alpha_s T_F}{2\pi}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[x^2 - (1-x)^2\right] \tag{2.89}$$

$$f_{\overline{q}/g}(x) = f_{q/g}(x) \tag{2.90}$$

[58] 'Singlet' and 'nonsinglet' refer to the transformation properties under the flavor group, $SU(n_f)$. Since the gluon fields are invariant under the action of this group and the nonsinglet distributions are not, these two sectors have no business with each other. The singlet distribution, on the other hand...

[59] Note the extra factor of 1/2 accompanying $\tilde{f}$. This compensates for the difference in spin between gluon and quark.

$$\tilde{f}_{\overline{q}/g}(x) \quad = \quad -\tilde{f}_{q/g}(x) \ . \tag{2.91}$$

On the other hand, gluons can enjoy their own hard scattering coefficient. Here, the fluctuation into a quark-antiquark pair occurs on the same time scale as the hard interaction, creating an effective photon-gluon coupling. This contribution will have the form $f_{g/g}\, t_g^{\mu\nu}$. The parton distribution which appears here must be expressed as a matrix element of a gluonic operator. The road from idea to matrix element is complicated in this case by the nonlinear transformation law for the gluonic fields. Here, we need somewhat more than a gauge link to form a gauge invariant distribution. In fact, the only local object which annihilates the one-gluon state and has linear gauge transformation properties is the field strength $\mathcal{F}^{\mu\nu}$. This implies that the matrix element

$$\begin{aligned} G_{g/a}^{\mu\nu\alpha\beta}(x) &= \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle p\eta \left| \mathcal{F}^{\mu\nu}\left(\lambda n\right) \mathcal{G}_n\left(\lambda n, 0\right) \mathcal{F}^{\alpha\beta}\left(0\right) \right| p\eta \right\rangle \\ &= -x f_{g/a}(x) \left(g^{\mu\alpha} p^\nu p^\beta - g^{\mu\beta} p^\nu p^\alpha + g^{\nu\beta} p^\mu p^\alpha - g^{\nu\alpha} p^\mu p^\beta \right) \\ &\quad + i\eta x \tilde{f}_{g/a}(x) \left(\epsilon^{\mu\alpha\gamma\delta} p^\nu p^\beta - \epsilon^{\mu\beta\gamma\delta} p^\nu p^\alpha + \epsilon^{\nu\beta\gamma\delta} p^\mu p^\alpha - \epsilon^{\nu\alpha\gamma\delta} p^\mu p^\beta \right) n_\gamma p_\delta \ , \end{aligned} \tag{2.92}$$

contains the vacuum as an intermediate state. $\mathcal{G}$ is the same gauge link as that in the quark distribution operator, with the generators T^a appearing in the gluon fields interpreted in the adjoint representation.[60] The normalization has been chosen so that the scalar distributions,[61]

$$f_{g/a}(x) = -\frac{1}{2x} \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle p\eta | \mathcal{F}^{\mu\alpha}\left(\lambda n\right) \mathcal{G}_n\left(\lambda n, 0\right) \mathcal{F}^{\nu}{}_{\alpha}\left(0\right) | p\eta \rangle n_\mu n_\nu \ , \tag{2.93}$$

$$\tilde{f}_{g/a}(x) = -\frac{i}{2\eta x} \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle p\eta | \mathcal{F}^{\mu\alpha}\left(\lambda n\right) \mathcal{G}_n\left(\lambda n, 0\right) \tilde{\mathcal{F}}^{\nu}{}_{\alpha}\left(0\right) | p\eta \rangle n_\mu n_\nu \ , \tag{2.94}$$

which can be isolated by contraction, have the leading behavior

$$\begin{aligned} f_{g/g}(x) &= \frac{x}{2} \left[\delta(1-x) + \delta(1+x) \right] + \mathcal{O}(\alpha_s) \\ \tilde{f}_{g/g}(x) &= \frac{x}{2} \left[\delta(1-x) - \delta(1+x) \right] + \mathcal{O}(\alpha_s) \ . \end{aligned} \tag{2.95}$$

As before, These distributions can be calculated in perturbation theory by introducing new fields with eikonal propagation and inserting a complete set of states.[62] The physical polarization vectors for gluons moving in the 3-direction are given by $\varepsilon_\eta^\mu = (0, 1, i\eta, 0)/\sqrt{2}$.

[60] See Appendix C. These indices have been suppressed as in the quark distribution operators.

[61] The dual field strength is defined by $\tilde{\mathcal{F}}^{\mu\nu} \equiv \frac{1}{2}\, \epsilon^{\mu\nu\alpha\beta} \mathcal{F}_{\alpha\beta}$.

[62] Since gluons are bosons, the creation and annihilation operators in each gluon field are of the same species.

Since this distribution appears only with $t_g^{\mu\nu}$, a quantity that starts with α_s, we do not need the next-to-leading behavior in our calculation. This also implies that we need not calculate the 'gluon distribution in a quark', $f_{g/q}$, since this object can only contribute to $T_q^{\mu\nu}$ at order α_s^2.

To see the connection between the above distributions and the idea of removing one gluon, we turn again to the singular light-cone gauge. As before, the gauge link is trivial in this gauge. Furthermore, this choice reduces $n_\alpha \mathcal{F}^{\alpha\mu}$ to $n \cdot \partial \mathcal{A}^\mu$. Introducing a complete set of intermediate states, we see that $f_{g/a}(x)$ becomes[63]

$$\begin{aligned} f_{g/a}(x) &= -\frac{1}{2x}\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \sum_X \langle ph\,|in\cdot\partial \mathcal{A}_a^\alpha(\lambda n)|\,X\rangle\,\langle X\,|in\cdot\partial \mathcal{A}_\alpha^a(0)|\,ph\rangle \\ &= x\sum_X \delta\left(p_X\cdot n-(1-x)\right)\langle ph\,|\mathcal{A}_a^\alpha(0)|\,X\rangle\,\left\langle X\left|\mathcal{A}_a^\beta(0)\right|\,ph\right\rangle\left(-\frac{g_{\alpha\beta}}{2}\right) \end{aligned} \tag{2.96}$$

Aside from the factor of x, this is exactly what we would have written down for the unpolarized gluon distribution if we did not have to worry about gauge invariance.[64] However, a proper normalization must take the issue of gauge invariance into account. If we wish to normalize all of our distributions in an unambiguous way, it is necessary to relate them to a physical observable. The conventional choice is the total + momentum carried by a certain parton in our target. Interpreting the parton distributions as probabilities, we would conclude that this object is simply a sum over the probabilities weighted by the momentum carried by each parton :

$$n\cdot P_{a/T} = \int_0^1 dx\, x f_{a/T}(x) \ . \tag{2.97}$$

On the other hand, the operator representing the momentum carried by various parton species can be obtained from the stress-energy tensor of QCD. According to the analysis of Appendix

This leads to the $\delta(1+x)$ terms above, which do not appear in the fermion case because the action of antiquark creation and annihilation operators cannot generate connected diagrams at this order. Note that this bose symmetry can be taken into account simply by requiring (anti)symmetry in the argument x for the (un)polarized distribution. This is a general property of the matrix element (2.92).

[63] neglecting the bose symmetry

[64] The factor $-g_{\alpha\beta}/2$ averages over transverse gluon polarizations with the weight +1; remember that in axial gauge $\mathcal{A}\cdot\mathcal{A} = -\vec{\mathcal{A}}_\perp\cdot\vec{\mathcal{A}}_\perp$.

D.2, we have

$$\begin{aligned}
\hat{P}^\mu &= \int d^3x\, \Theta^{0\mu}(x) && (2.98)\\
\Theta^{\mu\nu} &= \Theta_q^{\mu\nu} + \Theta_g^{\mu\nu} && (2.99)\\
\Theta_q^{\mu\nu} &= \overline{\psi}\gamma^{(\mu}\, i\overleftrightarrow{\mathcal{D}}^{\ \nu)}\psi && (2.100)\\
\Theta_g^{\mu\nu} &= \frac{1}{4}g^{\mu\nu}\mathcal{F}^2 - \mathcal{F}^{\mu\alpha}\mathcal{F}^{\nu}{}_{\alpha}\ . && (2.101)
\end{aligned}$$

Lorentz invariance implies

$$\langle P|\,\Theta_{q,g}^{\mu\nu}\,|P\rangle = 2a_{q,g}P^\mu P^\nu + 2b_{q,g}M^2 g^{\mu\nu}\ . \qquad (2.102)$$

Since $|P\rangle$ is an eigenstate of $\hat{P}^\mu$, we have[65]

$$\begin{aligned}
P^\mu\left(2P^0\right)(2\pi)^3\,\delta^{(3)}(0) &= \langle P|\int d^3x\,\Theta^{0\mu}(x)\,|P\rangle\\
&= \langle P|\,\Theta^{0\mu}(0)\,|P\rangle\,(2\pi)^3\,\delta^{(3)}(0)\ . && (2.103)
\end{aligned}$$

This immediately tells us that

$$a_q + a_g = 1 \qquad\qquad b_q + b_g = 0\ , \qquad (2.104)$$

which promotes the identification of $a_{q,g}$ with the fraction of momentum carried by quarks and gluons, respectively.[66] These parameters are pure numbers[67] and may be obtained in any way we wish. In particular, from Equation (2.102), we see that

$$a_{q,g} = \frac{1}{2}\,\langle P|\,\Theta_{q,g}^{++}\,|P\rangle\ . \qquad (2.105)$$

Our normalization condition, Eq.(2.97), then takes the convenient form

$$\frac{1}{2}\,\langle P|\,\Theta_{q,g}^{++}\,|P\rangle = \int dx\, x f_{q,g/T}(x)\ . \qquad (2.106)$$

[65] Our states are normalized so that $\langle P'|P\rangle = 2P^0(2\pi)^3\delta(\vec{P}' - \vec{P}\,)$.

[66] The quantity $b_{q,g}$ should not bother us too much here. Its presence can be understood in the rest frame of the nucleon, where Equation (2.102) would otherwise imply that no species of parton can have nontrivial momentum dynamics. These kinds of transverse effects are suppressed by powers of Q^2 in the Bjorken limit; in any case, the contribution of b to the + momentum of the proton for any appreciable value of the boost parameter Λ is quite negligible.

[67] ignoring, for the moment, their dependence on renormalization scale

It is now easy to see why the 'extra' factor of x adorning the gluon distributions is needed.[68]

Although the correctly normalized gauge-invariant distributions are given by $f_{g/T}$ and $\tilde{f}_{g/T}$, our perturbative calculation in the last section was performed with single gluon external states. For these states, distributions of the form in Eq.(2.96) are required. However, it is the proper distributions which will survive the transition from parton external states to physical hadronic ones. We can account for the difference between these two distributions simply by dividing our perturbative results by x since

$$\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle PS | \mathcal{A}_\perp^\mu(\lambda n) \mathcal{A}_\perp^\nu(0) | PS \rangle = \left[-\frac{2}{d-2} \frac{1}{x} g_\perp^{\mu\nu} f_{g/T}(x) + \frac{i}{x} \epsilon^{-+\mu\nu} \tilde{f}_{g/T}(x) \right] \quad (2.108)$$

in the light-cone gauge.

Armed with perturbative expressions for our parton densities, we are now ready to extract the hard scattering coefficients. Exploiting the fact that we work only to next-to-leading order, we write

$$\begin{aligned} T_q^{\mu\nu} &= \left(T_q^{\mu\nu}\right)^{(0)} + \frac{\alpha_s C_F}{4\pi} \left[\frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \left(T_q^{\mu\nu}\right)_{\rm IR}^{(1)} + \left(T_q^{\mu\nu}\right)^{(1)} \right] \\ f_{q'/q}(x) &= f_{q'/q}^{(0)}(x) + \frac{\alpha_s C_F}{4\pi} \frac{2}{\epsilon} \left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} f_{q'/q}^{(1)}(x) \quad (2.109) \\ t_{q'}^{\mu\nu} &= (t_{q'}^{\mu\nu})^{(0)} + \frac{\alpha_s C_F}{4\pi} (t_{q'}^{\mu\nu})^{(1)} \quad (2.110) \end{aligned}$$

for the quark amplitude. The infrared divergent term in $T_q^{\mu\nu}$ has been designed to include only the infrared divergence as seen in the $\overline{\rm MS}$ scheme. For example, the term

$$g^{\mu\nu} e_q^2 \frac{\alpha_s C_F}{4\pi} \frac{3}{x_B - 1} \frac{2}{\epsilon} \left(\frac{Q^2 e^{\gamma_E}}{4\pi\mu_0^2} \right)^{-\epsilon/2} \quad (2.111)$$

in $T^{\mu\nu}$ contributes

$$g^{\mu\nu} e_q^2 \frac{3}{x_B - 1} \quad (2.112)$$

[68] A somewhat easier, but less rigorous way to understand this factor comes from a different analogy with the quark case. Above, we have interpreted $\overline{\psi}$ as the hermitian conjugate to ψ. If instead we think of it as the momentum canonically conjugate to ψ, the the analogy with the gluon sector changes its nature completely. The canonically conjugate momentum to the gluon field is $\mathcal{F}$, so we would naively write

$$\langle \mathcal{F}^{+\alpha} \mathcal{A}_\alpha \rangle \quad (2.107)$$

for the (unpolarized) gluon distribution. In order to make this gauge-invariant, we must eliminate $\mathcal{A}$ in favor of $\mathcal{F}$. The 'extra' factor of x compensates for the necessary derivative.

to $\left(T_q^{\mu\nu}\right)^{(1)}_{\rm IR}$ and

$$- g^{\mu\nu} e_q^2 \, \frac{3}{x_B - 1} \, \log \frac{Q^2}{\mu^2} \tag{2.113}$$

to $\left(T_q^{\mu\nu}\right)^{(1)}$. Note that we are considering scattering on a quark of momentum p rather than xp here. The perturbative coefficient functions are designed not to care.

Equating terms order by order in α_s, we see that Equation (2.52) implies

$$\left(T_q^{\mu\nu}\right)^{(0)} = \sum_{q'} \int_{-1}^{1} dx \, f^{(0)}_{q'/q}(x) (t^{\mu\nu}_{q'})^{(0)}(x) \tag{2.114}$$

$$\left(T_q^{\mu\nu}\right)^{(1)} = \sum_{q'} \int_{-1}^{1} dx \, f^{(0)}_{q'/q}(x) (t^{\mu\nu}_{q'})^{(1)}(x) \tag{2.115}$$

$$\left(T_q^{\mu\nu}\right)^{(1)}_{\rm IR} = \sum_{q'} \int_{-1}^{1} dx \, f^{(1)}_{q'/q}(x) (t^{\mu\nu}_{q'})^{(0)}(x) \ . \tag{2.116}$$

Equations (2.114) and (2.115) are used to define $(t^{\mu\nu}_{q'})^{(0)}$ and $(t^{\mu\nu}_{q'})^{(1)}$. Equation (2.116) is a statement about the sources of soft physics in our calculation. If it is satisfied, all of the soft physics relevant to DIS is contained within the parton distribution functions. Otherwise, other sources of soft physics contribute to this process. This situation could imply that one cannot separate the hard and soft scales. Fortunately, this is not the case in inclusive DIS. Explicit substitution of the expressions above into (2.114), (2.115), and (2.116) and analogous relations in the gluon sector, shows that (2.116) is satisfied for polarized and unpolarized DIS on quark and gluon targets at order α_s. The results can be expressed for any target T in the simple form[69]

$$\begin{aligned} T_1(x_B) &= \int_{-1}^{1} \frac{dx}{x} \left[\sum_q f_{q/T}(x,\mu^2) C_q\left(\frac{x}{x_B}, \frac{Q^2}{\mu^2}\right) + f_{g/T}(x,\mu^2) C_g\left(\frac{x}{x_B}, \frac{Q^2}{\mu^2}\right) \right] \\ S_1(x_B) &= \int_{-1}^{1} \frac{dx}{x} \left[\sum_q \tilde{f}_{q/T}(x,\mu^2) \tilde{C}_q\left(\frac{x}{x_B}, \frac{Q^2}{\mu^2}\right) + \tilde{f}_{g/T}(x,\mu^2) \tilde{C}_g\left(\frac{x}{x_B}, \frac{Q^2}{\mu^2}\right) \right] \ , \end{aligned} \tag{2.117}$$

where[70]

$$C_q\left(\frac{x}{x_B}, \frac{Q^2}{\mu^2}\right) \quad = \quad e_q^2 \frac{x}{x_B - x} - \frac{\alpha_s(\mu^2) C_F}{4\pi} e_q^2 \left\{ \left(3 - \log\frac{Q^2}{\mu^2}\right) \frac{3x}{x_B - x} - 2 \log\frac{Q^2}{\mu^2} \right.$$

[69] The leading antiquark scattering amplitude is just the negative of the leading quark contribution evaluated at $-x$, so these contributions can be combined into the same hard scattering coefficient.

[70] Note the extra factors of $1/x$ in the gluonic coefficient functions coming from the difference in definition between the properly normalized gluon distributions and the distributions relevant to our calculated amplitudes.

$$+\left[\left(\frac{x_B+x}{x}+\frac{2x}{x_B-x}\right)\left(3-\log\frac{x_B-x}{x_B}-2\log\frac{Q^2}{\mu^2}\right)\right.$$
$$\left.-3\left(\frac{x_B-x}{x}+\frac{x}{x_B-x}\right)\right]\log\frac{x_B-x}{x_B}\Bigg\} \qquad (2.118)$$
$$+(x_B\to -x_B)\ ,$$

$$\tilde{C}_q\left(\frac{x}{x_B},\frac{Q^2}{\mu^2}\right) = e_q^2\frac{x}{x_B-x}-\frac{\alpha_s(\mu^2)C_F}{4\pi}e_q^2\left\{\left(3-\log\frac{Q^2}{\mu^2}\right)\frac{3x}{x_B-x}\right.$$
$$+\left[\left(\frac{x_B+x}{x}+\frac{2x}{x_B-x}\right)\left(3-\log\frac{x_B-x}{x_B}-2\log\frac{Q^2}{\mu^2}\right)\right.$$
$$\left.+7\,\frac{x_B-x}{x}-3\,\frac{x}{x_B-x}\right]\log\frac{x_B-x}{x_B}\Bigg\} \qquad (2.119)$$
$$-(x_B\to -x_B)\ ,$$

$$C_g\left(\frac{x}{x_B},\frac{Q^2}{\mu^2}\right) = \frac{\alpha_s(\mu^2)T_f}{2\pi}\left(\sum_q e_q^2\right)\left\{2\log\frac{Q^2}{\mu^2}-2\right.$$
$$+\left[\left(1+2x_B\frac{x_B-x}{x^2}\right)\left(4-\log\frac{x_B-x}{x_B}-2\log\frac{Q^2}{\mu^2}\right)\right.$$
$$\left.\left.-2\right]\log\frac{x_B-x}{x_B}\right\}+(x_B\to -x_B) \qquad (2.120)$$

$$\tilde{C}_g\left(\frac{x}{x_B},\frac{Q^2}{\mu^2}\right) = \frac{\alpha_s(\mu^2)T_f}{2\pi}\left(\sum_q e_q^2\right)$$
$$\times\left\{\left[\left(1+2\frac{x_B-x}{x}\right)\left(4-\log\frac{x_B-x}{x_B}-2\log\frac{Q^2}{\mu^2}\right)\right.\right.$$
$$\left.\left.-2\right]\log\frac{x_B-x}{x_B}\right\}-(x_B\to -x_B)\ . \qquad (2.121)$$

Renormalization has introduced violations of the simple scaling behavior we saw in the leading order amplitudes. The dependence on renormalization scale is illusory; the coefficient functions and parton distributions depend separately on μ in such a way that the product is independent. However, the dependence of our amplitude on Q^2 is quite real and can be measured in experiment. In the absence of asymptotic freedom, this behavior would persist even into the Bjorken region so scaling could never be experimentally observed. Asymptotic freedom ensures that scaling violations become smaller as Q^2 is increased.

Since these scattering coefficients are completely insensitive to the soft infrared physics, (2.117) is valid for *all* targets. In particular, we can derive expressions for the quantities measured in real DIS on proton targets simply by taking the imaginary part of T_1 and S_1 as x_B approaches the real axis from below.[71] The parton distribution functions are real, so the

[71] At this point, we must analytically continue our expressions to the region $|\,x_B\,|\leq 1$. It is only in this

measured structure functions are expressed as convolutions of the unknown structure functions with known scattering kernels. At leading order, we have simply[72]

$$F_1(\nu, Q^2) = \frac{1}{2}\sum_q e_q^2 \left[f_{q/T}(x_B, Q^2) + f_{\overline{q}/T}(x_B, Q^2) \right] \tag{2.122}$$

$$G_1(\nu, Q^2) = \frac{1}{2}\sum_q e_q^2 \left[\tilde{f}_{q/T}(x_B, Q^2) - \tilde{f}_{\overline{q}/T}(x_B, Q^2) \right] \quad . \tag{2.123}$$

The interpretation of this result is clear. At leading order, the photon simply couples to each parton completely incoherently. F_1 measures the total number of charge carriers with momentum $x_B P$ weighted by the square of the charge. Notice that the *sum* of the quark and antiquark distributions is relevant here, rather than the difference, since both contribute to DIS in the same way. Likewise, G_1 measures the helicity of charged carriers weighted by squared charge. Since the antiquark distribution has been defined in such a way that $\tilde{f}_{\overline{q}/T}$ measures the net *negative* helicity of antiquarks, it appears with a sign in this expression. The kinematical restriction $x_B = x$ corresponds to the requirement that the intermediate quark be 'physical'.

A measurement of the structure functions allows us to extract the parton distributions and obtain invaluable information on nonperturbative hadronic structure. In particular, integrals of the structure functions can tell us how much of a proton's momentum is carried by quarks or the total spin carried by quark helicity in tritium. The results of this section allow us to go even further and predict the leading QCD corrections to F_1 and G_1. All that is needed is the imaginary parts of the coefficient functions, Eqs.(2.118), (2.119), (2.120) and (2.121).[73]

region that the correspondence between our amplitude and physical measurements can be made.

[72] We choose $\mu^2 = Q^2$ because this is the most natural value for it. The full expression is indeed independent - if we calculate to all orders. Our lack of motivation for this task forces us to choose μ^2 in such a way that corrections are minimized.

[73] Taking the imaginary part of the logarithms in these expressions is somewhat tricky. For $x < x_B$,

$$\begin{aligned} \Re e \log \frac{x_B - x}{x_B} &= \log \frac{x_B - x}{x_B} \\ \Im m \log \frac{x_B - x}{x_B} &= 0 \ , \end{aligned} \tag{2.124}$$

while for $x > x_B$,

$$\begin{aligned} \Re e \log \frac{x_B - x}{x_B} &= \log \frac{x - x_B}{x_B} \\ \Im m \log \frac{x_B - x}{x_B} &= -\pi \ . \end{aligned} \tag{2.125}$$

Once the parton distributions are known at a certain scale, their scale evolution is fixed by perturbative QCD. This makes the study of scaling violations an especially clean way to view QCD. In fact, the prediction of these violations in DIS is one of the most precision tests of the theory.

The scale evolution of our distribution functions is interesting in its own right. Physically, it arises for the same reason that $\alpha_s(\mu^2)$ runs - things look different at different scales. From very far away, I may think that I am looking at a quark of momentum p. However, if I look closer by increasing my probing scale, I may find that that quark is really a quark of momentum xp and a gluon of momentum $(1-x)p$. This line of reasoning leads to the evolution equation

$$\mu^2 \frac{D_a}{D\,\mu^2} f_{a/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi} \sum_{a'} \int_0^1 dy f_{a'/T}(y,\mu^2) \int_0^1 dz P_{a'\to a}(z)\delta(x-zy) \quad (2.126)$$

$$= \frac{\alpha_s(\mu^2)}{2\pi} \sum_{a'} \int_x^1 \frac{dy}{y} P_{a'\to a}\left(\frac{x}{y}\right) f_{a'/T}(y,\mu^2) \ . \quad (2.127)$$

Here, $\alpha_s P_{a'\to a}(x) d\mu^2/2\pi\mu^2$ represents the probability that a parton of type a' and momentum p will generate a parton of type a and momentum xp due to the infinitesimal change $d\mu^2$ in scale. This activity will add to the distribution $f_{a/T}(x)$. The 'covariant' derivative, D, is designed to include the change in $f_{a/T}(x)$ due to partons of type a splitting into other partons. This is known as the endpoint contribution. Conceptually, it would seem that this term is just be some kind of sum over probabilities $P_{a\to a'}$. However, classical reasoning does not work here. In reality, we do not have conservation of particles in a quantum field theory. The *only* conserved quantities here are those quantum numbers associated with some symmetry of our lagrangian. In this case, the endpoint contributions are fixed by flavor and momentum conservation. As mentioned above,[74] conservation of flavor implies that

$$\int_0^1 dx \ \left[f_{q/T}(x,\mu^2) - f_{\overline{q}/T}(x,\mu^2)\right] \quad (2.128)$$

is independent of μ^2. Using

$$\mu^2 \frac{D_a}{D\mu^2} \equiv \mu^2 \frac{d}{d\mu^2} + \frac{\alpha_s(\mu^2)}{2\pi} A_a \ , \quad (2.129)$$

$-i\pi$ appears rather than $i\pi$ because for x_B slightly below the real axis, $x_B - x$ is a negative number with a negative imaginary part. Taking the branch cut along the negative real axis, we arrive at (2.125).

[74] At this level, it is conceptually easier to use the explicit forms of the quark and antiquark distributions. We can always simplify our expressions later using (2.75).

we see that this requirement leads to[75]

$$A_q = \int_0^x \frac{dy}{x} \left[P_{q\to q}(y/x) - P_{\overline{q}\to q}(y/x) \right] \quad . \tag{2.130}$$

Some of the relations

$$\begin{aligned}
P_{q\to q'}(x) &= P_{\overline{q}\to\overline{q}'}(x) && (2.131)\\
P_{q\to\overline{q}'}(x) &= P_{\overline{q}\to q'}(x) && (2.132)\\
P_{q\to g}(x) &= P_{\overline{q}\to g}(x) && (2.133)\\
P_{g\to q}(x) &= P_{g\to\overline{q}}(x) && (2.134)\\
(1-\delta_{qq'})P_{q\to q'}(x) &= (1-\delta_{qq'})P_{q\to\overline{q}'}(x) && (2.135)\\
&= (1-\delta_{qq'})P_{q\to\overline{q}}(x) && (2.136)\\
P_{q\to g}(x) &= P_{q'\to g}(x) && (2.137)\\
P_{g\to q}(x) &= P_{g\to q'}(x) && (2.138)\\
P_{q\to q}(x) &= P_{q'\to q'}(x) && (2.139)
\end{aligned}$$

are necessary to derive this result. All follow from the CP and flavor invariance of the QCD lagrangian and are valid to all orders in perturbation theory.[76] At the order to which we work, $P_{q\to\overline{q}}$ is zero. Hence conservation of flavor requires

$$A_q = \int_0^x \frac{dy}{x} P_{q\to q}\left(\frac{y}{x}\right) \quad . \tag{2.140}$$

On the other hand, we have calculated $f_{q'/q}(x,\mu^2)$ in perturbation theory. Manually taking the derivative of Eq.(2.62), we arrive at

$$\begin{aligned}
\mu^2\frac{d}{d\mu^2}f_{q/q}(x,\mu^2) &= \frac{\alpha_s(\mu^2)C_F}{2\pi}\left[\frac{1+x^2}{1-x} - \delta(1-x)\int_0^1 dy\,\frac{1+y^2}{1-y}\right]\\
&= \frac{\alpha_s(\mu^2)}{2\pi}\int_0^1 dydz\left(C_F\,\frac{1+z^2}{1-z}\right)f_{q/q}(y,\mu^2)\delta(x-zy) \qquad (2.141)\\
&\quad -\frac{\alpha_s(\mu^2)}{2\pi}\int_0^x \frac{dy}{x^2}\,C_F\,\frac{x^2+y^2}{x-y}f_{q/q}(x,\mu^2)
\end{aligned}$$

[75] In these kinds of manipulations, it is often easier to use the primordial form of the evolution equation, (2.126), rather than the streamlined (2.127). Note that we must be very careful deriving this result since the integrals involved are singular. In particular, the form of $P_{q\to q}(z)$ at leading order in $\alpha_s(\mu^2)$ leads to an anomalous dependence of A_q on x.

[76] Quark mass effects break flavor invariance. As such, they will alter relations (2.136), (2.137), (2.138), and (2.139). However, this will not change (2.130).

at leading order in $\alpha_s(\mu^2)$. Comparing with (2.126), we see that

$$P_{q\to q}(x) = C_F \, \frac{1+x^2}{1-x} \quad , \tag{2.142}$$

$$A_q = C_F \int_0^x \frac{dy}{x^2} \, \frac{x^2+y^2}{x-y} \tag{2.143}$$

at leading order in α_s. In light of (2.130), these two quantities are not independent. Since $P_{q\to q}(x)$ comes entirely from gluon intermediate states, the calculation of vacuum intermediate states is in some sense redundant. In the light-cone gauge, the absence of the gauge link causes the diagrammatic calculation to be incomplete. In order to obtain the full result in this gauge, one must *impose* (2.130).[77]

We are now in a position to understand the divergent integral in A_q. Its presence it required to *cancel* the divergent behavior of $\int P_{q\to q}$. To display this explicitly, one often combines the two terms :

$$\begin{aligned} \mu^2 \frac{d}{d\mu^2} f_{a/T}(x,\mu^2) &= \sum_{a'} \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \left[P_{a'\to a}\left(\frac{x}{y}\right) \right. \\ &\qquad \left. - A_a \delta_{aa'} \delta\left(1-\frac{x}{y}\right) \right] \, f_{a'/T}(y,\mu^2) \\ &= \sum_{a'} \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \, \mathcal{P}_{a'\to a}\left(\frac{x}{y}\right) f_{a'\to a}(y,\mu^2) \quad . \end{aligned} \tag{2.144}$$

In the quark case, we can write explicitly

$$\mathcal{P}_{q\to q}(x) = C_F \left[\frac{2}{(1-x)^+} - x - 1 + \frac{3}{2}\delta(1-x) \right] \quad . \tag{2.145}$$

The '+ function', $1/(1-x)^+$, is defined by

$$\frac{1}{(1-x)^+} \equiv \lim_{\varepsilon\to 0} \left[\frac{1}{1-x}\Theta(1-x-\varepsilon) - \delta(1-x) \int_0^{1-\varepsilon} dy \, \frac{1}{1-y} \right] \quad . \tag{2.146}$$

Its definition implies

$$\int_0^1 dx \, \frac{f(x)}{(1-x)^+} = \int_0^1 dx \, \frac{f(x)-f(1)}{1-x} \quad , \tag{2.147}$$

which is more useful in practice.

[77] This is true for a well-defined, consistent light-cone gauge choice. Due to the similarity between the pole prescription indicated in Fig. 2.4 and the PV-prescription, the pathological theory reproduces the correct end-point contribution at this order while ML does not. However, this does not reconcile our renormalizability issues with PV.

The other conservation law which must be satisfied by these distributions is that of + momentum. We have normalized our distributions in such a way that

$$\sum_a \int_0^1 dx\ x\ f_{a/T}(x,\mu^2) = n \cdot P_T \tag{2.148}$$

is the total + momentum of the target. Since this cannot be scale dependent, we have the relation

$$\sum_a \int_0^1 x\,dx\ A_a f_{a/T}(x,\mu^2) = \sum_a \sum_{a'} \left(\int_0^1 z\,dz\ P_{a'\to a}(z) \right) \left(\int_0^1 y\,dy\ f_{a'/T}(y,\mu^2) \right) \quad . \tag{2.149}$$

Using the above relations once more, along with the fact that $A_q = A_{\overline{q}}$ is independent of flavor and of *target*, we arrive at

$$A_g = \int_0^x \frac{y}{x}\frac{dy}{x} \left[2n_f P_{g\to q}\left(\frac{y}{x}\right) + P_{g\to g}\left(\frac{y}{x}\right) \right] \tag{2.150}$$

$$A_q = \int_0^x \frac{y}{x}\frac{dy}{x} \left[P_{q\to q}\left(\frac{y}{x}\right) + (2n_f - 1)\, P_{q\to \overline{q}}\left(\frac{y}{x}\right) + P_{q\to g}\left(\frac{y}{x}\right) \right] \quad . \tag{2.151}$$

The first of these relations allows us to deduce A_g without the use of vacuum insertions (or provides a useful check of our calculation), while the second takes the form of an unexpectedly strict consistency condition on the allowed probabilities for quark splitting. This equation can be used as an extremely strong check of one's expressions.

Our perturbative calculations of the parton distributions allow us to extract $P_{g\to q}(x)$ explicitly by differentiating $f_{q/g}(x)$. The result,

$$P_{g\to q}(x) = T_F \left[x^2 + (1-x)^2 \right] \quad , \tag{2.152}$$

allows us to write the full evolution equation

$$\mu^2 \frac{D_q}{D\mu^2} f_{q/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \left[P_{q\to q}\left(x/y\right) f_{q/T}(y,\mu^2) + P_{g\to q}\left(x/y\right) f_{g/T}(y,\mu^2) \right] \quad . \tag{2.153}$$

This equation is complicated by the presence of the gluon distribution.[78] As before, we can form nonsinglet distributions which decouple from the gluon sector using the diagonal genera-

[78] Note that the gluon distribution here has the leading behavior

$$f_{g/g}(x) \sim x\delta(1-x) \tag{2.154}$$

rather than Eq.(2.95) because we have chosen to eliminate negative arguments. In the quark case, this means that we must consider both a quark and an antiquark distribution. Since gluons are their own antiparticles, their distributions add.

tors of $SU(n_f)$. Due to the symmetry of our theory, these nonsinglet functions all satisfy the same evolution equation

$$\mu^2 \frac{D_q}{D\mu^2} f_{NS/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \left[P_{q\to q}\left(\frac{x}{y}\right) - P_{q\to\overline{q}}\left(\frac{x}{y}\right) \right] f_{NS/T}(y,\mu^2) \qquad (2.155)$$

to all orders in perturbation theory. Its appearance in this evolution equation as well as its role in the determination of A_q make

$$P_{NS}(x) \equiv P_{q\to q}(x) - P_{q\to\overline{q}}(x) \qquad (2.156)$$

deserving of the title 'nonsinglet splitting function'.

Equation (2.155) is also satisfied by the nonsinglet antiquark distributions and the difference between the singlet quark and antiquark distributions. Only the lone full singlet distribution,

$$f_{s/T}(x,\mu^2) \equiv \sum_q \left(f_{q/T}(x,\mu^2) + f_{\overline{q}/T}(x,\mu^2) \right) \qquad (2.157)$$

can mix with the gluon fields. Their evolution is coupled according to

$$\begin{aligned} \mu^2 \frac{D_q}{D\mu^2} f_{s/T}(x,\mu^2) &= \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \left[P_s\left(\frac{x}{y}\right) f_{s/T}(y,\mu^2) \right. \\ &\qquad \left. + 2n_f P_{g\to q}\left(\frac{x}{y}\right) f_{g/T}(y,\mu^2) \right] \ , \qquad (2.158) \\ \mu^2 \frac{D_g}{D\mu^2} f_{g/T}(x,\mu^2) &= \frac{\alpha_s(\mu^2)}{2\pi} \int_x^1 \frac{dy}{y} \left[P_{q\to g}\left(\frac{x}{y}\right) f_{s/T}(y,\mu^2) \right. \\ &\qquad \left. + P_{g\to g}\left(\frac{x}{y}\right) f_{g/T}(y,\mu^2) \right] \ , \qquad (2.159) \end{aligned}$$

where I have defined

$$P_s(x) \equiv 2n_f P_{q\to\overline{q}}(x) + P_{NS}(x) \ . \qquad (2.160)$$

The coupling of the gluon and singlet quark distributions makes data on these objects extremely difficult to analyze. Since one needs to know both distributions entirely at one scale in order to evolve either to another, measurements at different scales are not directly related to one another.

In the polarized sector, everything works in exactly the same way. The nonsinglet distributions decouple from the gluons and each other to evolve autonomously, while the fully singlet distribution mixes with the polarized gluon distribution. Here, the definition of the antiquark distribution implies that it is the *difference* of the quark and antiquark singlet distributions

that forms the fully singlet distribution, rather than the sum. The endpoint contributions, being properties of the fields themselves, are identical to those in the unpolarized case. Only the evolution kernels are different. In Appendix A.4, the full set of kernels at leading order in QCD is displayed. At next-to-leading order, one can find them in [43]. Note that Equation (2.151) is satisfied by these functions.

The expressions (2.144) are called Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution equations in honor of the physicists who first derived them [54]. The functions $\mathcal{P}_{a'\to a}(z)$, known either as *splitting functions* or DGLAP *evolution kernels*, allow us to sum certain effects to all orders in perturbation theory. For example, although the probability of finding an antiquark in a quark is zero at $\mathcal{O}(\alpha_s)$, the nonzero splitting functions $\mathcal{P}_{q\to g}$ and $\mathcal{P}_{g\to\overline{q}}$ will *generate* a nonzero probability via the DGLAP equation as the scale is varied. The distribution generated in this way will not be the same as that generated by using the correct nonzero splitting function $\mathcal{P}_{q\to\overline{q}}$, but it will certainly be closer than the direct leading order prediction of zero. This effect comes from our assumption (2.126) of the form the evolution will take. Once we assert this form, which is motivated on physical grounds, our equation already knows *part* of the result for the next order. The use of these kinds of evolution equations to sum certain contributions to all orders has been developed into a high art form and can be extremely useful in analyzing data.

2.4 Factorization to All Orders in DIS

Now that we have seen that the infrared divergences present in our one-loop amplitudes can be attributed to parton distributions in a parton, it is natural to ask if this property persists to all orders in perturbation theory. In this section, I will outline a general proof of factorization for inclusive DIS. This proof has been undertaken in a rigorous field-theoretic manner by several people [47, 48, 49, 55]. A very instructive recent application of the method under consideration here can be found in Ref. [56].

Our proof is structured as follows. First, we must identify the infrared-sensitive regions of our process. These regions can be characterized by certain reduced diagrams in which only the infrared behavior is emphasized. Since we work only at leading order in the Bjorken limit,[79]

[79] At higher orders, the arguments of Section 2.2 may no longer apply. Since these corrections are relevant




Figure 2.6: A representation of the all-orders scattering process. The nucleon 'blob' is intended to represent an all-orders sum, and the external photons are connected in all possible ways to this blob.

we will concern ourselves exclusively with regions that contribute to the leading behavior of the amplitude. These regions have special properties which allow us to show that their contribution does nothing more than generate the distribution functions introduced in the last section. Since all of the infrared-sensitive regions of our process can be segregated into these nonperturbative distributions, the remainder of the amplitude is due entirely to high energy physics and may be reliably calculated in perturbation theory.

To begin, we must find a way to isolate the regions of our process which are sensitive to soft scales. The easiest and most effective way to do this is to simply set all of the soft scales to zero. Since the perturbatively calculable hard scattering regions are insensitive to these scales, they will be unaffected by this procedure. On the other hand, the infrared regions of interest here will be forced to generate divergences. It must be emphasized that we are not assuming that the soft scales in our problem *are* zero. We are setting them to zero only as a means to *identify* the soft physics of our process.

To this end, we consider a general Feynman diagram in massless QCD. Since we work in principle to all orders in perturbation theory, we will consider an external nucleon state which we draw as a blob (shown in Figure 2.6). The propagator structure of an arbitrary diagram

for smaller values of Q^2, they will involve processes in which the partons do interact with each other during the scattering. Depending on the degree of this mixing of scales, we may or may not have a factorization theorem. Some of the necessary considerations for these power corrections are discussed in Chapter 4. For more detailed studies, especially with reference to factorization theorems, see [57].

may be combined into one denominator using a technique of Feynman's :[80]

$$\sum_i \alpha_i k_i^2(\ell_j^\mu) + i\varepsilon \ . \tag{2.161}$$

Here, the α_i are Feynman parameters and the k_i are propagator momenta (which are related linearly to the loop momenta ℓ_i^μ). Infrared divergences come from regions of the joint parameter-momentum space where (2.161) vanishes. Not all of these regions can give an infrared divergence. Consider the integral

$$\int_0^3 \frac{dx}{x-1} = \log 2 \ . \tag{2.162}$$

The integrand surely diverges at $x = 1$, but the integral itself is finite.[81] This is because the contour of integration can be deformed away from the singularity free of charge. These kinds of integrals can lead to branch cuts in our amplitudes and contribute to the imaginary part, but they can never give infrared divergences.

There are only two ways an integral can lead to an infrared divergence. The first is if the integrand is divergent at an endpoint of the integration domain. In this case, we cannot deform the contour away from the singularity. An example of an endpoint singularity is given by

$$\int_0^1 \frac{dx}{x} \ . \tag{2.165}$$

This integral is unavoidable infinite. Note that this is the kind of integral that generates the light-cone singularities in the PV-regulated axial gauge.

The second way to obtain an unavoidably infinite integral is by pinching the contour so that it cannot be deformed away from the singularity, for example

$$\lim_{\varepsilon\to 0}\int_{-1}^1 \frac{dx}{(x-i\varepsilon)(x+i\varepsilon)} = \lim_{\varepsilon\to 0}\frac{2}{\varepsilon}\,\mathrm{Tan}^{-1}\left(\frac{1}{\varepsilon}\right) \ . \tag{2.166}$$

[80] see Appendix A.3.

[81] Mathematically, this integral is ill-defined. However, a prescription such as

$$\lim_{\delta\to 0}\left[\int_0^{1-\delta}\frac{dx}{x-1} + \int_{1+\delta}^3 \frac{dx}{x-1}\right] = \log 2 \tag{2.163}$$

can be used to define the integral unambiguously. This is what the $+i\varepsilon$'s in our propagators do. In this case, we would obtain an imaginary part from the prescription :

$$\int_0^3 \frac{dx}{x-1+i\varepsilon} = \log 2 - i\pi \ . \tag{2.164}$$

This integral is also unavoidable infinite. However, note that the expression

$$\lim_{\varepsilon\to 0}\int_{-1}^{1}\frac{dx}{(x+i\varepsilon)^2}=-2 \tag{2.167}$$

is *not* divergent. This singularity is not pinched and therefore the integration contour may be deformed away from it.[82] These two cases are the only ways a contour can be forced through a singularity, so these are the only kinds on singularities we need consider in our search for infrared divergences.

Since (2.161) is linear in $\{\alpha_j\}$, singularities in this space *cannot* be pinched. We can deform the parameter contour to avoid all singularities unless either

$$\alpha_i=0 \qquad \text{or} \qquad k_i^2=0 \tag{2.169}$$

for *each* i. In the first case, the singularity occurs at an endpoint so cannot be deformed away.[83] In the second, (2.161) is independent of α_i so contour deformation produces no effect. This region of parameter-momentum space will produce an infrared divergence if it is pinched in momentum space. Since (2.161) is quadratic in k_i, there are two poles for each k_i. These poles are pinched only if they occur at the same point on the real line. The two zeros of a quadratic form $D(x)$ are equal iff $dD(x)/dx=0$ simultaneously with $D(x)$. The structure of Feynman diagrams is such that one can always choose loop momenta to appear with the same

[82]It may seem a little strange that this integral is not infinite. If we imagine adding up its contributions, we would certainly get infinity. However, we must not think of our manipulations in this way. What we are discussing here is the *analytic structure* of our integrals. The correct way to interpret this result is by considering instead

$$\lim_{a\to 0}\int_{-1}^{1}\frac{dx}{(x+a)^2}=\lim_{a\to 0}\frac{2}{a^2-1}=-2\ . \tag{2.168}$$

In some sense, it is only a numerical accident that 0 lies between the two endpoints of integration. The true singularities in this family of integrals lie at $a=\pm 1$, the endpoints. This may seem like a swindle, but in reality it is the only way to interpret objects like (2.167). These kinds of expressions are ill-defined only in specific regions of our parameter space. The only sensible thing to do is evaluate them away from these regions and analytically continue the result later. This is exactly what was done in Section 2.2, where we took $x_B\gg x$ to simplify the calculation and analytically continued our amplitudes to $x_B\leq x$ at the end.

[83]The other endpoint, $\alpha_i=1$, requires all other parameters to be zero (since Feynman parameters are constrained by $\sum_i\alpha_i=1$). If the denominator (2.161) vanishes in this case, $k_i^2=0$. Hence this also reduces to (2.169).

sign in *all* of the propagators.[84] Since propagator momenta always appear quadratically in the denominator, we can choose this sign to be positive.[85] With this choice of routing, the condition for a pinched singularity in ℓ_j^μ becomes

$$\sum_i \alpha_i k_i^\mu = 0 \ . \tag{2.170}$$

The sum goes over all propagators through which ℓ_j^μ flows. In order to meet the criteria for an infrared divergence, Equation (2.169) must be satisfied by each propagator *and* Equation (2.170) must be satisfied by each loop momentum. The conditions are collectively referred to as the Landau equations [50].

Equation (2.170) has a nice physical interpretation due to Coleman and Norton [58]. Since $\alpha_i k_i^\mu$ is proportional to the 4-velocity of propagator i, we can think of it as the distance traversed by the particle it represents in a time $\alpha_i k_i^0$.[86] (2.170) states that as we go around the loop ℓ_j, we get back to the point we started from. Since propagators with $\alpha_i \neq 0$ are onshell ($k_i^2 = 0$), the Landau equations state that pinched surfaces come only from *classically allowed* propagation.[87] Each solution of the Landau equations has a corresponding 'pinched diagram' in which all lines representing propagators with $\alpha_i = 0$ have been shrunk to a point[88] and all other lines represent onshell propagation.

For DIS on a hadron target, the most general pinched diagram for $T^{\mu\nu}$ is shown in Figure 2.7. Here, we see that any number of (collinear) parton lines may connect the hard scattering to the hadron blob. However, the hard scattering itself must be considered a point interaction. Diagrams in which this is not the case cannot satisfy (2.169) and (2.170) simultaneously since two or more onshell propagators leaving the first photon interaction vertex must be moving in different directions and therefore can never (classically) recombine at the other photon vertex.

[84] This can be seen by inspection.

[85] All we are saying here is that one can always choose a momentum routing such that $k_i = \sum_j \ell_j + q$, where q is some external momentum and the sum extends over all loops whose momenta flow through k_i.

[86] Since the units of α_i are arbitrary, we can consider it a frame-independent ratio of time to energy k_i^0.

[87] We can obtain classically unallowed propagation in quantum mechanics since momentum eigenstates do not have well-defined spacetime locations. To see this in another way, recall that momentum conservation at each vertex is imposed *after* integration over the spacetime location of the vertex.

[88] This is because $\alpha_i = 0$ implies no propagation. In this region of parameter space, the integral is independent of k_i^μ so represents an position eigenstate.




Figure 2.7: The most general reduced diagram for DIS. Straight lines represent collinear (or soft) partons of *any* species.

Shrinking all offshell propagators to a point thus gives us a pointlike hard interaction. This implies that summing over all final states in DIS cancels the infrared divergences associated with final state interactions.[89] Physically, we can understand this as a consequence of uniterity. By definition, final state interactions take place on the scale of nonperturbative QCD. For large Q^2, the arguments of Section 2.2 imply that these interactions can only occur long after the hard scattering has taken place. As such, they cannot affect the probability that the scattering will take place. By the time final state interactions come into play, that decision has already been made. The only thing these interactions can affect is the final form of the outgoing physical state. Therefore, their effects *must* cancel out in the inclusive cross-section.[90]

Although all diagrams of the type in Figure 2.7 can contribute to infrared divergences, they will not all contribute at leading order in the Bjorken limit. The Bjorken suppression of various graphs may be deduced from Q^2 power counting. A detailed analysis [47] reveals that only those diagrams in which the hard scattering is connected to the hadron blob by two 'physical'

[89] Without the sum over final states, we could not obtain the cross-section for DIS from $T^{\mu\nu}$.

[90] As Q^2 is lowered, the time the hard interaction takes grows until there is no longer a clear separation between the (not-so-)hard interaction and final (and initial) state interactions. At this point, the above analysis breaks down. These nonperturbative interactions certainly can and will affect the total inclusive cross-section. In our classification of reduced diagrams, the assumption of large Q^2 is reflected in the fact that we take point-like couplings for the photon interactions. Less virtual photons have spatially extended couplings which allow classical propagation between them, reinstating final state interactions.

parton propagators and any number of collinear scalar[91] gluons give leading contributions in the Bjorken limit. The appearance of scalar gluons can be understood by considering the simplest case of one gluon interacting directly with the intermediate quark propagation.[92] The gluon insertion breaks the quark propagator into two, giving a suppression of Q^2 unless this can be canceled by the numerator factors. Since large terms in the numerator appear only with $\not{n}$, leading behavior occurs only when the interaction involves $\not{p}$. In light-cone gauge, this is not possible since $\not{p}$ is related to $\mathcal{A}^+ = 0$. In covariant gauge, we can obtain $\not{p}$ by contracting the other side of the propagator with p^μ :

$$p^\mu \left(-g_{\mu\nu} + (1-\xi)\frac{p_\mu p_\nu}{p^2} \right) \gamma^\nu = -\xi \not{p} \ . \tag{2.171}$$

This procedure gives zero for strictly collinear gluons in light-cone gauge,[93]

$$p^\mu \left(-g_{\mu\nu} + p_\mu n_\nu + p_\nu n_\mu \right) = 0 \ , \tag{2.172}$$

so we can say in either case that leading behavior comes only from scalar gluons. Generalizing this analysis to higher loops and other couplings leads to the above result.

A simple way to understand why these diagrams are favored over others comes from light-cone dimensional analysis. Since the dimension of an amplitude is fixed, all soft mass dimensions must be compensated by the hard scale Q^2. Assuming covariant normalization for the external states ($\langle p|p\rangle = 2p^0(2\pi)^3\delta^3(0)$), every external wavefunction contributes mass dimension -1. The collinear quarks and gluons in a soft hadron vertex have effective mass dimensions depending on their polarizations. A Dirac field ψ can be written as a sum of good (ψ_+) and bad (ψ_-) components, where $\psi_\pm = P_\pm \psi$ and $P_\pm = \frac{1}{2}\gamma^\mp\gamma^\pm$. The good (bad) component has effective light-cone mass dimension 1 (2).[94] The vector potential A^μ has light-cone components A^+, $A^\perp$, and A^-, which have effective mass dimensions 0, 1, and 2, respectively.[95] For the

[91] Scalar gluons can be thought of as gluons whose polarization vectors are dotted with their momenta. I.e. the 'scalar gluon field' is written $\partial_\mu \mathcal{A}^\mu$.

[92] Like Figure 2.2c in Section 2.2.

[93] In these manipulations, we take $p^2 = 0$ wherever it will not cause a divergence.

[94] Since we consider collinear quarks moving in the 3-direction, the free quark wavefunction is annihilated by $\not{p}$. This means that the 'bad' component cannot freely propagate. We will see in Chapter 4 that this part of the quark field really represents a quark-gluon composite rather than a simple quark.

[95] As in the quark case, this is related to equation of motion constraints.




Figure 2.8: A graphical statement of factorization in the axial gauge.

reduced diagram shown in Figure 2.7, the only soft mass dimension comes from the nucleon-quark-gluon blob. Using the above rule, we find it is $\Sigma m_i - 2$. Here, m_i is the light-cone mass dimension of the ith collinear parton line. Our scattering process requires at least two physical ($m \neq 0$) partons, so $\Sigma m_i \geq 2$. Since physical parton lines add soft dimension to our matrix element, they lead to suppression in our amplitudes. Only the scalar gluon field $\mathcal{A}^+$ can be added without penalty. Hence, we see that the most general leading pinched collinear surface is represented by Fig. 2.7, with two parton lines with $m = 1$ and any number of scalar gluons connecting the nucleon blob to the hard scattering. Noncollinear lines cannot classically reach the hard interaction, so the only other possibility for a pinched singularity involves soft lines. Since zero is in some sense collinear to everything, these lines are in principle contained in Figure 2.7. However, we must consider them explicitly as they can lead to divergences that will not factorize into our parton distributions.

The power counting involving soft quark and gluon lines is more subtle, and some discussion may be found in Ref. [56]. Essentially, one applies a general theorem due to Kinoshita, Lee and Nauenberg (KLN) [59] which states that infrared divergences cancel in the sum over initial *and* final states. According to the above arguments, soft initial state interactions are suppressed by Q^2 while collinear ones only serve to enforce the gauge-invariance of our process. Hence our sum over final state interactions only is enough to cancel the *leading* soft divergences. Because of this cancelation, we can consider any reduced diagram with soft lines connecting the hard scattering blob to the nucleon jet subleading. The situation here is discussed in detail in Sterman [60].

In the light-cone gauge, where collinear scalar gluons do not exist, the jump from this anal-

ysis to a factorization theorem is in some sense obvious. Here, $\Psi = \psi$ and only those diagrams with two physical partons connecting the hard and soft physics contribute at leading order. In this case, factorization is represented graphically by Figure 2.8. However, the complications associated with renormalization in light-cone gauge[96] make it desirable to consider covariant gauges. In these gauges, factorization follows from the Ward-Takehashi identities introduced in Appendix D.2. Diagrammatically, these identities tell us that the sum of all insertions of a scalar gluon is zero. This is true for all physical matrix elements of gauge-invariant operators. In particular, we can apply it both to the hard scattering kernel and to the eikonal line. A graphical illustration of this application is shown in Figures 2.9a and b. Since the right-hand-sides of these graphical equations are identical, we are free to make the identification in 2.9c. Physically, the scalar gluons are *coherently* absorbed by the hard jet, which is represented by an eikonal line. They do not possess the immense amounts of energy necessary to resolve the individual constituents of the jet, so only the *total* color charge and momentum are relevant. This kind of coherent absorption cannot affect the hard scattering in a nontrivial way; its only effect is to induce gauge invariance in the final result. As we saw in the last section, this is exactly the effect of the gauge link in our parton distributions. Seen from this angle, it is not surprising that the scalar gluons generate our distributions. The proof to all orders is by induction; applying the Ward identities iteratively, one easily arrives at the factorization theorem expressed in Equation (2.117).

2.5 The Operator Product Expansion

In the last three sections, we have shown that the hard and soft physics of the amplitude[97]

$$T^{\mu\nu} = i \int d^4z \, e^{-iq\cdot z} \, \langle PS \,|\mathrm{T}\, J^\mu(0)\, J^\nu(z)|\, PS\rangle \tag{2.173}$$

can be consistently separated from each other if the virtuality $Q^2 = -q^2 \gg M\Lambda_{\mathrm{QCD}}$. Furthermore, we have argued that this separation is *independent* of the target. Since the target dependence of this amplitude is contained entirely within the external states, it is natural to

[96] see Section 1.6

[97] The difference between this expression and that in Eq.(2.22) is purely aesthetic, as can be shown immediately by shifting the argument and changing $z \to -z$.




Figure 2.9: A graphical illustration of the use of Ward identities to show that scalar gluons generate the gauge link in our parton distribution functions. Dashed lines represent the propagation of a scalar gluon.

ask whether or not we can strip these states away and express factorization as an operator relation. This idea was first introduced by Wilson in 1969 [29] and has been used extensively in deep-inelastic scattering and other perturbative QCD processes. The fundamental idea is that for large Q^2 the integral in (2.173) has support only in the region of small z. This means that our amplitude receives contributions only from regions in which the operator product $J^\mu J^\nu$ is nearly local, which suggests a kind of Taylor expansion of the operator $J^\mu(z)$ about $z = 0$. In this way, we express the operator

$$i \int d^4z \, e^{-iq\cdot z} \, \mathrm{T} J^\mu(0) \, J^\nu(z) \tag{2.174}$$

as an infinite series, known as an Operator Product Expansion (OPE), of local operators. For the product of two currents separated near the light-cone, the expansion is threefold. Primarily, it is a twist expansion, in which twist-2 contributions are leading whereas the higher twist terms are suppressed by powers of M/Q. Each term in the twist expansion contains an infinite number of local operators of the relevant twist. Finally, the coefficients of these operators (Wilson coefficients) are themselves expansions in the strong coupling constant. In this section, I will show how we can convert the results of Section 2.3 into next-to-leading order results for the Wilson coefficients of the OPE relevant to DIS. The Wilson coefficients for the unpolarized DIS process were first calculated at order α_s in the $\overline{\mathrm{MS}}$ scheme in [61]. For the polarized case, one can find them in [62].

To get a better handle on the meaning of the OPE, it is useful to construct the leading order Wilson coefficients directly from the operator, Eq.(2.174). Since we take z near zero, we will be concerned only with the most singular terms in this limit. By inspection, one can see that singular behavior occurs only when we contract two quark fields from the different currents together to form a propagator.[98] For $z^0 < 0$, we have

$$\begin{aligned}
&i \int d^4z \, \Theta(-z^0) e^{-iq\cdot z} J^\mu(0) \, J^\nu(z) \\
&\quad = i \int d^4z \, \Theta(-z^0) e^{-iq\cdot z} \int \frac{d^4k}{(2\pi)^4} e^{ik\cdot z} \sum_q e_q^2 \overline{\psi}_q(0) \gamma^\mu \, \frac{i}{\not{k}} \, \gamma^\nu \psi_q(z) \\
&\quad = i \int d^4z \, \Theta(-z^0) e^{-iq\cdot z} \int \frac{d^4k}{(2\pi)^4} \sum_q e_q^2 \overline{\psi}_q(0) \gamma^\mu \left(\frac{i}{-i \overleftarrow{\not{\partial}}_z} e^{ik\cdot z} \right) \gamma^\nu \psi_q(z)
\end{aligned}$$

[98] This implies that we need not consider terms in the operator product that couple different flavors.

$$= -\sum_q e_q^2 \overline{\psi}_q(0)\gamma^\mu \, \frac{1}{i\not{\partial}+\not{q}} \, \gamma^\nu \psi_q(0) \ . \tag{2.175}$$

In the last line, I have integrated by parts[99] (ignoring a surface term) and performed the integrals over z and k. The result for $z^0 > 0$ is identical to this one, but with $q \to -q$ and $\mu \leftrightarrow \nu$.[100]

These results are very similar to those obtained in Section 2.2, as they must be. We can get those results simply by tacking on external states and allowing the leftover quark fields to contract with them. Alternatively, using

$$\frac{1}{\not{p}+\not{q}} = \frac{\not{p}+\not{q}}{q^2} \sum_{n=0}^{\infty} \left(-\frac{p^2 + 2p\cdot q}{q^2} \right)^n \ , \tag{2.176}$$

we can turn this expression into the expansion

$$\begin{aligned} i\int d^4z\, e^{-iq\cdot z}\, \mathrm{T}J^\mu(0)\, J^\nu(z) \quad &= \quad \sum_q e_q^2\, \overline{\psi}_q \sum_{n=0}^{\infty} \left[\gamma^\mu \frac{i\not{\partial}+\not{q}}{Q^2} \left(\frac{2q\cdot i\partial - \partial^2}{Q^2} \right)^n \gamma^\nu \right. \\ &\left. + \gamma^\nu \frac{i\not{\partial}-\not{q}}{Q^2} \left(\frac{-2q\cdot i\partial - \partial^2}{Q^2} \right)^n \gamma^\mu \right] \psi_q \end{aligned} \tag{2.177}$$

involving only local operators. The operator ∂^2 cannot hope to compete with Q^2 in the Bjorken limit unless we intend to consider external states with incredible amounts of ambient energy, so we can safely neglect this term. On the other hand, if ∂_μ chooses its orientation wisely, $q\cdot i\partial$ can survive this limit.

Simplifying the Dirac structure and cleaning house a little, we arrive at

$$\begin{aligned} & i\int d^4z\, e^{-iq\cdot z} \mathrm{T}J^\mu(0)\, J^\nu(z) \\ &= \quad \frac{2}{Q^2} \sum_q e_q^2 \left(g^{\mu\alpha} + \frac{q^\mu q^\alpha}{Q^2} \right) \left(g^{\nu\beta} + \frac{q^\nu q^\beta}{Q^2} \right) \\ &\qquad \times {\sum_{n=0}^{\infty}}' \overline{\psi}_q \left(i\partial_\alpha \gamma_\beta + i\partial_\beta \gamma_\alpha \right) \left(\frac{2q\cdot i\partial}{Q^2} \right)^n \psi_q \\ &\quad - \frac{2}{Q^2} \sum_q e_q^2 \left(g^{\mu\nu} + \frac{q^\mu q^\nu}{Q^2} \right) \end{aligned} \tag{2.178}$$

[99] The action of the derivative on the Θ-function generates a contact term which can be subtracted by considering a modified time-ordering operation, T^*. This technical point is discussed in Ref. [63]. Here, we will assume that the appropriate modifications have been made and ignore this term.

[100] In order to show this, we have to use the fact that we consider only *diagonal* matrix elements. As full-fledged operators, these objects differ by a total derivative.

$$\times \sum_{n=1}^{\infty}{}' \overline{\psi}_q \not{q} \left(\frac{2q\cdot i\partial}{Q^2}\right)^n \psi_q \tag{2.179}$$

$$- i\epsilon^{\mu\nu\alpha\beta}\frac{2}{Q^2}\sum_q e_q^2$$

$$\times \sum_{n=0}^{\infty}{}' \overline{\psi}_q \left(\frac{2q\cdot i\partial}{Q^2} i\partial + q\right)_\alpha \gamma_\beta\gamma_5 \left(\frac{2q\cdot i\partial}{Q^2}\right)^n \psi_q \tag{2.180}$$

$$+ \frac{2}{Q^2}\sum_q e_q^2 \left(\frac{q^\mu q^\alpha}{Q^2} g^{\nu\beta} - \frac{q^\nu q^\beta}{Q^2} g^{\mu\alpha}\right)$$

$$\times \sum_{n=0}^{\infty}{}' \overline{\psi}_q \left(i\partial_\alpha\gamma_\beta - i\partial_\beta\gamma_\alpha\right)\left(\frac{2q\cdot i\partial}{Q^2}\right)^n \psi_q \tag{2.181}$$

$$- \frac{2}{Q^2} g^{\mu\nu}\sum_q e_q^2 \sum_{n=0}^{\infty}{}' \overline{\psi}_q i\not{\partial}\left(\frac{2q\cdot i\partial}{Q^2}\right)^n \psi_q \ . \tag{2.182}$$

Here, a primed summation indicates that only even or odd numbers are summed over, depending on whether the sum starts with 0 or 1.

Contracting with q_μ, we immediately see that the first two terms automatically conserve current. However, the last three operators do not satisfy the Ward identity. To see why, let us imagine taking diagonal hadronic matrix elements of our expression. Since the operators on the right-hand-side do not have any intrinsic dependence on q^μ, Lorentz invariance requires all of their vector indices to be carried by the hadronic momentum and all of their axial indices to be carried by the spin. This implies that (2.182) vanishes by the equation of motion,[101] and (2.181) does so identically. In order to have some other behavior, we need to introduce another momentum into the problem - an action that invariably leads to other contributions which will combine with these to restore gauge invariance. The third term, (2.180), retains the nonvanishing gauge dependent piece

$$\epsilon^{\mu\nu\alpha\beta} P_\alpha S_\beta \tag{2.183}$$

after we have taken our matrix element. In the Bjorken limit, this tensor must vanish since S is parallel to P. Although it's more difficult to see, any deviation from this limit will imply other contributions to the amplitude. We will see in Chapter 4 that these contributions do indeed cancel this gauge dependence. For now, let us consider only leading contributions and take this object as zero.

[101] which, to the order at which we work, is simply $i\not{\partial}\psi = 0$.

Now that we have taken care of electromagnetic gauge invariance, it behooves us to consider chromodynamic gauge invariance. As it stands, none of the operators on the right-hand-side of our expansion are gauge-invariant. The reason for this is very simple - we are working only to leading order in QCD. The culprit is our replacement of $\psi(0)\overline{\psi}(z)$ with the Dirac propagator. To maintain explicit gauge invariance without calculating to all orders, we can introduce a gauge link between the two remaining fields simultaneously with the replacement. The effect of this procedure is a simple replacement of all partial derivatives with covariant derivatives.

Stripping away the factors of q^μ, we see that our expansion involves local operators of the form

$$\overline{\psi}\gamma^{\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_n}\psi \quad . \tag{2.184}$$

As explained in Appendix B.3, n-index tensors such as these can be separated into several disjoint pieces. Each of these pieces transforms separately from the others under the action of the Lorentz group. As a symmetry of our theory, Lorentz covariance restricts the possible divergences that can appear. For this reason, operators which begin in the same irreducible representation of a symmetry group will remain together in that representation after renormalization. However, Lorentz invariance cannot impose relations between two *different* representations. The operators that appear in our expansion actually represent sums of operators in different representations, all of which renormalize independently of one another. Rather than this mess, we would prefer to separate our operators into groups which stay together after renormalization. The procedure for this separation is straightforward and outlined in the appendix. Here, we mention only that the tensors required for our purpose are those with the highest spin. All other structures are subleading.[102] In view of this, we define

$${}_q\mathcal{O}_n^{\mu_1\mu_2\cdots\mu_n} \equiv \overline{\psi}_q\gamma^{(\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_n)}\psi_q \tag{2.185}$$

$${}_q\tilde{\mathcal{O}}_n^{\mu_1\mu_2\cdots\mu_n} \equiv \overline{\psi}_q\gamma^{(\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_n)}\gamma_5\psi_q \quad . \tag{2.186}$$

In terms of these operators, we can write our expansion as

$$i\int d^4z\, e^{-iq\cdot z}\,\mathrm{T}J^\mu(0)J^\nu(z) = \quad \frac{2}{Q^2}\sum_q e_q^2 \sum_{n=0}^{\infty}{}' \frac{(2q_{\mu_1})(2q_{\mu_2})\cdots(2q_{\mu_n})}{(Q^2)^n}$$

[102] This can be seen simply be taking matrix elements. The hadron's momentum and helicity cannot support any antisymmetric tensors and the traces all lead to powers of M^2.

$$\times \left\{ \left[2 \left(\delta^{\mu}_{\alpha} + \frac{q^{\mu} q_{\alpha}}{Q^2} \right) \left(\delta^{\nu}_{\beta} + \frac{q^{\nu} q_{\beta}}{Q^2} \right) \right. \right. \tag{2.187}$$
$$\left. -2 \, \frac{q_{\alpha} q_{\beta}}{Q^2} \left(g^{\mu\nu} + \frac{q^{\mu} q^{\nu}}{Q^2} \right) \right]_{q} \mathcal{O}^{\alpha\beta\mu_1 \cdots \mu_n}_{n+2}$$
$$\left. - i \epsilon^{\mu\nu\alpha\beta} q_{\alpha} g_{\beta\lambda \, q} \tilde{\mathcal{O}}^{\lambda\mu_1 \cdots \mu_n}_{n+1} \right\} \quad .$$

This is the desired expansion of the product of two electromagnetic currents in terms of an infinite series of local operators. Each operator appears with a power of Q to compensate its mass-dimension and a power of q_{α}/Q for most of its tensor indices. Removing trivial kinematical factors,[103] a general operator of spin s and mass-dimension m appears in our expansion appears as[104]

$$\frac{(2q_{\mu_1})(2q_{\mu_2}) \cdots (2q_{\mu_s})}{(Q^2)^s} \frac{1}{Q^{m-s-2}} {}_{q}\mathcal{O}^{\mu_1 \mu_2 \cdots \mu_s} \quad . \tag{2.188}$$

Taking matrix elements with momentum P such that $P \cdot q = \nu$, we see that the contribution of this operator is suppressed by Q^{m-s-2}. Since it governs the suppression of these operators, the *twist*

$$t \equiv m - s \tag{2.189}$$

is of great importance. A look at Equations (2.185) and (2.186) tells us that the twist of the operators in our leading expansion is $(n-1) + 3 - n = 2$. Hence these operators are not suppressed in the Bjorken limit.

As mentioned above, operators of different spin cannot mix under renormalization due to the Lorentz invariance of our theory. Likewise, the absence of a scale[105] in massless QCD prevents operators with different mass-dimension from mixing with one-another. This implies that only operators of the same twist can mix. An exhaustive classification of QCD operators in terms of twist [64] reveals that the lowest possible twist is twist-2. Furthermore, the only

[103] This amounts to ignoring each index that is 'bogged down' in a tensor structure involving $\mu\nu$, and along with it a power of Q. In addition, factors of q_{α} adorning these tensor structures should also be ignored along with a corresponding factor of Q. By taking matrix elements and comparing with (2.28), one can see that this is the proper procedure to determine the suppression of the structure function itself.

[104] This is purely a dimensional argument. Note that the mass-dimension of our operator $\int \mathrm{T}\, JJ$ is 2.

[105] The scale generated through the coupling by dimensional transmutation is of no help here. Its effect is limited to producing scale variations; μ^2 can appear only in a ratio.

other operators of twist-2 in the nonsinglet sector,[106]

$$ {}_q\hat{\mathcal{O}}_n^{\alpha\mu_1\cdots\mu_n} = \overline{\psi}_q \sigma^{\alpha(\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_n)}\psi_q \quad , \tag{2.190} $$

must completely decouple from our operators in massless QCD because of their chiral-odd structure.[107] Since the operators (2.185) and (2.186) have different transformation properties under parity, they cannot mix with each other. This leads us to the result that all of our operators must have diagonal matrix elements which evolve multiplicatively.[108] Hence if we can extract the value of one of these matrix elements at any scale, we automatically know it at all (perturbative) scales. This represents a tremendous simplification from the results of Section 2.3, where we found that knowledge of the entire parton distribution at one scale is necessary to evolve any part of it to another. Unfortunately, we will see that this simplification is not devoid of distasteful attributes.

To see the relation between our expansion (2.187) and the structure functions of (2.28), we must take forward matrix elements. Using Lorentz symmetry, we can express the matrix elements of ${}_q\mathcal{O}$ and ${}_q\tilde{\mathcal{O}}$ in terms of pure numbers :

$$ \left\langle P \left| {}_q\mathcal{O}_n^{\mu_1\cdots\mu_n} \right| P \right\rangle \equiv 2a_n^q P^{(\mu_1}\cdots P^{\mu_n)} \tag{2.191} $$

$$ \left\langle P \left| {}_q\tilde{\mathcal{O}}_n^{\mu_1\cdots\mu_n} \right| P \right\rangle \equiv 4\tilde{a}_n^q M S^{(\mu_1}\cdots P^{\mu_n)} \quad . \tag{2.192} $$

In this way, we arrive at

$$ T_1(\nu, Q^2) = 2\sum_{n=2}^{\infty}{}' \left(\sum_q e_q^2 a_n^q \right) \left(\frac{1}{x_B} \right)^n \tag{2.193} $$

$$ T_2(\nu, Q^2) = 4\sum_{n=1}^{\infty}{}' \left(\sum_q e_q^2 a_{n+1}^q \right) \left(\frac{1}{x_B} \right)^n \tag{2.194} $$

$$ S_1(\nu, Q^2) = 2\sum_{n=1}^{\infty}{}' \left(\sum_q e_q^2 \tilde{a}_n^q \right) \left(\frac{1}{x_B} \right)^n \quad . \tag{2.195} $$

These equations express a direct relation between the structure functions of the Compton amplitude and the matrix elements of local QCD operators. In particular, barring belligerent

[106] Here, we consider only operators with nonzero forward matrix elements. We will meet other twist-2 operators in the next chapter.

[107] These operators relate left- and right-handed quarks. As such, they cannot contribute perturbatively in massless QCD.

[108] Once gluons are introduced, we will that the flavor singlet quark operators mix with the gluonic operators.

behavior on the part of a_n and $\tilde{a}_n$, they tell us that our structure functions are analytic functions of x_B near $x_B = \infty$. In addition, the hermiticity of ${}_q\mathcal{O}$ and ${}_q\tilde{\mathcal{O}}$ implies that our structure functions are real for x_B on the real axis. This means that the true measured structure functions, F_1, F_2, and G_1, can only be nonzero in nonanalytic regions of the x_B-plane. In these regions, branch cuts can allow discontinuities across the real axis which generate an imaginary part. This fact implies that our operator product expansion *cannot* converge in the physical region. However, it can be used as a powerful tool to study the analytic structure of the Compton amplitude. Moreover, properties such as the Callan-Gross relation, $T_2(\nu, Q^2) = 2x_B T_1(\nu, Q^2)$, can be studied more easily from this vantage.

To go further, we must find some way to relate the scalar matrix elements to one-another. This is done following a hint from Section 2.3. There, we found that the integral of x times a structure function is related to the matrix element of a local operator appearing in $\Theta^{\mu\nu}$. Generalizing this procedure, we write

$$\begin{aligned}
&\int dx x^{n-1} f_{q/T}(x) \\
&\qquad = \frac{1}{2}\int dx \frac{d\lambda}{2\pi} e^{i\lambda x} x^{n-1} \left\langle PS \left| \overline{\psi}\,(0)\, \mathcal{G}_n\,(0,\lambda n) \not{n}\, \psi\,(\lambda n) \right| PS \right\rangle \\
&\qquad = \frac{1}{2}\int dx \frac{d\lambda}{2\pi} \left(\frac{1}{i}\frac{d}{d\lambda}\right)^{n-1} e^{i\lambda x} \left\langle PS \left| \overline{\psi}(0) \mathcal{G}_n\,(0,\lambda n) \not{n}\, \psi(\lambda n) \right| PS \right\rangle \\
&\qquad = \frac{1}{2}\int dx \frac{d\lambda}{2\pi} e^{i\lambda x} \left\langle PS \left| \overline{\psi}(0) \mathcal{G}_n\,(0,\lambda n) \not{n}\, \left(n\cdot i\mathcal{D}(\lambda n)\right)^{n-1} \psi(\lambda n) \right| PS \right\rangle \\
&\qquad = \frac{1}{2} \left\langle PS \left| \overline{\psi}(0) \not{n}\, \left(n\cdot i\mathcal{D}\right)^{n-1} \psi(0) \right| PS \right\rangle \quad , \qquad (2.196)
\end{aligned}$$

where we have ignored a surface term. Performing the same manipulations for the polarized distribution and comparing with (2.191) and (2.192), we see that

$$\int dx\, x^{n-1}\, f_{q/T}(x) = a_n^q \qquad (2.197)$$

$$\int dx\, x^{n-1}\, \tilde{f}_{q/T}(x) = \tilde{a}_n^q \quad . \qquad (2.198)$$

Integrals of this form are called *moments*. Mathematically, the correspondence between a_n^q and $f_{q/T}(x)$ is contained in *Mellin* transformation. Substituting these expressions into (2.193), (2.194), and (2.195), switching the order of summation and integration,[109] and re-summing,

[109] I refuse to justify this mathematically.

we arrive at

$$T_1(\nu,Q^2) = \int \frac{dx}{x}\ \frac{x^2}{x_B}\ \frac{1}{x_B - x}\ \sum_q e_q^2\ f_{q/T}(x) + (x_B \to -x_B) \tag{2.199}$$

$$T_2(\nu,Q^2) = \int \frac{dx}{x}\ 2x^2\ \frac{1}{x_B - x}\ \sum_q e_q^2\ f_{q/T}(x) - (x_B \to -x_B) \tag{2.200}$$

$$S_1(\nu,Q^2) = \int \frac{dx}{x}\ \ x\ \frac{1}{x_B - x}\ \sum_q e_q^2\ \tilde{f}_{q/T}(x) - (x_B \to -x_B)\ . \tag{2.201}$$

These expressions are familiar from Section 2.2, as they should be. At this point, we are free to take imaginary parts and obtain the physical structure functions[110]

$$F_1(\nu,Q^2) \ = \ \frac{1}{2}\sum_q e_q^2\left[f_{q/T}(x_B) + f_{\overline{q}/T}(x_B)\right] \tag{2.202}$$

$$F_2(\nu,Q^2) \ = \ x_B\sum_q e_q^2\left[f_{q/T}(x_B) + f_{\overline{q}/T}(x_B)\right] \tag{2.203}$$

$$G_1(\nu,Q^2) \ = \ \frac{1}{2}\sum_q e_q^2[\tilde{f}_{q/T}(x_B) - \tilde{f}_{\overline{q}/T}(x_B)]\ . \tag{2.204}$$

Note that we have introduced the antiquarks. Their inclusion does not significantly alter the derivation above.

Since we have only considered the leading order in α_s, Equation (2.187) is guaranteed to have QCD corrections. A calculation in the spirit of the above derivation would invariably lead to arthritis. Alternatively, we can turn the above analysis on its head and use the results of Section 2.3 to obtain these corrections. Our amplitudes have the form

$$T \sim \int \frac{dx}{x} f(x) C(x, x_B)\ . \tag{2.205}$$

In the unphysical region $x_B \gg x$, we write

$$T \sim \sum_{n=0}^{\infty} \int \frac{dx}{x} f(x) c_n\ \left(\frac{x}{x_B}\right)^n\ . \tag{2.206}$$

Substituting our expressions for the moments, we obtain an expansion analogous to (2.193). Writing

$$\left(\frac{1}{x_B}\right)^n = \frac{(2q_{\mu_1})(2q_{\mu_2})\cdots(2q_{\mu_n})}{(Q^2)^n} P^{\mu_1} P^{\mu_2} \cdots P^{\mu_n} \tag{2.207}$$

[110] The mathematical indiscretion above generated these imaginary parts. It can be justified in the unphysical region where the expansion converges. Once this has been done, the re-summed unphysical amplitudes can be analytically continued to the physical region.

and interpreting the resulting expressions as matrix elements of the local operators (2.185) and (2.186), we can strip away the matrix elements and arrive at the operator product expansion[111]

$$
\begin{aligned}
& i\int d^4z\, e^{iq\cdot z}\ \mathrm{T}\, J^\mu(z)J^\nu(0) \\
&\quad = \left(\frac{q^\mu q^\nu}{q^2} - g^{\mu\nu}\right)\sum_{n=2}^{\infty}{}' \frac{(2q_{\mu_1})\cdots(2q_{\mu_n})}{(Q^2)^n} \\
&\qquad\qquad \times\left[\sum_q e_q^2 c_n^q\ {}_q\mathcal{O}_n^{\mu_1\cdots\mu_n} + \left(\sum_q e_q^2\right) c_n^g\ {}_g\mathcal{O}_n^{\mu_1\cdots\mu_n}\right] \\
&\quad - i\epsilon^{\mu\nu\alpha\beta} q_\alpha g_{\beta\lambda}\frac{2}{Q^2}\sum_{n=1}^{\infty}{}' \frac{(2q_{\mu_2})\cdots(2q_{\mu_n})}{(Q^2)^{n-1}} \\
&\qquad\qquad \times\left[\sum_q e_q^2 \tilde{c}_n^q\ {}_q\tilde{\mathcal{O}}_n^{\lambda\mu_2\cdots\mu_n} + \left(\sum_q e_q^2\right) \tilde{c}_n^g\ {}_g\tilde{\mathcal{O}}_n^{\lambda\mu_2\cdots\mu_n}\right] \quad . \qquad (2.208)
\end{aligned}
$$

I have ignored the longitudinal contribution, although it appears at the twist-2 level, because it is kinematically suppressed. The gluonic operators, defined by

$$
{}_g\mathcal{O}_n^{\mu_1\cdots\mu_n} = \mathcal{F}^{\alpha(\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_{n-1}}\mathcal{F}_\alpha^{\mu_n)} \qquad (2.209)
$$

$$
{}_g\tilde{\mathcal{O}}_n^{\mu_1\cdots\mu_n} = \mathcal{F}^{\alpha(\mu_1} i\mathcal{D}^{\mu_2}\cdots i\mathcal{D}^{\mu_{n-1}} i\tilde{\mathcal{F}}_\alpha^{\mu_n)} \quad , \qquad (2.210)
$$

are related to the gluon distributions in exactly the same way as their quark counterparts. These operators are also twist-2 and mix with the singlet quark operators, as we will see below. The coefficients of the expansion are given by[112]

$$
\begin{aligned}
c_n^q &= 1 - \frac{\alpha_s C_F}{4\pi}\left[9 - \frac{8}{n} + \frac{2}{n+1} + 4S_2(n-1) - 4T_1^1(n-1)\right. \\
&\qquad\qquad \left. - S_1(n-1)\left(3 + \frac{2}{n} + \frac{2}{n+1}\right)\right] \quad , \qquad (2.211) \\
\tilde{c}_n^q &= 1 - \frac{\alpha_s C_F}{4\pi}\left[9 - \frac{8}{n+1} + \frac{2}{n} + 4S_2(n-1) - 4T_1^1(n-1)\right. \\
&\qquad\qquad \left. - S_1(n-1)\left(3 + \frac{2}{n} + \frac{2}{n+1}\right)\right] \quad , \qquad (2.212) \\
c_n^g &= \frac{\alpha_s T_F}{2\pi}\left[-\frac{2}{n+2} - 2S_1(n-1)\left(\frac{1}{n} - \frac{2}{n+1} + \frac{2}{n+2}\right)\right] \quad , \qquad (2.213) \\
\tilde{c}_n^g &= \frac{\alpha_s T_F}{2\pi}\left[2 + 2S_1(n-1)\right]\left(\frac{1}{n} - \frac{2}{n+1}\right) \quad , \qquad (2.214)
\end{aligned}
$$

[111] The factor e_q^2 does not appear as the quark coefficient to all orders. At higher orders, it is only the singlet and nonsinglet distributions which make sense. For these distributions, one can factor out an electromagnetic charge factor to all orders. Since the distinction is irrelevant at this order, I have not made it.

[112] These coefficients do not agree with those in [62] because their choice of γ_5 scheme is different from mine.

where I have introduced

$$S_j(n) \equiv \sum_{i=1}^{n} \frac{1}{i^j} \ ,$$
$$T_j^k(n) \equiv \sum_{i=1}^{n} \frac{S_j(i)}{i^k} \ . \tag{2.215}$$

These coefficients can be obtained simply by expanding the hard scattering kernels about $x_B = \infty$. Note the implication that the kernels are finite in this limit. This is one of the requirements of a local operator product expansion.

Since the moments of the distribution functions are mathematically equivalent to the functions themselves,[113] it is interesting to study their evolution. According to (2.126), we have

$$\mu^2 \frac{d}{d\mu^2} a_n^a(\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi} \sum_{a'} \int_0^1 dz \, z^{n-1} \, \mathcal{P}_{a' \to a}(z) \, a_n^{a'}(\mu^2) \ . \tag{2.216}$$

Hence the evolution of the moments is governed by the *anomalous dimension*

$$\gamma_{aa'}^n \equiv \int_0^1 dz \, z^{n-1} \, \mathcal{P}_{a' \to a}(z) \ . \tag{2.217}$$

In the nonsinglet sector, we have the simple result

$$\gamma_{NS}^n = C_F \left[\frac{3}{2} + \frac{1}{n(n+1)} - 2S_1(n) \right] \ , \tag{2.218}$$
$$a_n^{NS}(\mu^2) \sim \left(\frac{\mu^2}{\mu_0^2} \right)^{\alpha_s \gamma_{NS}^n / 2\pi} a_n^{NS}(\mu_0^2) \ . \tag{2.219}$$

This explains why γ is called an anomalous dimension. a_n is dimensionless, but it varies with scale as though it had dimension $\alpha_s \gamma_{NS}^n / \pi$.[114] γ is related to the renormalization constant required to remove its ultraviolet divergences. One can think of it as the residue leftover by

[113] The Legendre polynomials are finite linear combinations of $\{x_i\}$, so we can always trade Mellin moments for Legendre moments. Since the latter form a complete basis of L_2 on the interval $[-1, 1]$, we can fix the distribution functions in L_2. To do better, we determine endpoint contributions from the conservation requirements, Eqs.(2.130) and (2.150).

[114] If we take the leading variation of $\alpha_s(\mu^2)$ with μ^2 into account, the factor

$$\left(\frac{\mu^2}{\mu_0^2} \right)^{\alpha_s \gamma_{\rm NS}^n / 2\pi} \to \left(\frac{\alpha_s(\mu_0^2)}{\alpha_s(\mu^2)} \right)^{2\gamma_{\rm NS}^n / b_0} \ . \tag{2.220}$$

However, since we only use the leading value of $\gamma_{\rm NS}^n$, this represents only a partial resummation of higher order effects. This so-called 'leading log' approximation is used extensively in fits to data. In any case, the two factors coincide in the limit of small α_s.

renormalization. Note that $\gamma_{NS}^n \leq 0$ for all values of n. This implies that as the scale is raised, less and less of the operators in our expansion are important. It reflects the fact that as we look closer and closer at a parton distribution, more and more partons appear. The more partons there are to share the total momentum of our target, the less momentum each one has. Hence as the scale is increased, the parton distributions become sharply peaked at the origin. Since each higher moment penalizes small-x contributions more than the last, it sees less of the bulk of the distribution. One more important aspect of our result is that $\gamma_{NS}^1 = 0$. Since the first moment of the quark distribution is conserved, its anomalous dimension is constrained to be zero.

In the singlet sector, the quark and gluon operators mix with each other. Mathematically, this is because they contain divergences that they can't absorb by themselves. Physically, it is because 'quark' and 'gluon' are not good quantum numbers in this context. If we want something to evolve independently of other things, it had better be an eigenstate of the evolution operator. Since the gluon and singlet quark fields fluctuate into one another, it doesn't really make sense to separate them in this way. To find out what *does* make sense, one can calculate the anomalous dimension matrix,

$$\mu^2 \frac{d}{d\mu^2} \begin{pmatrix} a_n^s \\ a_n^g \end{pmatrix} = \frac{\alpha_s(\mu^2)}{2\pi} \begin{pmatrix} \gamma_s^n & \gamma_{sg}^n \\ \gamma_{gs}^n & \gamma_g^n \end{pmatrix} \begin{pmatrix} a_n^s \\ a_n^g \end{pmatrix} \quad , \tag{2.221}$$

where

$$\gamma_s^n = C_F \left[\frac{3}{2} + \frac{1}{n(n+1)} - 2S_1(n) \right] \tag{2.222}$$

$$\gamma_{sg}^n = 2n_f T_F \left[\frac{1}{n} - \frac{2}{(n+1)(n+2)} \right] \tag{2.223}$$

$$\gamma_{gs}^n = C_F \left[\frac{1}{n+1} + \frac{2}{n(n-1)} \right] \tag{2.224}$$

$$\gamma_g^n = 2C_A \left[\frac{1}{n(n-1)} + \frac{1}{(n+1)(n+2)} - S_1(n) \right] - \frac{4n_f T_F - 11C_A}{6} \quad , \tag{2.225}$$

and diagonalize it. The eigenstates of this matrix evolve independently of one another. If we now invert the Mellin transform, we will find ourselves with two singlet distributions which make sense (at this order) independently of one another. In particular, the evolution of the second moment is governed by the matrix

$$\gamma^2 = \begin{pmatrix} -\frac{4}{3} C_F & \frac{2}{3} n_f T_F \\ \frac{4}{3} C_F & -\frac{2}{3} n_f T_F \end{pmatrix} \quad . \tag{2.226}$$

This matrix has the eigenvalues

$$\gamma_1^2 = 0 \qquad\qquad \gamma_2^2 = -\frac{4}{3}\,C_F - \frac{2}{3}\,n_f T_F \quad . \tag{2.227}$$

Since the second eigenvalue is negative, the associated eigenvector will be 'run into the ground' as the scale is increased and the only relevant combination becomes that associated with the eigenvalue zero. Since the second moment of our distributions represents the fraction of + momentum carried by quarks and gluons, this eigenvector tells us these fractions at asymptotically large momentum scales :

$$P_q(\infty) = \frac{2N_c n_f T_F}{2(N_c^2-1)+2N_c n_f T_F} = \frac{3n_f}{16+3n_f} \tag{2.228}$$

$$P_g(\infty) = \frac{2(N_c^2-1)}{2(N_c^2-1)+2N_c n_f T_F} = \frac{16}{16+3n_f} \quad . \tag{2.229}$$

This result was first discovered by D. Gross and F. Wilczek in 1974 [65]. Since there are (N_c^2-1) gluons each with two physical polarization states and $N_c n_f$ quarks each with two polarizations and 'strength' T_F, this result has the simple interpretation of equal momentum sharing among all of the relevant degrees of freedom. Corrections to this result vanish as

$$\left(\frac{\mu^2}{\mu_0^2}\right)^{-(16/9+n_f/3)\alpha_s/2\pi} \tag{2.230}$$

as the scale is increased.

The polarized matrix,

$$\tilde{\gamma}^n = \begin{pmatrix} \tilde{\gamma}_s^n & \tilde{\gamma}_{sg}^n \\ \tilde{\gamma}_{gs}^n & \tilde{\gamma}_g^n \end{pmatrix} \quad , \tag{2.231}$$

$$\tilde{\gamma}_s^n = \tilde{\gamma}_{NS}^n = \gamma_s^n \tag{2.232}$$

$$\tilde{\gamma}_{sg}^n = 2n_f T_F \left(\frac{2}{n+1} - \frac{1}{n}\right) \tag{2.233}$$

$$\tilde{\gamma}_{gs}^n = C_F \left(\frac{2}{n} - \frac{1}{n+1}\right) \tag{2.234}$$

$$\tilde{\gamma}_g^n = 2C_A \left[\frac{2}{n(n+1)} - S_1(n)\right] - \frac{4n_f T_F - 11C_A}{6} \quad , \tag{2.235}$$

is quite unrelated to γ^n. This is because the separate helicity of the gluon and singlet quark fields also does not make sense by itself; only the combination is important. Since this is in a different sector of the theory, there is no reason that the two eigenstates should be related. In

particular, the unpolarized anomalous dimension matrix in the singlet sector is divergent for $n = 1$ owing to the fact that the first moment of the gluon distributions is either *not* local[115] or *not* gauge-invariant.[116] However, the polarized matrix

$$\tilde{\gamma}^1 = \begin{pmatrix} 0 & 0 \\ \frac{3}{2} C_F & \frac{1}{6}(11 C_A - 4 n_f T_F) \end{pmatrix} \tag{2.236}$$

is a singular but perfectly finite object. This matrix implies that the singlet helicity is conserved[117] while the 'gluon helicity' *grows* with the *positive* anomalous dimension

$$\tilde{\gamma}_g^1 = \frac{33 - 2 n_f}{6} \ . \tag{2.237}$$

This number is, of course, familiar to us as the leading coefficient of the β-function. It can be shown that this behavior is canceled by an equally large orbital contribution in any truly physical quantities[118] [66]. The steady-state solution,

$$\tilde{a}_1^s = \frac{4 n_f T_F - 11 C_A}{6} \ , \tag{2.238}$$

$$\tilde{a}_1^g = \frac{3}{2} C_F \ , \tag{2.239}$$

is not as useful as the unpolarized result because we have no way to normalize it short of experiment.

2.6 Summary of DIS

In this chapter, we have studied one of the most important tools of modern particle physics - deep-inelastic scattering. In a special kinematic limit, this process can be considered incoherent; one can calculate the cross-section independently for each constituent of a complicated bound state and simply average over the constituents to obtain the total cross-section. This is due to a separation of scales between the soft physics that dominates the bound state interactions and the hard physics relevant to the virtual photon scattering. Using analytic arguments

[115] Since factors of x turn into derivatives, we can think of factors of $1/x$ as turning into integrals.

[116] Instead of living with a nonlocal distribution, we could choose to of cancel the x by working in the light-cone gauge. This process leads to a local operator, but one that has no readily accessible physical interpretation.

[117] The anomaly does not make its presence known until the next order.

[118] Note that according to (2.214), this first moment of this distribution does *not* appear in polarized DIS. All higher moments have *negative* anomalous dimensions, as they should.

about the generation of singularities, one can show that this *factorization* of hard and soft scales persists to all orders in perturbation theory.

The weighting of the average over constituents is done via parton distribution functions, which can be shown within QCD to be related to probabilities for finding partons within our target hadron. Although these distributions are nonperturbative and cannot be calculated reliably using contemporary methods, they are process-independent and appear over and over again in different experiments. As such, these distributions can be measured in one experiment and used to predict the outcome of another as a test of the theory. Moreover, the information on nonperturbative hadronic wavefunctions they contain is interesting in itself. This information can be used to test the accuracy of models, gain insight into hadronic structure, and help make contact between the infinitesimal world of QCD degrees of freedom and the 'macroscopic' world of hadronic interactions.

Scaling, a somewhat robust model-independent prediction of current algebra, is broken by the evolution of these parton distributions. Completely calculable within perturbation theory, violations of scaling in deep-inelastic scattering provide a very useful test of the accuracy of QCD which is not obscured by hadronic effects we do not understand. Experimental observation of scaling violations agrees with the predictions of QCD to better than 10% over a span in Q^2 of four orders of magnitude and in x_B of one order of magnitude.

Expanding our amplitudes in the unphysical region of large x_B, we can use DIS to derive a local operator expansion for the product of two electromagnetic currents. This expansion not only allows us to study the analytic structure of the amplitude itself, but also provides a useful playing field on which to test various assumptions about the behavior of operators in a quantum field theory. Furthermore, several of the operators that appear in the expansion are important in their own right as representatives of flavor currents, axial currents, and the stress-energy tensor of QCD. We will find this aspect very useful in motivating the topic of the next chapter.

As a whole, deep-inelastic scattering has made tremendous contributions to many aspects of modern physics.

Chapter 3

Deeply Virtual Compton Scattering

In a recent paper, X. Ji introduced deeply virtual Compton scattering (DVCS) as a probe to a novel class of "off-forward" parton distributions (OFPD's) [34]. DVCS is a process in which a highly virtual photon scatters on a nucleon target (polarized or unpolarized), producing an exclusive final state consisting of a high-energy real photon and a slightly recoiled nucleon.

Several interesting theoretical papers which study the DVCS process further have since appeared in the literature. In Ref. [67], the single-quark scattering was recalculated using a different, but equivalent definition of the parton distributions. The evolution equations of the distributions were derived and some general aspects of factorization were discussed. In Ref. [68], the evolution equations for OFPD's were derived and the leading-twist DVCS cross sections were calculated at order α_s^0. Some past and recent studies of OFPD's can be found in [69]. The DVCS process was considered as a limit of unequal mass Compton scattering, which was studied from the point of view of the operator product expansion, in Ref. [70]. Some early studies of unequal mass Compton processes can be found in Refs. [71, 72]. $\mathcal{O}(\alpha_s)$ corrections to DVCS were studied in Refs. [73, 74, 75] from different perspectives.

The main motivation for the present study is to see if the theoretical basis for the DVCS process is up to par with other well-known perturbative QCD processes. More explicitly, we discuss the existence of a factorization theorem for this process. For general two virtual photon processes in the Bjorken limit, factorizability is suggested by the analysis of DIS in the last chapter. In the case of DVCS, where one of the photons is onshell, the situation could be different. Potential infrared problems can arise because of the additional light-like vector in this special kinematic limit. However, it is believed that these complications will not ruin the

factorization properties [68].

The material presented in this chapter was originally published by X. Ji and I in [35]. It is structured as follows. In the first section, I will outline one of the theoretical motivations for the study of DVCS. The kinematics of this process allow measurement of certain matrix elements which are related to the *total* angular momentum carried by quarks in the proton. Since the angular momentum carried by quark *spin* is measured in DIS, DVCS can help isolate the contribution due to quark *orbital* angular momentum. This quantity can give us great insight into the wavefunction of the proton. Section 3.2 is devoted to the kinematics of DVCS and the introduction of OFPD's which embody the nonperturbative physics at leading twist in DVCS.

To see factorization at work, it is instructive to work out one-loop examples. We will do this explicitly in Section 3.3. As in Section 2.2, we consider the unphysical process of DVCS on onshell quark and gluon targets. For completeness we consider both the symmetric and antisymmetric parts of the amplitudes, which are related to helicity-independent and dependent parton distributions, respectively. The only omission is the gluon helicity flip amplitude, which is discussed in Ref. [76]. We will see that the collinear infrared divergences can be interpreted as the one-loop perturbative parton distributions in Section 3.4. This property is independent of the special kinematic limit of DVCS.

A general proof of factorization in DVCS was first given by Radyushkin in his approach based on the α-representation [67]. Here, we give an alternative proof using the tools introduced in the last chapter. According to these, the infrared sensitive contributions in a generic Feynman diagram can be represented by reduced diagrams. In Section 3.5, we will see that the leading reduced diagrams for DVCS are identical to those present in DIS. Therefore, the proof of factorization at this twist level is identical to that for DIS.

The factorization properties of general two virtual photon processes can be summarized beautifully in terms of Wilson's operator product expansion. This expansion requires operators with total derivatives to describe the off-forward nature of the process [70, 71, 72]. In Section 3.6, I show how we can convert our one-loop results into Wilson coefficients of the twist-2 operators in the $\overline{\text{MS}}$ scheme. The resulting operator product expansion is valid for forward as well as a certain class of off-forward matrix elements.

A summary and conclusion of this chapter appears in Section 3.7.

3.1 Motivation

One of the most interesting contemporary theoretical studies in nuclear physics involves the origin of nucleon spin. Along with the discovery that protons are composite objects comes the realization that their spins must somehow be generated dynamically by QCD. As described in the Introduction, the earliest quark models saw hadronic spin as a simple sum of the spins of the constituents. The three-quark structure of baryons led to spins of 1/2 and 3/2 for ground states, with orbital excitations providing higher half-integer spins. However, as viewed from QCD, these models are fundamentally flawed. They make no mention either of the sea quarks, which certainly must be taken into account in a full-fledged quantum field theory, or of the gluons, which are expected to be responsible for the quark interactions. Furthermore, they tacitly assume that the ground state contains no orbital excitations. While this is certainly true in non-relativistic quantum mechanics, there is no reason to expect it to be the case in a highly relativistic quantum field theory. These objections motivate us to take another look at the origins of nucleon spin, this time from the standpoint of QCD.

As we have seen in the last chapter, the polarized quark density, $\tilde{f}_{q'/q}(x)$, represents the difference between the probability densities for positive and negative helicity quarks carrying momentum fraction x. This promotes the identification

$$s_{q'/T} = \frac{1}{2}\int dx \tilde{f}_{q'/T}(x) \tag{3.1}$$

of the first moment of the polarized quark distribution with the amount of target spin carried by quark helicity. We will see below that this relation is indeed correct. This contribution has been measured in several experiments, most notably the European Muon Collaboration (EMC) in 1987, with the result that quark spin accounts for only $12 \pm 17\%$ of the proton's spin at a scale of approximately 10 GeV^2.[1] This completely counterintuitive result caused the so-called 'spin-crisis' and put to bed any notion that the naive quark model could be used for

[1]The fit used to extract this value depends on an assumption of $SU(3)$ flavor symmetry. This assumption accounts for much of the quoted error bar. A recent global analysis of the relevant data can be found in Ref. [77].

anything more than the crudest of calculations. Apparently, $\sim 85\%$ of nucleon spin must be attributed to orbital and gluonic effects. To further classify the spin sources, it was proposed that the first moment of the spin-dependent gluon distribution contains information on the contribution due to gluon spin. Unfortunately, this is not the case. The subtleties associated with gauge invariance forbid us from separating the gluon contribution to hadronic spin into separate pieces representing 'orbital angular momentum' and 'helicity angular momentum'. These two quantities do not make sense separately for gauge fields.[2]

A correct analysis of nucleon spin begins with the Pauli-Ljubanskii vector

$$\hat{W}^\mu = -\frac{1}{2}\epsilon^{\mu\nu\alpha\beta}\hat{P}_\nu \hat{J}_{\alpha\beta} \tag{3.2}$$

introduced in Appendix B.2. As in the appendix, we classify particle states by eigenvalues of the momentum operator $\hat{P}$ and one of the components of the Pauli-Ljubanskii vector. In the rest frame of a nucleon, $p^\mu = (M, 0, 0, 0)$, the 3-component of $\hat{W}$ is $M\hat{J}^{12}$. The statement that a proton has spin-1/2 means that there are two eigenstates of $\hat{W}^3$:

$$\begin{aligned} \hat{W}^3 \left| p^\mu, \uparrow \right\rangle &= \quad \frac{1}{2} M \left| p^\mu, \uparrow \right\rangle \\ \hat{W}^3 \left| p^\mu, \downarrow \right\rangle &= -\frac{1}{2} M \left| p^\mu, \downarrow \right\rangle \quad . \end{aligned} \tag{3.3}$$

The normalization of our states is such that

$$\langle p', h' \mid p, h \rangle = 2p^0 \delta_{hh'} (2\pi)^3 \delta\left(\vec{p} - \vec{p}\,'\right) \quad , \tag{3.4}$$

where h and h' represent $\uparrow$ and $\downarrow$, so

$$\left\langle p, \uparrow \left| \hat{J}^{12} \right| p, \uparrow \right\rangle = \frac{1}{2}\, 2M(2\pi)^3 \delta^{(3)}(0) \quad . \tag{3.5}$$

Our choice of the 3-direction was arbitrary, as was our choice of frame; in general, for an arbitrary spin direction s^μ and nucleon momentum p^μ, we have[3]

$$\left\langle ps \left| \hat{W}^\mu \right| ps \right\rangle = M s^\mu\, 2p^0 (2\pi)^3 \delta^{(3)}(0) \quad . \tag{3.6}$$

[2] This fact is exemplified by the discussion following Eq.(2.237). The separation of gluon spin and orbital angular momentum leads to this positive anomalous dimension.

[3] My normalization is such that $s^2 = -1/4$ rather than -1.

In order to understand how the spin of a proton is generated, we need to decompose it into several contributions. As shown in Appendix D.2, the spatial rotation generators[4]

$$\hat{J}^{ij} = \int d^3x \left(J^{ij}_{qs} + J^{ij}_{qo} + J^{ij}_{g} \right) \tag{3.7}$$

$$J^{ij}_{qs} = \frac{1}{2}\, \psi^\dagger \sigma^{ij} \psi \tag{3.8}$$

$$J^{ij}_{qo} = \psi^\dagger \left(x^i\, i\mathcal{D}^j - x^j\, i\mathcal{D}^i \right) \psi \tag{3.9}$$

$$J^{ij}_{g} = -\left(x^i \mathcal{F}^{j\alpha} - x^j \mathcal{F}^{i\alpha} \right) \mathcal{F}^0_{\ \alpha} \tag{3.10}$$

split nicely into three separately gauge-invariant pieces. Writing $\hat{S} \equiv \hat{W}/M$ and substituting these expressions into the definition of $\hat{W}$, we find that[5]

$$\hat{\vec{S}} = \int d^3x \left(\vec{S}_{qs} + \vec{S}_{qo} + \vec{S}_{g} \right) \tag{3.11}$$

$$\vec{S}_{qs} = \frac{1}{2}\, \overline{\psi}\, \vec{\gamma}\, \gamma_5\, \psi \tag{3.12}$$

$$\vec{S}_{qo} = \psi^\dagger \left(\vec{x} \times -i\vec{\mathcal{D}} \right) \psi \tag{3.13}$$

$$\vec{S}_{g} = \vec{x} \times \left(\vec{\mathcal{E}} \times \vec{\mathcal{B}} \right) \ , \tag{3.14}$$

where $\mathcal{E}$ and $\mathcal{B}$ are the chromo-electric and -magnetic fields, respectively, defined by $\mathcal{E}^i \equiv -\mathcal{F}^{0i}$ and $\mathcal{B}^i \equiv -\tilde{\mathcal{F}}^{0i}$. Equation (3.12) has the obvious interpretation as the contribution from quark intrinsic angular momentum since it gives a different sign to the left- and right-handed components of the quark fields. Similarly, (3.13) is what we expect from an orbital contribution to angular momentum once we recognize $-i\vec{\mathcal{D}}$ as the momentum operator in a gauge theory. The remaining term is attributed to gluons. The nonlinear behavior of the gauge field under gauge transformations prevents and further decomposition. The canonical approach to angular momentum discussed in Appendix D.2 shows conclusively that this separation is unavoidably gauge-dependent. However (3.14) does have a nice physical interpretation. Since $\vec{\mathcal{E}} \times \vec{\mathcal{B}}$ is the Poynting vector representing classical energy flow, this term has the expected form of an angular momentum. In fact, it is exactly what we would write down for the angular momentum of a classical electromagnetic field.

[4] As always, neglected $SU(3)$ and flavor indices are fully contracted.

[5] As usual, a vector hat on a 4-vector denotes the 3-vector formed from its spatial components - with indices *upstairs*.

The contribution of each operator in our decomposition is given by the matrix element

$$\begin{aligned}\left\langle ps \left| \int d^3x \, \vec{S}_i(x) \right| ps \right\rangle &= (2\pi)^3 \delta^{(3)}(0) \left\langle ps \left| \vec{S}_i(0) \right| ps \right\rangle \\ &= 2Ma_i \vec{s} (2\pi)^3 \delta^{(3)}(0) \ . \end{aligned} \tag{3.15}$$

The last equality follows from rotational invariance and the fact that QCD conserves parity.[6] The constants a_i characterize the contribution of operator i to the nucleon spin. In view of (3.6) and (3.11), we have the sum rule

$$a_{qs} + a_{qo} + a_g = 1 \ . \tag{3.16}$$

At this point, we are free to use any means necessary to extract the a_i's. These objects are inherently nonperturbative and so must be measured in experiment (at least in this stage of the game).

As we saw in the last chapter, certain matrix elements show up naturally in physical scattering processes. We have only to calculate their coefficients and perform the measurements. In particular, hadronic matrix elements of the operator $\overline{\psi} \not{n} \gamma_5 \psi$ can be measured in polarized DIS. Exploiting Lorentz invariance[7], we can extract a_{qs} from these measurements :

$$\begin{aligned}\left\langle ps \left| \overline{\psi} \gamma^\mu \gamma_5 \psi \right| ps \right\rangle &= 4Ma_{qs} s^\mu \\ \left\langle ps \left| \overline{\psi} \not{n} \gamma_5 \psi \right| ps \right\rangle &= 4ha_{qs} \ , \end{aligned} \tag{3.17}$$

where in the last line we have chosen a frame in which the hadron is moving ultra-relativistically along the 3-axis. In light of Equation (2.79), a_{qs} is just the first moment of the polarized singlet quark distribution,[8]

$$a_{qs} = \int_{-1}^{1} dx \, \tilde{f}_{S/T}(x) = \Delta N_{S/T} \ . \tag{3.18}$$

As mentioned in Section 2.3, this quantity is not conserved by the interactions of QCD. It a scale-dependent object, like the QCD coupling. In fact, all three of the operators in Equation (3.11) evolve with scale. This is a completely natural phenomenon. Only the sum

[6] The left hand side of this equation is odd (doesn't change sign) under parity, so it must be proportional to a vector which is also odd under parity. The only such vector in this problem is $\vec{s}$.

[7] and the fact that QCD conserves parity

[8] The antiquark contribution must, of course, be included. As expected, it simply extends the integration into the negative real axis according to (2.75).

of all three contributions must be scale-independent. This means that it makes no sense to ask how much spin is carried, for example, by quark orbital angular momentum. Only when a scale is specified do questions such as this have meaning. At high scales, one can calculate the evolution of these operators in perturbation theory. As the scale runs to infinity, only the eigenvector of the anomalous dimension matrix with the smallest (negative) eigenvalue becomes relevant. In this limit, one can show [34, 66] that the percentage of spin carried by quarks and gluons becomes

$$S_q(\infty) = \frac{2N_c n_f T_F}{2(N_c^2-1) + 2N_c n_f T_F} = \frac{3n_f}{16+3n_f} \tag{3.19}$$

$$S_g(\infty) = \frac{2(N_c^2-1)}{2(N_c^2-1) + 2N_c n_f T_F} = \frac{16}{16+3n_f} \quad . \tag{3.20}$$

This result is identical to that found in Section 2.5 for the distribution of longitudinal momentum! Apparently, it is a somewhat general feature of the asymptotic limit. This is not completely surprising since its interpretation is simply benevolent sharing among partons. However, what this result implies about the distribution at moderate scales is wholly unclear.

As mentioned above, experimental results indicate that the quark spin contribution to proton spin is $\sim 15\%$ at moderate probing scales (10 GeV^2). This implies that a full 85% of the nucleon spin is attributed to gluonic and orbital effects at these scales. Unfortunately, the other operators are not readily available for comment. In order to measure a_{qo} directly, one would have to find a process which is sensitive only to the quark orbital angular momentum - a formidable, if not impossible, task. Luckily, all is not lost. The primordial form of $\hat{J}$ involves the energy-momentum tensor of QCD. Hopefully, we can entice this somewhat less exotic operator to help us extract the information we desire.

We have already studied a process in which matrix elements of the energy-momentum tensor appear - unpolarized DIS. According to (2.99) and (2.106), the second moments of the unpolarized nucleon structure functions are proportional to its diagonal matrix elements. Unfortunately, things are not as simple here as they were in the quark spin sector. The operator that we want diagonal matrix elements of is

$$\hat{J}^{ij} = \int d^3x \left(x^i \Theta^{0j} - x^j \Theta^{0i} \right) \quad . \tag{3.21}$$

Writing

$$\begin{aligned}
\langle \hat{J}^{ij} \rangle &= \lim_{\Delta^\mu \to 0} \left\langle P + \frac{\Delta}{2} \left| \hat{J}^{ij} \right| P - \frac{\Delta}{2} \right\rangle \\
&= \lim_{\Delta^\mu \to 0} \int d^3x \, e^{-i\vec{x}\cdot\vec{\Delta}} \left\langle P + \frac{\Delta}{2} \left| x^i \Theta^{0j} - x^j \Theta^{0i} \right| P - \frac{\Delta}{2} \right\rangle \\
&= \lim_{\Delta^\mu \to 0} \int d^3x \left\langle P + \frac{\Delta}{2} \left| \left(i \frac{\partial}{\partial \Delta^i} e^{-i\vec{x}\cdot\vec{\Delta}} \right) \Theta^{0j} - (i \leftrightarrow j) \right| P - \frac{\Delta}{2} \right\rangle \\
&= \lim_{\Delta^\mu \to 0} \left[-i \frac{\partial}{\partial \Delta^i} \left\langle P + \frac{\Delta}{2} \left| \Theta^{0j} \right| P - \frac{\Delta}{2} \right\rangle - (i \leftrightarrow j) \right] (2\pi)^3 \, \delta\left(\vec{\Delta}\right) \quad ,
\end{aligned} \tag{3.22}$$

we see that the spatial coordinates in the definition of $\hat{J}$ can be converted into derivatives on Θ's *off-diagonal* matrix elements.

Thanks to the separable form of Θ, this expression for $\langle \hat{J} \rangle$ can also be split into independent quark and gluon pieces. Off-diagonal matrix elements of these two pieces can be separately parameterized by form factors :[9]

$$\begin{aligned}
\left\langle P + \frac{\Delta}{2} \left| \Theta^{\mu\nu}_{q,g} \right| P - \frac{\Delta}{2} \right\rangle = \overline{U}\left(P + \frac{\Delta}{2}\right) \quad & \Big[A_{q,g}(\Delta^2) \gamma^{(\mu} P^{\nu)} \\
& + B_{q,g}(\Delta^2) P^{(\mu} i\sigma^{\nu)\alpha} \Delta_\alpha / 2M \\
& + C_{q,g}(\Delta^2) \left(\Delta^\mu \Delta^\nu - g^{\mu\nu} \Delta^2 \right) / M \\
& + \overline{C}_{q,g}(\Delta^2) M g^{\mu\nu} \Big] U\left(P - \frac{\Delta}{2}\right) \quad .
\end{aligned} \tag{3.23}$$

Here, $(\overline{U})$ U is the free wavefunction for an (outgoing) incoming nucleon. Since we wish to consider a derivative of this object at $\Delta = 0$, only quantities linear in Δ can contribute. Lorentz invariance[10] implies that the scalar $\overline{U}U$ can depend only on Δ^2, so contributions to $\langle \hat{J} \rangle$ can come only from the A and B terms. Using the *Gordon identity*,

$$\begin{aligned}
\overline{U}\left(P + \frac{\Delta}{2}\right) \gamma^{(\mu} P^{\nu)} U\left(P - \frac{\Delta}{2}\right) = \overline{U}\left(P + \frac{\Delta}{2}\right) & \left[P^\mu P^\nu / M \right. \\
& \left. + P^{(\mu} i\sigma^{\nu)\alpha} \Delta_\alpha / 2M \right] U\left(P - \frac{\Delta}{2}\right) \quad ,
\end{aligned} \tag{3.24}$$

it becomes obvious that

$$\langle P | \, \hat{J}^{ij}_{q,g} \, | P \rangle = \frac{1}{2} \left[A_{q,g}(0) + B_{q,g}(0) \right] \overline{U}(P) \, \sigma^{ij} U(P) \, (2\pi)^3 \, \delta^{(3)}(0) \tag{3.25}$$

[9] Conservation of the full energy-momentum tensor implies that $\overline{C}_q = -\overline{C}_g$.

[10] along with the fact that $\Delta \cdot P = 0$ since both external states are on the same mass-shell

in the rest frame of the nucleon. Contracting with $\epsilon^{ijk}/2$ and comparing with (3.6) in the rest frame, we see that the quark and gluon contributions to hadronic spin are given by[11]

$$S_{q,g} = A_{q,g}(0) + B_{q,g}(0) \qquad ; \qquad \sum_{q,g} [A_i(0) + B_i(0)] = 1 \ . \tag{3.26}$$

This is an extremely interesting result. It states that we can obtain information about the spin structure of a hadron from *unpolarized* scattering experiments. This is not a completely crazy idea since polarization concerns the *alignment* of a particle's spin rather than the way it is distributed among its constituents. Still, it is somewhat unexpected that we can obtain such useful information on spin without polarization.

Although this result still does not tell us how to measure the quark orbital contribution to proton spin directly, knowledge of the above quantities coupled with the information obtained from polarized DIS allows us to extract this information. At present, we have very little information on the nonperturbative momentum distribution of partons within hadrons. The distributions measured in DIS give us information only about longitudinal momentum distributions. The quark orbital contribution to the proton spin will give us valuable information on the *transverse* momentum distribution. In addition, since the proton represents the absolute baryonic ground state, information on its orbital distribution tells us information about the structure of this global energy minimum of QCD.

However simple it seems, Equation (3.26) does have a catch. As mentioned above, DIS allows us to probe *diagonal* matrix elements of $\Theta^{\mu\nu}$. According to (3.23), the form factor $A(0)$ can be extracted from these measurements. Unfortunately, the other form factor necessary for our present purpose does not contribute to diagonal matrix elements. In order to extract $B(0)$, we will have to find some way to probe slightly off-diagonal matrix elements of this operator.

We discovered in Section 2.5 that the local operators appearing in the QCD stress-energy tensor also contribute to the light-cone operator product expansion of two electromagnetic currents. Although this expansion is valid only for diagonal matrix elements, it is certainly plausible to assume that the operators of interest will also contribute to a generalized expansion which extends the range of validity. If this is the case, then we can extract information on the off-diagonal matrix elements of Θ by studying off-forward two-photon processes. In fact,

[11] I have used the fact that $U^\dagger \, \vec{\gamma} \, \gamma_5 U = 4M\vec{s}$ in the rest frame.

it is easy to show that *all* of Chapter 2 generalizes quite easily to off-forward processes in which both photons are deeply virtual (though with different virtualities). The calculations are almost identical. However, there is a major philosophical difference between these two processes. Deep-inelastic scattering only becomes a two-photon process when we sum over final states and use the optical theorem to express its cross-section as the imaginary part of the Compton amplitude. Experimentally, it is electron-proton scattering with only stray photons in the initial or final states. On the other hand, the general two-photon processes proposed to help us extract $B(0)$ are intrinsic Compton processes. Since the initial and final states are necessarily different, the imaginary parts of these amplitudes are not related to *any* cross-section. To get the cross-section for a physical process, we have to square *these* amplitudes! Furthermore, in order for these processes to be physical, the initial and final states must be onshell. In DIS, the virtual photon is not a part of the initial or final state; it is exchanged between the electron and the proton during the process. We can generate the initial virtual photon the same way in the present case, but the deeply virtual photon in the 'final' state presents a real problem. It is simply too much to ask for the electron to bear the burden of both virtual photons. Unfortunately, spacelike virtual photons are in no position to bargain. If they are not quickly absorbed by other particles, they simply will not form in the first place.

One way out of this dilemma is to move into a region where the final photon is not spacelike, but timelike. Timelike virtual photons have many options open to them. Depending on their energy, they can decay into many physical final states. However, the kinematics associated with timelike virtual photons requires a large momentum transfer between the initial and final states. Since we intend to extrapolate our form factors to zero momentum transfer, this is not ideal for our purpose.

Alternatively, we can relax the condition that both photons have a large virtuality and allow the final state photon to go onshell. This process, called deeply virtual Compton scattering (DVCS), is the subject of the present chapter. We will find that it is almost ideal for our needs in that it probes certain nonperturbative structure functions, the off-forward parton distributions (OFPD's), whose moments are related to off-diagonal matrix elements of the stress-energy tensor. Furthermore, its kinematics permit the square of the momentum transfer

to vanish with the nucleon mass. This fact allows a smooth extrapolation to our region of interest.

3.2 Kinematics and Parton distributions

Although our ultimate interest is in deeply virtual Compton scattering, we start by considering a general Compton process involving two offshell photons with different virtualities. This and a suitable choice of kinematic variables allow us to exploit the full symmetry of the problem. In the general Compton process, a virtual photon of momentum $q+\Delta/2$ is absorbed by a hadron of momentum $P-\Delta/2$, which then emits a virtual photon with momentum $q-\Delta/2$ and recoils with momentum $P+\Delta/2$. The three independent external momenta can be expanded in terms of the light-cone vectors

$$\begin{aligned} p^\mu &= \left(p^+, 0, 0, p^+\right) \ , \\ n^\mu &= \frac{1}{2p^+}\left(1, 0, 0, -1\right) \ , \end{aligned} \tag{3.27}$$

where the 3-direction is chosen as the direction of the average hadron momentum (P), and two transverse vectors. As in previous chapters, we call the coefficient of p^μ the + component and that of n^μ the − component. Thus we write

$$\begin{aligned} P^\mu &= p^\mu + \frac{M^2 - t/4}{2} n^\mu \ , \\ q^\mu &= -\zeta p^\mu + \frac{Q^2}{2\zeta} n^\mu \ , \\ \Delta^\mu &= -2\xi p^\mu + \xi(M^2 - t/4) n^\mu + \Delta_\perp^\mu \ , \end{aligned} \tag{3.28}$$

where M is the hadron mass (which is taken to be the same for the initial and final hadrons), $t = \Delta^2$, Q^2 is the virtuality of q^μ, ξ is a measure of the difference of the virtualities of the two external photons, ζ is defined as

$$\zeta = \frac{Q^2}{2x_B(M^2 - t/4)} \left(-1 + \sqrt{1 + \frac{4x_B^2(M^2 - t/4)}{Q^2}} \right) , \tag{3.29}$$

and $\Delta_\perp^\mu$ is a vector in the transverse directions which has squared length $-t\left(1-\xi^2\right) - 4\xi^2 M^2$. We have also introduced $x_B = Q^2/(2P\cdot q)$, the analogue of the Bjorken scaling variable in this off-forward process. These expressions limit the range of ξ to

$$\xi^2 \leq \frac{-t}{-t + 4M^2} \ , \tag{3.30}$$

for fixed t, or the range of t to

$$-t \geq \frac{4\xi^2 M^2}{1-\xi^2} \ , \tag{3.31}$$

for fixed ξ.

According to the last section, our region of interest is $t = 0$. In order for the final state photon to be onshell,[12] we must have $\xi \neq 0$. In light of the above expressions, this is only possible if we take the limit $M^2 \to 0$ *before* $t \to 0$. Unfortunately,[13] we do not control the mass of the nucleon. However, it is kinematically allowed to take ξ small so that $-t$ can be much smaller than M^2. Performing several of these measurements will provide a smooth extrapolation to the required values.

In the Bjorken limit, these expressions simplify considerably. Since we consider only the leading twist in this chapter, we may neglect all but the + components of P^μ and Δ^μ (in order to form large scalars, one must dot the + component of a vector with the − component of q). Hence, in the limit $Q^2 \to \infty$ (x_B remaining finite), we may write

$$\begin{aligned} P^\mu &\sim p^\mu \ , \\ q^\mu &\sim -x_B p^\mu + \frac{Q^2}{2x_B} n^\mu \ , \\ \Delta^\mu &\sim -2\xi p^\mu \ . \end{aligned} \tag{3.32}$$

We note that the external invariants have been reduced from six to three by enforcing kinematics and taking the Bjorken limit. We express these three scalars in terms of one mass scale, Q^2, and two dimensionless parameters, x_B and ξ. Apparently, the limit $\xi \to 0$ recovers the DIS results from the last chapter. We will find this to be a useful check of our calculation. The DVCS limit is given by $\xi \to x_B$. As mentioned above, we cannot do DVCS when $\xi = 0$.[14] However, measurements at small x_B may allow $-t$ small enough that the variation to zero holds no surprises. In particular, for the relatively large value $x_B \sim 0.1$, we can have $-t$ as low as ~ 0.04 GeV^2. Since typical hadronic scales are ~ 0.1 GeV^2 and larger, this value of $-t$ is actually quite close to zero.

[12] in the Bjorken limit

[13] Well, maybe fortunately...

[14] Otherwise, we could've just extracted the required information from DIS measurements!

Our goal is to factorize the short and long distance physics of the Compton amplitude[15]

$$T^{\mu\nu} = i \int d^4z\, e^{iq\cdot z} \left\langle P + \frac{\Delta}{2} \left| \mathrm{T} J^\mu \left(\frac{z}{2}\right) J^\nu \left(-\frac{z}{2}\right) \right| P - \frac{\Delta}{2} \right\rangle \tag{3.33}$$

in the Bjorken limit, where $J^\mu = \sum_q e_q \overline{\psi}_q \gamma^\mu \psi_q$ is the electromagnetic current and ψ_q is the bare quark field of flavor q and charge e_q. Note that the operator here is identical to that considered in DIS. Only the external states are different. This immediately implies that all of the diagrams we need to calculate are topologically identical to those of the last chapter. The only difference is in the momenta of the external states.

As before, the nonperturbative contribution to the Compton amplitude in Eq.(3.33) can be expressed in terms of the off-forward parton distributions contained in parton density matrices [68]. For quarks, we define the light-cone correlation function

$$\begin{aligned} \mathcal{M}^q_{\alpha\beta}(x,\xi) &= \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P + \frac{\Delta}{2} \left| \overline{\psi}^q_\beta \left(\frac{\lambda}{2} n\right) \mathcal{G}_n \left(\frac{\lambda}{2} n, -\frac{\lambda}{2} n\right) \psi^q_\alpha \left(-\frac{\lambda}{2} n\right) \right| P - \frac{\Delta}{2} \right\rangle \\ &= \frac{1}{2} F_{q/T}(x,\xi,t) \not{n}_{\alpha\beta} + \tilde{F}_{q/T}(x,\xi,t) \left(\gamma_5 h \not{n}\right)_{\alpha\beta} + \cdots \ , \end{aligned} \tag{3.34}$$

where h is the helicity of the proton. The ellipses denote contributions either of higher twist or chiral-odd structure, which do not contribute to the leading process under consideration. The gauge link, $\mathcal{G}_n$, is defined as in Section 2.3 with a contour that runs along the n direction. Note that this density matrix is just an off-diagonal matrix element of the *same* field operator we considered there.[16]

Isolating the scalar distributions, we find

$$\begin{aligned} &F_{q/T}(x,\xi,t) \qquad (3.35) \\ &\quad = \frac{1}{2} \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P + \frac{\Delta}{2} \left| \overline{\psi}_q \left(\frac{\lambda}{2} n\right) \mathcal{G}_n \left(\frac{\lambda}{2} n, -\frac{\lambda}{2} n\right) \not{n} \psi_q \left(-\frac{\lambda}{2} n\right) \right| P - \frac{\Delta}{2} \right\rangle \ , \\ &\tilde{F}_{q/T}(x,\xi,t) \qquad (3.36) \\ &\quad = \frac{1}{4h} \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P + \frac{\Delta}{2} \left| \overline{\psi}_q \left(\frac{\lambda}{2} n\right) \mathcal{G}_n \left(\frac{\lambda}{2} n, -\frac{\lambda}{2} n\right) \not{n} \gamma_5 \psi_q \left(-\frac{\lambda}{2} n\right) \right| P - \frac{\Delta}{2} \right\rangle \ . \end{aligned}$$

[15] Here, we are actually interested in this amplitude rather than the analogue of $W^{\mu\nu}$.

[16] The difference in definition between this correlation function and that in Section 2.3 (the arguments of the parton fields) is not important there since the matrix element is diagonal. However, here the off-diagonal nature of the matrix element creates a dependence on the positions of the fields. I have chosen the symmetric form for later convenience.

The ever-present renormalization scale, μ, has been suppressed here for brevity. Dependence on the square of the momentum transfer, t, will also largely be ignored in the following as it will not affect most of our discussion.

Before we go on, let us see how these distributions are related to the form factors $A(t)$ and $B(t)$ of the stress-energy tensor. Writing

$$\begin{aligned}\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P+\frac{\Delta}{2}\left|\overline{\psi}_q\left(\frac{\lambda}{2}n\right)\mathcal{G}_n\left(\frac{\lambda}{2}n,-\frac{\lambda}{2}n\right)\gamma^\mu\psi_q\left(-\frac{\lambda}{2}n\right)\right|P-\frac{\Delta}{2}\right\rangle \qquad (3.37)\\ = \overline{U}\left(P+\frac{\Delta}{2}\right)\left[\gamma^\mu\, H_{q/T}(x,\xi,t)+i\sigma^{\mu\alpha}\Delta_\alpha/2M\; E_{q/T}(x,\xi,t)\right]\, U\left(P-\frac{\Delta}{2}\right)\ ,\end{aligned}$$

we see that

$$\begin{aligned}F_{q/T}(x,\xi,t) = \frac{1}{2}\,\overline{U}\left(P+\frac{\Delta}{2}\right)\left[\not{n}\, H_{q/T}(x,\xi,t)\right.\\ \left.+i\sigma^{\mu\alpha}n_\mu\Delta_\alpha/2M\; E_{q/T}(x,\xi,t)\right]\, U\left(P-\frac{\Delta}{2}\right)\ . \qquad (3.38)\end{aligned}$$

On the other hand, the second x-moment of $F_{q/T}(x,\xi,t)$ is an off-diagonal matrix element of the quark contribution to the stress-energy tensor :[17]

$$\int dx\, x\, F_{q/T}(x,\xi,t) = \frac{1}{2}\,\left\langle P+\frac{\Delta}{2}\left|\overline{\psi}_q\,\not{n}\, n\cdot i\overleftrightarrow{\mathcal{D}}\,\psi_q\right|P-\frac{\Delta}{2}\right\rangle\ . \qquad (3.39)$$

Comparison with (3.23) gives

$$\begin{aligned}\int dx\, x\, F_{q/T}(x,\xi,t) = \frac{1}{2}\,\overline{U}\left(P+\frac{\Delta}{2}\right)\left[\not{n}\, A_q(t)+i\sigma^{\mu\nu}n_\mu\Delta_\nu/2M\; B_q(t)\right.\\ \left.+4\xi^2/M\; C_q(t)\right]\, U\left(P-\frac{\Delta}{2}\right)\ . \qquad (3.40)\end{aligned}$$

The Gordon identity, Equation (3.24), allows us to deduce

$$\int dx\, x\, H_{q/T}(x,\xi,t) = A_q(t)+4\xi^2\, C_q(t)\ , \qquad (3.41)$$

$$\int dx\, x\, E_{q/T}(x,\xi,t) = B_q(t)-4\xi^2\, C_q(t)\ , \qquad (3.42)$$

from which we immediately see that

$$S_q = A_q(0)+B_q(0) = \int dx\, x\, \left[H_{q/T}(x,\xi,0)+E_{q/T}(x,\xi,0)\right]\ . \qquad (3.43)$$

[17]Note that here the '$\leftrightarrow$' adorning $\mathcal{D}$ is quite important. For off-diagonal matrix elements, it must not be forgotten that $\Theta^{\mu\nu}$ is a benevolent operator which treats both external states with the utmost care and respect.

Luckily, the quark contribution to angular momentum is not sensitive to ξ. The lower bound on $-t$ causes us to want ξ small anyway, but it's certainly nice to know that we need not perform an extrapolation to $\xi = 0$; only $t = 0$ is strictly required.

Many properties of the distributions $E(x,\xi,t)$ and $H(x,\xi,t)$, as well as their sister distributions in the polarized, chiral-odd, and gluon sector are summed up nicely in Ref. [78]. There, one can find details of their parton model interpretation, form factor decompositions for their moments, models representing their behavior, and sum rules that constrain them.

The contribution of gluons to DVCS requires off-forward distribution functions. In analogy with Section 2.3, these are written[18]

$$F_{g/T}(x,\xi,t) \tag{3.44}$$
$$= -\frac{1}{2x}\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P+\frac{\Delta}{2}\left| \mathcal{F}^{+\alpha}\left(\frac{\lambda}{2}n\right) \mathcal{G}_n\left(\frac{\lambda}{2}n,-\frac{\lambda}{2}n\right) \mathcal{F}^{+}_{\ \alpha}\left(-\frac{\lambda}{2}n\right)\right| P-\frac{\Delta}{2}\right\rangle \ ,$$
$$\tilde{F}_{g/T}(x,\xi,t) \tag{3.45}$$
$$= -\frac{i}{4hx}\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P+\frac{\Delta}{2}\left| \mathcal{F}_a^{+\alpha}\left(\frac{\lambda}{2}n\right) \mathcal{G}_n\left(\frac{\lambda}{2}n,-\frac{\lambda}{2}n\right) \tilde{\mathcal{F}}_b^{+}{}_{\alpha}\left(-\frac{\lambda}{2}n\right)\right| P-\frac{\Delta}{2}\right\rangle \ .$$

In this chapter, the fact that the external wavefunctions do not correspond to the same momenta implies that the identity $u\overline{u} = (1+2h\gamma_5)\not{p}/2$ cannot be used. Instead, we calculate the amplitudes via a direct substitution of the expansions of the parton distributions. Interpreting these functions as their leading order values (which are simple δ-functions for partonic targets), we will see that the one-loop calculation generates their evolution. For example, the polarized quark amplitude is obtained by tracing with $\gamma_5 h\not{p}$ because this is the structure that appears multiplying $\tilde{F}_{q/T}$ in (3.34). The subtleties associated with defining γ_5 in d dimensions make the polarized result ambiguous. To resolve the ambiguity, one must use a well-defined γ_5 scheme. Several different schemes can be found in the literature;[19] the prescription used here is equivalent to that introduced by 't Hooft and Veltman [10]. The fact that this choice favors certain directions over others makes it necessary to multiply the polarized contribution of diagram 3.1e by the factor $(1+\delta\epsilon)$,[20] where δ is determined by the requirement that the nonsinglet axial current is conserved, whenever one wishes to consider the chiral properties of

[18] Here, Δ leads to an additional twist-2 term involving gluon helicity flip. This term is considered in [76].

[19] See Appendix E.

[20] This amounts to a redefinition of $\gamma^\alpha\gamma_5\gamma_\alpha$.

a theory [79]. Since we are not concerned with the chiral nature of QCD at the moment, we will ignore this technicality.

In the gluon sector, the fact that the conceptual idea of removing one gluon and the actual gauge invariant distributions that describe this process do not coincide leads to extra factors. In this case, one can show that

$$\int \frac{d\lambda}{2\pi} e^{-i\lambda x} \left\langle P + \frac{\Delta}{2} \middle| \mathcal{A}_\perp^\mu \left(\frac{\lambda}{2} n\right) \mathcal{A}_\perp^\nu \left(-\frac{\lambda}{2} n\right) \middle| P - \frac{\Delta}{2} \right\rangle$$
$$= -\frac{x}{x_+ x_-} \left[\frac{2}{d-2} g_\perp^{\mu\nu} F_{g/T}(x,\xi) - i\epsilon^{\mu\nu\alpha\beta} (2hp)_\alpha n_\beta \tilde{F}_{g/T}(x,\xi) \right] \quad , (3.46)$$

where I have defined $x_\pm \equiv x \pm \xi$. Here, we see explicitly the extra factor of $1/(1-\epsilon/2)$ multiplying the unpolarized gluon amplitude.

3.3 One-Loop Compton Amplitudes on Quark and Gluon Targets

In this section, we perform a one-loop calculation of the general Compton scattering on onshell quark and gluon targets in the Bjorken limit. The result will be used in the next section to show that the infrared divergences present in our amplitude are generated exclusively by the off-forward parton distributions. This property does not change in the limit of a real final state photon.

We begin with an onshell quark target of momentum p. Here, there are two diagrams at leading order (LO) and eight at next-to-leading order (NLO). The renormalization constants necessary for this process cancel themselves out in exactly the same manner we saw in Section 2.2. Half of the diagrams are shown in Fig. 3.1. The other half will be taken into account by using the crossing symmetry, i.e., the simultaneous replacement of $q \to -q$ and $\mu \leftrightarrow \nu$. The terms $\pm(x_B \to -x_B)$ in the following formulae reflect this contribution. Because of time reversal invariance, the Compton amplitude is also an even function of ξ. This relates the left and right vertex diagrams (Figs. 3.1c and 3.1d, respectively) to each other. The quark self-energy diagram and the box diagram are themselves ξ-symmetric. This symmetry not only allows us to reduce the number of graphs at NLO from four to three, but also becomes a powerful tool which helps us compute each amplitude.




Figure 3.1: Diagrams for Compton scattering on an onshell quark to order α_s.




Figure 3.2: The momentum flow in the box diagram

Examining our graphs, we see that the self-energy diagram contains a loop integral with two Feynman denominators, the vertex diagram contains one with three, and the box with four. A one-loop integral with two propagators is straightforward. Difficulties arise, however, with the calculation of three- and especially four-propagator integrals. These difficulties may be avoided in this calculation because of several simplifications. Consider first the box diagram. The loop integral is of the form (with momentum routing as shown in Fig. 3.2)

$$\int \frac{d^d k}{(2\pi)^d} \frac{\mathrm{Tr}[\gamma^\alpha\,(\not{k} - \xi\ \not{p})\,\gamma^\mu\,(\not{k} + \not{q})\,\gamma^\nu\,(\not{k} + \xi\ \not{p})\,\gamma_\alpha \not{p}(\gamma_5)]}{(k+\xi p)^2 (k-\xi p)^2 (k-p)^2 (k+q)^2}\ , \tag{3.47}$$

where I have replaced Δ^μ by $-2\xi p^\mu$. In order to simplify the integral, we express the trace as a sum of terms which cancel one of the propagators. This can be done because both k^2 and $2p\cdot k$ can be written as linear combinations of $(k+\xi p)^2$ and $(k-\xi p)^2$, and the trace vanishes whenever k^- and $k_\perp^2$ do.

Now that we have shown that a denominator can be canceled, we have effectively reduced

the four-propagator problem to a three-propagator one. Since we have only shown that the numerator will be a linear combination of two different denominators, rather that proportional to one, the four-propagator integral will in general become two three-propagator integrals. However, we may use the ξ symmetry by writing

$$\begin{aligned} k^2 &= \frac{1}{2}\ (k+\xi p)^2 + (\xi \to -\xi)\ , \\ 2p\cdot k\ &= \frac{1}{2\xi}\,(k+\xi p)^2 + (\xi \to -\xi)\ . \end{aligned} \tag{3.48}$$

In this way, we consider only the $(k+\xi p)^2$ cancelation and let the symmetry take care of the rest. The integral we now have is exactly the same (up to numerator differences) as that arising from the right vertex correction. We will see that this basic integral is the only one we must calculate to obtain both the polarized and unpolarized amplitudes for the quark and gluon contributions (if we forget, for the moment, the simple self-energy diagram).

The integral has the form

$$\int \frac{d^dk}{(2\pi)^d}\frac{Numerator}{[(k-p)^2+i\varepsilon][(k-\xi p)^2+i\varepsilon][(k+q)^2+i\varepsilon]}\ . \tag{3.49}$$

We have found that this integral is easily done in light-cone coordinates by expanding k^μ in terms of the light cone momenta, p^μ and n^μ. As in Section 2.2, we first do the k^- integration by contour in the unphysical region of large x_B and then the transverse integrations. The k^+ integration is left until the end. Writing $k^+ = yp^+$, the value of the integral (3.49) is

$$\begin{aligned} &\frac{i}{16\pi^2}\left(\frac{Q^2}{4\pi}\right)^{-\epsilon/2}\frac{\Gamma(\epsilon/2)}{1-\xi}\left\{\left(\frac{x_B-a}{x_B}\right)^{-\epsilon/2}\right. \\ &\qquad\left.\times\int_a^{x_B} dy\left(\frac{y-a}{x_B-a}\right)^{1-\epsilon/2}\left(1-\frac{y-a}{x_B-a}\right)^{-\epsilon/2}N(a)\right\}\Bigg|_{a=\xi}^{a=1}\ , \end{aligned} \tag{3.50}$$

where $N(a)$ is defined by

$$\begin{aligned} Numerator\Big|_{k^-=k_\perp^2/2p^+(y-a)} &= \alpha+\beta k_\perp^2\ \ ; \\ N(a) &= \beta - \frac{\alpha}{Q^2}\frac{x_B(x_B-a)}{(y-a)(x_B-y)}\ \ . \end{aligned} \tag{3.51}$$

Doing the y-integrals requires some care because a delicate cancelation must occur if one is to get a finite result, but the treatment is straightforward. After the ξ and crossing symmetries



Figure 3.3: Diagrams for gluon Compton scattering at one-loop.

are used, we find the full NLO result for the symmetric quark amplitude[21]

$$T_q^{(ij)} = -g^{ij}\sum_{q'} e_{q'}^2\delta_{qq'}\left\{\frac{1}{x_B-1} - \frac{\alpha_s C_F}{4\pi}\left(\frac{Q^2e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left\{\frac{3}{x_B-1}\left(\frac{2}{\epsilon}+3\right)\right.\right.$$
$$-\frac{1}{\xi}\left[\left(\frac{2\xi}{x_B^2-1}+\frac{x_B+\xi}{1-\xi^2}\right)\left(\frac{4}{\epsilon}+3-\log\left(1-\frac{\xi}{x_B}\right)\right)-3\,\frac{x_B-\xi}{1-\xi^2}\right]\log\left(1-\frac{\xi}{x_B}\right)$$
$$+\left[\left(\frac{x_B+1}{1-\xi^2}+\frac{2}{x_B-1}\right)\left(\frac{4}{\epsilon}+3-\log\left(1-\frac{1}{x_B}\right)\right)-3\,\frac{x_B-1}{1-\xi^2}\right. \tag{3.52}$$
$$\left.\left.\left.-\frac{3}{x_B-1}\right]\log\left(1-\frac{1}{x_B}\right)\right\}+(x_B\to -x_B)\right\}$$

and the antisymmetric amplitude

$$T_q^{[ij]} = -i\epsilon^{\alpha\beta ij}n_\alpha(2hp)_\beta\sum_{q'} e_{q'}^2\delta_{qq'}\left\{\frac{1}{x_B-1} - \frac{\alpha_s C_F}{4\pi}\left(\frac{Q^2e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left\{\frac{3}{x_B-1}\left(\frac{2}{\epsilon}+3\right)\right.\right.$$
$$-\left[\left(\frac{2x_B}{x_B^2-1}+\frac{x_B+\xi}{1-\xi^2}\right)\left(\frac{4}{\epsilon}+3-\log\left(1-\frac{\xi}{x_B}\right)\right)+7\frac{x_B-\xi}{1-\xi^2}\right]\log\left(1-\frac{\xi}{x_B}\right)$$
$$+\left[\left(\frac{x_B+1}{1-\xi^2}+\frac{2}{x_B-1}\right)\left(\frac{4}{\epsilon}+3-\log\left(1-\frac{1}{x_B}\right)\right)+7\,\frac{x_B-1}{1-\xi^2}\right. \tag{3.53}$$
$$\left.\left.\left.-\frac{3}{x_B-1}\right]\log\left(1-\frac{1}{x_B}\right)\right\}-(x_B\to -x_B)\right\}\ .$$

The divergences in these amplitudes are infrared divergences since renormalization has removed all ultraviolet singularities. Their presence signals the existence of nonperturbative physics in the process. As mentioned earlier, they will be factorized into nonperturbative matrix elements whose values can be extracted from experiment. We will see this explicitly in the next section. For now, we summarize the contributions from the gluon sector.

The six graphs that contribute to the LO amplitude for gluon-photon scattering can be reduced to three by reversing the fermion number flow, and one of these can be eliminated

[21] We take $\mu^2 = Q^2$ and $\mu\nu$ in the $\perp$ directions as before.

by crossing symmetry. The two distinct graphs we are left with are shown in Fig. 3.3. The denominator of Fig. 3.3a is identical to that of the quark box. Again, the numerator is seen to vanish whenever k^2 and $2p\cdot k$ do, which allows us to cancel one of the propagators exactly as above. Fig. 3.3b is somewhat more tricky. This diagram is itself symmetric under both crossing and ξ symmetry. Labeling the momenta as shown, we see that under an integral the symmetry $q\to -q$ is equivalent to $A\leftrightarrow B$ and $C\leftrightarrow D$ and $\xi\to -\xi$ is equivalent to $B\leftrightarrow D$ and $A\leftrightarrow C$. We also note that here it is possible to represent 1 as a linear combination of Feynman denominators, which guarantees our ability to cancel one. Of course, since 1 does not depend on q or ξ, it may be represented in the symmetric form

$$1=-\frac{1}{2(1-\xi)p\cdot q}A^2+(\xi\to-\xi)+(q\to-q). \tag{3.54}$$

Now it remains only to calculate the trace in a symmetric way and substitute the result into the formulae of the quark calculation. Averaging the gluon polarization for the symmetric amplitude in d dimensions, one finds

$$\begin{aligned}T_g^{(ij)}=&-\frac{\alpha_s T_F}{2\pi}g^{ij}\left(\sum_q e_q^2\right)\left(\frac{Q^2e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\frac{1}{1-\xi^2}\\ &\times\left\{-2x_B\left[\left(\frac{x_B}{1-\xi^2}-\frac{1}{2\xi}\right)\left(\frac{4}{\epsilon}+4-\log\left(1-\frac{\xi}{x_B}\right)\right)+\frac{1}{\xi}\right]\left(1-\frac{\xi}{x_B}\right)\log\left(1-\frac{\xi}{x_B}\right)\right.\\ &+\left[\left(1+2x_B\frac{x_B-1}{1-\xi^2}\right)\left(\frac{4}{\epsilon}+4-\log\left(1-\frac{1}{x_B}\right)\right)-2\right]\log\left(1-\frac{1}{x_B}\right)\\ &\left.+(x_B\to-x_B)\right\}\ ,\end{aligned} \tag{3.55}$$

for the symmetric amplitude and

$$\begin{aligned}T_g^{[ij]}=&-i\frac{\alpha_s T_F}{2\pi}\epsilon^{\alpha\beta ij}n_\alpha(\eta p)_\beta\left(\sum_q e_q^2\right)\left(\frac{Q^2e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\frac{1}{1-\xi^2}\\ &\times\left\{-2x_B\left[\frac{1}{1-\xi^2}\left(\frac{4}{\epsilon}+4-\log\left(1-\frac{\xi}{x_B}\right)\right)\right]\left(1-\frac{\xi}{x_B}\right)\log\left(1-\frac{\xi}{x_B}\right)\right.\\ &+\left[\left(1+2\frac{x_B-1}{1-\xi^2}\right)\left(\frac{4}{\epsilon}+4-\log\left(1-\frac{1}{x_B}\right)\right)-2\right]\log\left(1-\frac{1}{x_B}\right)\\ &\left.-(x_B\to-x_B)\right\}\ ,\end{aligned} \tag{3.56}$$

where $\eta=\pm1$ is the helicity of the gluon 'target', for the antisymmetric amplitude.

3.4 One-Loop Factorization in DVCS

We now turn to the infrared divergences present in all of the amplitudes. These divergences arise from regions of the loop-momentum integration where some of the internal propagators are near their 'mass shells'. In these regions, perturbative calculations are clearly meaningless. The standard procedure of fixing this problem is to factorize the amplitudes into their infrared safe (i.e. devoid of infrared divergences) and infrared divergent pieces, interpreting the latter as nonperturbative QCD quantities. The goal of this section is to show that all infrared divergences present in the Compton amplitudes can be associated with the off-forward parton distributions.

Since we consider factorization within the framework of perturbation theory, we need to compute the parton distributions in quark and gluon targets in perturbative QCD. At the leading order in α_s, one has

$$F^{(0)}_{q'/q}(x,\xi) = \delta_{qq'}\delta(x-1)\ . \tag{3.57}$$

At next-to-leading order,

$$F^{(1)}_{q'/q}(x,\xi) = \frac{\alpha_s(\mu^2)}{2\pi}\delta_{qq'}\left(-\frac{2}{\epsilon} + \log(\mu^2 e^{\gamma_E}/4\pi\mu_0^2)\right)\left(P_{qq}(x,\xi) + A_q(x,\xi)\delta(x-1)\right)\quad , \tag{3.58}$$

where

$$A_q(x,\xi) = C_F\left[\frac{3}{2} + \int_\xi^x \frac{dy}{y-x-i\epsilon} + \int_{-\xi}^x \frac{dy}{y-x-i\epsilon}\right] \tag{3.59}$$

can be determined from conditions analogous to (2.130) and

$$\begin{aligned} P_{qq}(x,\xi) &= C_F\frac{x^2+1-2\xi^2}{(1-x+i\epsilon)(1-\xi^2)}\ , && x>\xi \\ &= C_F\frac{x+\xi}{2\xi(1+\xi)}\left(1+\frac{2\xi}{1-x+i\epsilon}\right)\ , && -\xi<x<\xi \\ &= 0\ , && x<-\xi \end{aligned} \tag{3.60}$$

is calculated in the same way as $P_{q\to q}(x)$ in Section 2.3. The quark distributions contain infrared divergences signaled by the presence of the $1/\epsilon$ terms. These divergences reflect the soft physics intrinsic to the parton distributions. $F^{(1)}_{q'/q}$, calculated for the quark target, satisfies the evolution equation derived in Ref. [68] :

$$\mu^2\frac{D_a\, F_{a/T}(x,\xi,\mu^2)}{D\,\mu^2} = \frac{\alpha_s(\mu^2)}{2\pi}\int_x^1\frac{dy}{y}P_{ab}\left(\frac{x}{y},\frac{\xi}{y}\right)F_{b/T}(y,\xi,\mu^2), \tag{3.61}$$

where b is summed over all parton species and the P's are the off-forward Alterelli-Parisi kernels, or splitting functions. The 'covariant' derivative is defined to include A_a, the endpoint contribution in Eq.(3.58). Note that these kernels reduce to the standard forward splitting function (2.142) in the DIS limit $\xi \to 0$.

We can re-express the symmetric part of the quark Compton amplitude in terms of the unpolarized, off-forward quark distribution[22]

$$\begin{aligned} T_q^{(ij)} = & \\ & -g^{ij} \sum_{q'} e_{q'}^2 \int dx F_{q'/q}(x,\xi) \left\{ \frac{1}{x_B - x} - \frac{\alpha_s(Q^2) C_F}{4\pi} \left\{ \frac{9}{x_B - x} \right.\right. \\ & \quad -\frac{x}{\xi} \left[\left(\frac{2\xi}{x_B^2 - x^2} + \frac{x_B + \xi}{x_+ x_-} \right) \left(3 - \log\left(1 - \frac{\xi}{x_B} \right) \right) - 3 \frac{x_B - \xi}{x_+ x_-} \right] \log\left(1 - \frac{\xi}{x_B} \right) \\ & \quad + \left[\left(\frac{x_B + x}{x_+ x_-} + \frac{2}{x_B - x} \right) \left(3 - \log\left(1 - \frac{x}{x_B} \right) \right) \right. \\ & \quad\quad \left. -3 \frac{x_B - x}{x_+ x_-} - \frac{3}{x_B - x} \right] \log\left(1 - \frac{x}{x_B} \right) \Big\} \\ & \quad\quad \left. + (x_B \to -x_B) \right\} \ , \end{aligned} \tag{3.62}$$

and find that the infrared physics is completely contained within $F_{q'/q}(x,\xi)$. Analogously, we find that the antisymmetric part of the quark Compton amplitude can be expressed in terms of the polarized off-forward quark distribution

$$\begin{aligned} T_q^{[ij]} = & \\ & -i\epsilon^{\alpha\beta ij} n_\alpha (2hp)_\beta \sum_{q'} e_{q'}^2 \int dx \tilde{F}_{q'/q}(x,\xi) \left\{ \frac{1}{x_B - x} - \frac{\alpha_s(Q^2) C_F}{4\pi} \left\{ \frac{9}{x_B - x} \right.\right. \\ & \quad - \left[\left(\frac{2x_B}{x_B^2 - x^2} + \frac{x_B + \xi}{x_+ x_-} \right) \left(3 - \log\left(1 - \frac{\xi}{x_B} \right) \right) + 7 \frac{x_B - \xi}{x_+ x_-} \right] \log\left(1 - \frac{\xi}{x_B} \right) \\ & \quad + \left[\left(\frac{x_B + x}{x_+ x_-} + \frac{2}{x_B - x} \right) \left(3 - \log\left(1 - \frac{x}{x_B} \right) \right) \right. \\ & \quad\quad \left. +7 \frac{x_B - x}{x_+ x_-} - \frac{3}{x_B - x} \right] \log\left(1 - \frac{x}{x_B} \right) \Big\} \\ & \quad\quad \left. - (x_B \to -x_B) \right\} \ . \end{aligned} \tag{3.63}$$

We now turn to Compton scattering on a gluon target. Infrared divergent contributions come from regions where the quarks in the box diagrams are nearly onshell. Therefore, it is

[22] For convenience, we have identified the factorization scale μ with Q.

natural to associate these divergences with the quark distributions in a gluon target. At order α_s, off-forward gluonic matrix elements of the quark distribution operators have the value

$$F^{(1)}_{q/g}(x,\xi) = \frac{\alpha_s(\mu)}{2\pi}\left(-\frac{2}{\epsilon} + \log(\mu^2 e^{\gamma_E}/4\pi\mu_0^2)\right) P_{qg}(x,\xi) \ , \tag{3.64}$$

where for $x > \xi$

$$P_{qg}(x,\xi) = 2T_F \frac{x^2 + (1-x)^2 - \xi^2}{(1-\xi^2)^2} \ , \tag{3.65}$$

and for $-\xi < x < \xi$

$$P_{qg}(x,\xi) = T_F \frac{(x+\xi)(1-2x+\xi)}{\xi(1+\xi)(1-\xi^2)} \ . \tag{3.66}$$

$F^{(1)}_{q/g}(x,\xi) = 0$ for $x < -\xi$[23]

On the other hand, the finite contributions come from regions where large momenta run through the quark loop. In these regions, the photon has an effective pointlike coupling with the gluons in the target. At leading order, the off-forward gluon distribution in a gluon target is just

$$F^{(0)}_{g/g}(x,\xi) = \frac{1}{2}(\delta(x-1) - \delta(x+1)) \ . \tag{3.67}$$

Using the above off-forward distributions, we re-express the symmetric part of the gluon Compton scattering amplitude

$$\begin{aligned} T_g^{(ij)} = & \\ & -\frac{\alpha_s(Q^2)T_F}{2\pi} g^{ij}\left(\sum_q e_q^2\right)\int dx \frac{x}{x_+ x_-} F_{g/g}(x,\xi) \\ & \times\left\{\left[\left(1 + 2x_B\frac{x_B - x}{x_+ x_-}\right)\left(4 - \log\left(1 - \frac{x}{x_B}\right)\right) - 2\right]\log\left(1 - \frac{x}{x_B}\right)\right. \\ & \left. - 2x_B\left[\left(\frac{x_B}{x_+ x_-} - \frac{1}{2\xi}\right)\left(4 - \log\left(1 - \frac{\xi}{x_B}\right)\right) + \frac{1}{\xi}\right]\left(1 - \frac{\xi}{x_B}\right)\log\left(1 - \frac{\xi}{x_B}\right)\right\} \\ & -g^{ij}\sum_q e_q^2 \int dx F_{q/g}(x,\xi)\frac{1}{x_B - x} \\ & + (x_B \to -x_B) \ . \end{aligned} \tag{3.68}$$

Similarly, we can re-express the antisymmetric part of the gluon Compton amplitude in terms of helicity-dependent, off-forward quark and gluon distributions

$$T_g^{[ij]} =$$

[23]This accounts for the factor of 2 in Eq.(3.65). The antiquark contribution is treated with the quarks for simplicity.

$$
\begin{aligned}
&-i\frac{\alpha_s(Q^2)T_F}{2\pi}\epsilon^{\alpha\beta ij}n_\alpha(\lambda p)_\beta\left(\sum_q e_q^2\right)\int_{-1}^1 dx\frac{x}{x_+x_-}\tilde{F}_{g/g}(x,\xi)\\
&\quad\times\left\{\left[\left(1+2x\frac{x_B-x}{x_+x_-}\right)\left(4-\log\left(1-\frac{x}{x_B}\right)\right)-2\right]\log\left(1-\frac{x}{x_B}\right)\right.\\
&\qquad\left.-2x_B\left[\frac{x}{x_+x_-}\left(4-\log\left(1-\frac{\xi}{x_B}\right)\right)\right]\left(1-\frac{\xi}{x_B}\right)\log\left(1-\frac{\xi}{x_B}\right)\right\}\\
&\qquad -i\epsilon^{\alpha\beta ij}n_\alpha p_\beta\sum_q e_q^2\int_{-1}^1 dx\tilde{F}_{q/g}(x,\xi)\frac{1}{x_B-x}\\
&\qquad -(x_B\to -x_B)\ .
\end{aligned}
\tag{3.69}
$$

The complete set of off-forward splitting functions at leading order can be found in Ref. [68].

The one-loop Compton amplitude on a target T may be summarized by the factorization formula[24]

$$
\begin{aligned}
T_T^{ij} &= -g^{ij}\int_{-1}^1\frac{dx}{x}\left[\sum_q F_{q/T}(x,\xi)C_q\left(\frac{x}{x_B},\frac{\xi}{x_B}\right)+F_{g/T}(x,\xi)C_g\left(\frac{x}{x_B},\frac{\xi}{x_B}\right)\right]\\
&\quad -i\epsilon^{ij\alpha\beta}n_\alpha(2hp)_\beta\int_{-1}^1\frac{dx}{x}\left[\sum_q\tilde{F}_{q/T}(x,\xi)\tilde{C}_q\left(\frac{x}{x_B},\frac{\xi}{x_B}\right)+\tilde{F}_{g/T}(x,\xi)\tilde{C}_g\left(\frac{x}{x_B},\frac{\xi}{x_B}\right)\right]\ ,
\end{aligned}
\tag{3.70}
$$

where C_q, C_g, $\tilde{C}_q$ and $\tilde{C}_g$ are shown to order α_s in Eqs. (3.62, 3.68, 3.63, 3.69), respectively.[25] In the above form, all infrared sensitive contributions have been isolated in the relevant parton distributions, which must be calculated nonperturbatively or measured in experiments. In the DVCS limit $\xi\to x_B$, the coefficient functions remain finite, although they have branch cuts there. This indicates that factorization holds for two-photon amplitudes even when one of the photons is onshell. We will argue in the next section that the above formula, one of the main results of this thesis, remains valid to all orders in perturbation theory.

3.5 All Order Factorization of DVCS

In this section, we generalize the one-loop result of the previous section, showing that the factorization formula Eq.(3.70) is valid in the DVCS limit to all orders in perturbation theory.

[24]This form is correct for spin-1/2 targets. For spin-1, the replacement $2h\to\lambda$ is required. This convention is fixed by the normalization of the distributions.

[25]I emphasize here that the contributions from longitudinally polarized photons and from photon helicity flip have been neglected. Both effects start at order α_s, although longitudinal contributions are kinematically suppressed.

The one-loop result indicates that all soft divergences - those associated with integration regions where all components of some internal momenta are zero - cancel, whereas all collinear divergences can be factorized into the off-forward parton distributions. To see that this happens also at higher orders in perturbation theory, it is important to understand how the soft cancelation happens in the simplest case.

The self-energy diagram in Fig. 3.1b does not contain any infrared divergences because the intermediate quark is far offshell. The vertex corrections in Figs. 3.1c and 3.1d potentially have infrared divergences, but a simple power counting argument indicates that these diagrams are in fact infrared convergent. Thus infrared divergences appear only in the box diagram. In the region where the gluon momentum k is soft, we can approximate the integral as

$$\sim \int \frac{d^4k}{(2\pi)^4} p' \cdot p \, \frac{1}{p' \cdot k} \, \frac{1}{p \cdot k} \, \frac{1}{k^2} \ , \tag{3.71}$$

where p and p' are the momenta of the two external quark lines. On the other hand, the wavefunction renormalization of an 'onshell' quark, δZ, also contains infrared divergences. Grouping these divergent terms together, we have the entire soft contribution

$$\sim -\frac{1}{2} \int \frac{d^4k}{(2\pi)^4} \left(\frac{p^\mu}{p \cdot k} - \frac{p'^\mu}{p' \cdot k} \right)^2 \frac{1}{k^2} \ . \tag{3.72}$$

In the collinear approximation, p' is proportional to p and thus the above integral vanishes.

For higher-order Feynman diagrams, a systematic method of identifying, regrouping and factorizing infrared-sensitive contributions has been developed by Sterman, Libby, Collins, and others [47, 48, 49, 55]. As detailed in Section 2.4, this method essentially consists of the following steps: 1) simplify the Feynman integrals by setting all the soft scales to zero, including the quark masses; 2) identify the regions of loop integration which give rise to infrared divergences; 3) use infrared power counting to find the leading infrared-divergent regions; 4) show that all soft and collinear divergences either cancel or factorize into some nonperturbative quantities. In the remainder of this section, we examine the validity of the factorization formula Eq.(3.70) in the limit of $x_B = \xi$ following the above steps.

In DVCS, the leading contributions come essentially from a massless collinear process in which the external momenta take the form shown in Eq.(3.32). In this simplified kinematic region, all infrared-sensitive contributions appear as $1/\epsilon$ poles in dimensional regularization. If a contribution contains no infrared divergences, it comes from regions of loop momenta




Figure 3.4: General reduced diagrams for the DVCS process.

comparable to the hard scale Q^2, and thus is insensitive to the soft scales. An infrared-divergent contribution must come from the integration regions where some internal propagators are near their mass shells. The parts of these regions which will actually generate infrared divergences can be identified from the Landau equations, as we have seen, and characterized by reduced diagrams. We will argue below that the general *leading* reduced diagram for DVCS is the one shown in Fig. 3.4a. Here, an incoming virtual photon and an outgoing real one are attached to the hard interaction blob, which in turn is connected to the forward nucleon jet with two collinear quark lines or two physically polarized gluon lines. An arbitrary number of longitudinally polarized collinear gluon lines can also connect the hard diagram to the soft one. This diagram is identical to that of DIS. Indeed, the only real difference between DVCS and DIS lies in the fact that the final state photon is real.

A real photon can have a pointlike coupling to quarks as well as an extended coupling via its soft wavefunction. In this case, one has an additional reduced diagram in which a jet of quarks and gluons emerges in the n^μ direction and combines into a real photon long after the hard scattering (Fig. 3.4b). Such a reduced diagram has already been considered in Ref. [56] and is $\mathcal{O}(Q^{-1})$ by infrared power counting. Indeed, according to the discussion in Section 2.4, the photon wavefunction vertex has a soft mass dimension $1 = 2$ (quark lines) -1 (photon state). A negative hard power (Q^{-1}) is needed in $T^{\mu\nu}$ to balance it out. Hence the soft coupling of quarks to photons will appear only at next-to-leading twist.

The only remaining difficulties lie in the soft divergences shown in Figure 3.4c. However, since we have already shown that real-photon effects are sub-leading, one can use the same arguments here as for DIS. The differences appear only at higher orders in our twist expansion.

At this point, the only leading reduced diagram is identical to that in DIS. The arguments presented in Section 2.4 may also be applied here in almost exactly the same form.[26] The eikonal line together with the physical quarks and gluons and the nucleon jets form the off-forward parton distributions defined in Section 3.2. In the hard scattering, only the total momentum and charge supplied by collinear partons are important. Thus, one can calculate it with incoming physical partons carrying the total momentum and charge of all the collinear longitudinally-polarized gluons. In this way, we have a complete factorization of the soft and hard physics in the DVCS process.

3.6 Generalized Operator Product Expansion and Wilson's Coefficients to the NLO Order

As we saw in Section 2.5, the calculation of the Compton amplitude can be exploited to derive an expansion for the product of two electromagnetic current operators in terms of local QCD operators. Since the kinematics of that chapter require identical initial and final states, the OPE we derived there is valid only for diagonal matrix elements. Here, however, we have considered a somewhat more general case in which certain off-diagonal effects have been included. It stands to reason that we can use these results to derive a *generalized* OPE which is valid for these special off-forward matrix elements.

In our operator derivation of the leading OPE, Equation (2.187), we ignored several 'total derivative' operators which cannot contribute to forward matrix elements. By treating the initial and final states symmetrically, systematically using $\overleftrightarrow{\mathcal{D}}$ in the place of $\mathcal{D}$, we will see that these total derivative operators still do not contribute to the leading expansion. However, at next-to-leading order, they become indispensable as generators of the ξ-dependence of our amplitude. In the remainder of this section, we will recast our factorization formula in its generalized OPE form. In the process, we identify these total derivative operators and obtain their Wilson coefficients to next-to-leading order in α_s. The final result agrees with the known OPE, Eq.(2.208), when diagonal matrix elements are considered.

To derive the generalized OPE, we expand T^{ij} in a power series about $x_B = \infty$. In this

[26] The only difference is in the off-forward nature of the distributions relevant to this case.

way, we can express the amplitude in terms of moments of the parton distributions rather than the distributions themselves. Eventually, we will relate the moments of parton distributions to the matrix elements of local operators. After the aforementioned expansion, we have

$$
\begin{aligned}
T_T^{ij} &= -g^{ij} T_S - i\epsilon^{\alpha\beta ij} n_\alpha (2hp)_\beta T_A \ ; \\
T_S &= 2 {\sum_{n=2}^{\infty}}' {\sum_{m=0}^{\infty}}' \int \frac{dx}{x} \left(\frac{x}{x_B}\right)^n \left(\frac{\xi}{x_B}\right)^m \sum_{a=q,g} c_{nm}^a F_{a/T}(x,\xi) \ , \qquad (3.73) \\
T_A &= 2 {\sum_{n=1}^{\infty}}' {\sum_{m=0}^{\infty}}' \int \frac{dx}{x} \left(\frac{x}{x_B}\right)^n \left(\frac{\xi}{x_B}\right)^m \sum_{a=q,g} \tilde{c}_{nm}^a \tilde{F}_{a/T}(x,\xi) \ . \qquad (3.74)
\end{aligned}
$$

The coefficients for the moments of the quark distributions in the expansion are

$$
\begin{aligned}
c_{nm}^q &= \delta_{m0} - \frac{\alpha_s C_F}{4\pi} \left\{ \left[9 - \frac{8}{n} + \frac{2}{n+1} + 4S_2(n-1) - 4T_1^1(n-1) \right.\right. \\
&\quad \left. - S_1(n-1)\left(3 + \frac{2}{n} + \frac{2}{n+1}\right)\right] \delta_{m0} \\
&\quad + \left[\frac{6n}{m(m+n)} - \frac{2}{(n+m)(n+m+1)} + \frac{4}{m} S_1(m-1) \right. \qquad (3.75) \\
&\quad \left.\left. - 2S_1(n+m-1)\left(\frac{1}{n+m} + \frac{1}{n+m+1}\right)\right](1-\delta_{m0})\right\} \ , \\
\tilde{c}_{nm}^q &= \delta_{m0} - \frac{\alpha_s C_F}{4\pi} \left\{ \left[9 - \frac{8}{n+1} + \frac{2}{n} + 4S_2(n-1) - 4T_1^1(n-1) \right.\right. \\
&\quad \left. - S_1(n-1)\left(3 + \frac{2}{n} + \frac{2}{n+1}\right)\right] \delta_{m0} \\
&\quad + \left[\frac{6n}{m(n+m)} + \frac{8}{(n+m)(n+m+1)} + \frac{4}{m} S_1(m-1) \right. \qquad (3.76) \\
&\quad \left.\left. - 2S_1(n+m-1)\left(\frac{1}{n+m} + \frac{1}{n+m+1}\right)\right](1-\delta_{m0})\right\} \ ,
\end{aligned}
$$

where I have introduced

$$
\begin{aligned}
S_j(n) &\equiv \sum_{i=1}^{n} \frac{1}{i^j} \ , \\
T_j^k(n) &\equiv \sum_{i=1}^{n} \frac{S_j(i)}{i^k} \ , \qquad (3.77)
\end{aligned}
$$

as in Section 2.5. The above expansion contains only positive powers of x and ξ. This result is not immediately obvious because of the x_+x_- denominators in the amplitudes. In the case of the gluon distribution functions, we have an additional factor of x_+x_- in the denominator. Since the final OPE contains only local operators, these factors have to be canceled in the process of expansion. This is indeed the case; the coefficients of the positive moments of the

gluon distributions are

$$\begin{aligned} c^g_{nm} &= \frac{\alpha_s T_F}{2\pi}\left[\frac{m}{n+m}-\frac{m+2}{n+m+2}\right. \\ &\qquad \left. -2S_1(n+m-1)\left(\frac{1}{n+m}-\frac{m+2}{n+m+1}+\frac{m+2}{n+m+2}\right)\right] \ , \\ \tilde{c}^g_{nm} &= \frac{\alpha_s T_F}{2\pi}\left[2+2S_1(n+m-1)\right]\left(\frac{m+1}{n+m}-\frac{m+2}{n+m+1}\right) \ . \end{aligned} \tag{3.78}$$

Having obtained an expansion involving the moments of the distributions, we move toward a general form of the OPE. To this end, we consider the moments of the parton distributions. As before, these moments can be made into local operators via an appropriate use of integration by parts. The result is[27]

$$\int_{-1}^{1} dx x^{n-1} F_q(x,\xi) = \frac{1}{2}\left\langle P_f \left| \overline{\psi}_q(0)\, i\overleftrightarrow{\mathcal{D}}_{(\mu_1}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,\gamma_{\mu_n)}\psi_q(0)\right| P_i\right\rangle n^{\mu_1}\cdots n^{\mu_n} \ . \tag{3.79}$$

Defining

$${}_q\mathcal{O}^n_{\mu_1\mu_2\cdots\mu_n} = \overline{\psi}_q(0)\, i\overleftrightarrow{\mathcal{D}}_{(\mu_1}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,\gamma_{\mu_n)}\psi_q(0), \tag{3.80}$$

where $(\cdots)$ signifies that the indices are symmetrized and the trace has been removed, we see that the replacement

$$(n\cdot i\partial)\ {}_q\mathcal{O}^{+\cdots+}_n \to 2\xi\mathcal{O}^{+\cdots+}_n \tag{3.81}$$

is valid for the matrix elements we consider. This prompts us to define

$$\begin{aligned} {}_q\mathcal{O}^{n,m}_{\mu_1\mu_2\cdots\mu_n} &= i\partial_{(\mu_1}\cdots i\partial_{\mu_m}\overline{\psi}(0)\, i\overleftrightarrow{\mathcal{D}}_{\mu_{m+1}}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,\gamma_{\mu_n)}\psi(0) \ , \\ {}_q\tilde{\mathcal{O}}^{n,m}_{\mu_1\mu_2\cdots\mu_n} &= i\partial_{(\mu_1}\cdots i\partial_{\mu_m}\overline{\psi}(0)\, i\overleftrightarrow{\mathcal{D}}_{\mu_{m+1}}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,\gamma_{\mu_n)}\gamma_5\psi(0) \ , \\ {}_g\mathcal{O}^{n,m}_{\mu_1\mu_2\cdots\mu_n} &= i\partial_{(\mu_1}\cdots i\partial_{\mu_m}F_{\mu_{m+1}\alpha}(0)\, i\overleftrightarrow{\mathcal{D}}_{\mu_{m+2}}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,F^{\alpha}_{\mu_n)}(0) \ , \\ {}_g\tilde{\mathcal{O}}^{n,m}_{\mu_1\mu_2\cdots\mu_n} &= i\partial_{(\mu_1}\cdots i\partial_{\mu_m}F_{\mu_{m+1}\alpha}(0)\, i\overleftrightarrow{\mathcal{D}}_{\mu_{m+2}}\cdots i\overleftrightarrow{\mathcal{D}}_{\mu_{n-1}}\,i\tilde{F}^{\alpha}_{\mu_n)}(0) \ . \end{aligned} \tag{3.82}$$

After replacing the moments of parton distributions in Eq.(3.74) with matrix elements of these operators and interpreting the result as an operator relation, we find the following generalized OPE,

$$i\int d^4z\ e^{iq\cdot z}\ TJ^{\mu}\left(\frac{z}{2}\right)J^{\nu}\left(-\frac{z}{2}\right)$$

[27] As mentioned above, the off-diagonal nature of these matrix elements complicates this process slightly. The derivatives that actually appear are $\overleftrightarrow{\mathcal{D}} \equiv 1/2(\overrightarrow{\mathcal{D}}-\overleftarrow{\mathcal{D}})$, where the arrow indicates the direction the derivative is meant to operate. In DIS, this distinction is unnecessary since $\overleftrightarrow{\mathcal{D}} = \overrightarrow{\mathcal{D}}$ in this case.

$$= \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{q^2}\right) \sum_{n=2}^{\infty}{}' \sum_{m=0}^{n}{}' \left(\frac{2^{n-m} q_{\mu_1} \cdots q_{\mu_n}}{(Q^2)^n}\right) \sum_{a=q,g} c^a_{n-m,m}\ {}_a\mathcal{O}^{\mu_1 \cdots \mu_n}_{n,m}$$
$$-i\epsilon^{\mu\nu\alpha\beta} q_\alpha\, g_{\beta\lambda} \sum_{n=1}^{\infty}{}' \sum_{m=0}^{n}{}' \left(\frac{2^{n-m} q_{\mu_2} \cdots q_{\mu_n}}{(Q^2)^n}\right) \sum_{a=q,g} \tilde{c}^a_{n-m,m}\ {}_a\tilde{\mathcal{O}}^{\beta\mu_2 \cdots \mu_n}_{n,m} \quad , \qquad (3.83)$$

where the primed summations once again imply even or odd values only, depending on the initial value. It must be pointed out that the generalized OPE does not have a unique form. One can define x_B as any nontrivial dimensionless invariant formed from the external momenta which remains finite in the Bjorken limit and expand the amplitude in inverse powers of this variable. This will lead to a different set of coefficient functions, but the physical content is the same. The choice of which OPE to use is determined by the specifics of the problem at hand.

Since the operators $\mathcal{O}_{n,m}$ are symmetrized and traceless, their rank is n. The mass dimension of these operators is just the dimension of the fields plus that of the derivatives, or $3 + n - 1 = 4 + n - 2$. Hence these operators are all twist 2. As we saw in Section 2.5, the small number of possible twist-2 operators in QCD led to autonomous evolution of the diagonal matrix elements of our operators in the nonsinglet sector. Here, we see that for a given spin n there are in fact n independent nonsinglet twist-2 operators with the same quantum numbers that contribute to our generalized OPE. Since they are not constrained by symmetry to evolve autonomously, the off-diagonal matrix elements of these operators will mix as the scale is varied. It is this behavior that leads to the ξ dependence of the off-forward splitting kernels above.

Although Lorentz invariance cannot distinguish between these operators, the larger symmetry group of conformal invariance can. It is the irreducible representations of *this* symmetry group, which are definite combinations of our operators $\mathcal{O}_{n,m}$, that diagonalize the leading order anomalous dimension matrix. Using this larger symmetry group, one can determine *all* of the above generalized Wilson coefficients *from the results of Section 2.5* [73]. Unfortunately, conformal invariance is broken by a quantum anomaly[28] beyond this order and can no longer constrain the form of the OPE.

[28] Conformal invariance is based on the fact that the (massless) QCD lagrangian contains no scale. However, dimensional transmutation generates a scale (μ^2) on which parameters can depend. Stated succinctly, the conformal invariance of QCD is broken by the β-function.

The above expression contains only the contributions at leading order in $1/Q^2$. At the next order in $1/Q$ one has to consider operators of higher twist. Some of these operators are discussed in the next chapter.

3.7 Summary of DVCS

In this chapter, we have studied the QCD factorization for deeply virtual Compton scattering explicitly at one loop and then to all orders in perturbation theory. Our conclusion is that DVCS is factorizable in perturbation theory at leading twist. This statement has the same level of rigor as that for DIS studied in Chapter 2. The OFPD's probed in this process are not only interesting themselves, but also can be manipulated to yield information on the total spin carried by quarks in the proton. In conjunction with DIS data, we can use this to obtain the quark orbital contribution to proton spin. This quantity has tremendous implications on the nature of the proton wavefunction and is not readily obtainable from any other known process. This makes DVCS a prime candidate for experimental study in the next few years. Indeed, several experiments are already proposed at Jefferson Lab.

Expanding our one-loop result for general off-forward Compton scattering about $x_B = \infty$, we were able to obtain a generalization of Wilson's OPE. Our expression is valid at leading twist for off-diagonal as well as diagonal matrix elements. This generalization requires the introduction of total derivative operators whose forward matrix elements vanish. Our one-loop result gives the coefficients of these operators to NLO in QCD. A full generalization would also require an analysis of longitudinal photon scattering, which has been done in Ref. [74], and the photon-helicity flip amplitude [76].

The scale evolution of total derivative operators can best be studied using conformally-symmetric operators [73]. In fact, it has been known for a long time that at the leading-log level, operators of the same twist and dimension evolve multiplicatively in Gegenbauer polynomial combinations. It is a simple exercise to transform Eq.(3.83) into this basis. This fact is a consequence of conformal symmetry in massless QCD. Although this symmetry is broken by a quantum anomaly [80], it still has implications on the behavior of the OPE. One can actually obtain the one-loop generalized OPE coefficients from the OPE introduced in Section 2.5 by exploiting the constraints from conformal symmetry. Unfortunately, the quantum anomaly

causes this method to become uneconomical for higher-order calculations.

Measurements of the DVCS amplitude are made difficult by the fact that it competes with the simple bremsstrahlung process in which the electron deposits the full momentum transfer Δ with one virtual photon, emitting the final state photon on its own. Unfortunately, due to the propagator of the virtual photon in this process, it wins the competition in the region of small $-t$ relevant to spin physics. However, this competition is a mixed blessing since the interference between the two amplitudes allows a measurement of the phase associated with DVCS. Furthermore, in regions where the direct emission dominates, the interference actually *enhances* the effect. The Bethe-Heitler amplitude for direct emission is quite well-known. Since the only hadronic physics that enters it are standard electromagnetic form factors, there should be no difficulty in performing a direct subtraction to extract the DVCS contribution. Some estimates of the cross-section in several different kinematical regions is presented in Ref. [68].

Chapter 4

The Structure Function $f_T(x, \mu^2)$ - Higher Twist Evolution

Until this point, we have been studying only the leading contributions to scattering processes in the Bjorken limit. This led to distribution functions whose moments could be classified as twist-2. As we have seen, partons described by twist-2 distributions move collinear to the parent hadron. This is what we expect since transverse behavior is suppressed in the Bjorken limit. However, a complete understanding of hadrons requires information that is not contained in these collinear distribution functions. In particular, transverse momentum effects are intimately related to partonic correlations within hadrons.

A simple example of coherent parton scattering is the interference of a single quark with a quark *and* a gluon in a nucleon target. To describe this phenomenon, it is necessary to introduce a three-parton light-cone correlation function

$$\mathcal{M}^{\alpha}(x, y, \mu^2) = \int \frac{d\lambda}{2\pi} \frac{d\mu}{2\pi} e^{-i\lambda y} e^{-i\mu(x-y)} \langle PS|\overline{\psi}(\lambda) i D^{\alpha}(\mu n)\psi(0)|PS\rangle \ , \tag{4.1}$$

rather than the usual two-parton functions of the last two chapters. General parton correlations involve more than one Feynman variable, and hence their scale evolution is more complicated than those we have studied.

Experimental study of parton correlations is challenging for a number of reasons. One is the lack of processes in which all Feynman variables in a parton correlation can be kinematically controlled. For instance, in lepton-nucleon deep-inelastic scattering with trasverse polarization, one can measure the structure function $f_T(x, \mu^2)$. In the Bjorken limit, $f_T(x, \mu^2)$ is related

to a y-moment of the above correlation function. Since a moment of $M^\alpha(x, y, \mu^2)$ does not evolve autonomously, knowing the entire $f_T(x, \mu^2)$ at one scale is not sufficient to calculate it at another. This makes an analysis of $f_T(x, \mu^2)$ data at different scales difficult.

Several years ago, Ali, Braun, and Hiller (ABH) [36] made a remarkable discovery that in the limit of large number of colors, N_c, the twist-3 part of $f_T(x, \mu^2)$ does evolve autonomously at the one-loop level in the nonsinglet sector. Their result has since been widely used in model calculations and analyses of experimental data [81]. More recently, similar results have been found for the evolutions of other twist-3 functions, $h_L(x, \mu^2)$ and $e(x, \mu^2)$ [82]. Given the practical importance of the ABH result, a deeper understanding of the large-N_c simplification is clearly desirable. Moreover, it is interesting to investigate the possibility of a similar simplification at two or more loops and for analogous twist-4 correlations.

In this chapter, I present a direct calculation of the large-N_c evolution of $f_T(x, \mu^2)$ in the light-cone gauge. This calculation is based on original research by X. Ji and I published in [37]. Section 4.1 is devoted to a discussion of the kinematics and relevant correlations in our process. In Section 4.2, the details of the calculation are discussed. Here, we will see that the autonomy of the twist-3 evolution arises from a special property of one particular Feynman diagram. Since this property is independent of the γ-matrix structure of the composite operators inserted, the ABH result generalizes immediately to the twist-3 parts of $h_L(x, \mu^2)$ and $e(x, \mu^2)$. Unfortunately, for various reasons, it seems there is no similar large-N_c simplification for twist-4 functions nor for $f_T(x, \mu^2)$ beyond one loop.

A summary of this chapter appears in Section 4.3.

4.1 DIS With Transverse Polarization

The process we wish to consider here has the same kinematics as that of Chapter 2, but here we take the proton spin along a transverse direction (either the 1- or 2-direction). Of course, this means that we cannot take the proton mass strictly to zero because massless fermions do not have transverse spin components. However, we will ignore this quantity as small whenever it is subleading.

Since $q \cdot S = 0$, the spin-dependent part of $T^{\mu\nu}$ can be written

$$T^{[\mu\nu]} = -i\epsilon^{\mu\nu\alpha\beta} q_\alpha S_\beta \frac{2M}{\nu} \left(S_1(\nu, Q^2) + S_2(\nu, Q^2) \right) \quad . \tag{4.2}$$

Assuming a factorization theorem for this process, we can express its amplitude as a convolution of a perturbative scattering coefficient function with certain nonperturbative distributions. As before, we write the quark density matrix[1]

$$\mathcal{M}_{\alpha\beta}(x) = \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle PS| \overline{\psi}_\beta(\lambda n) \psi_\alpha(0) |PS\rangle = \frac{1}{2} f(x) (\not{P})_{\alpha\beta} + f_T(x) M (\gamma_5 \not{S})_{\alpha\beta} \quad . \tag{4.3}$$

The first term is just the spin-averaged quark distribution function of Chapter 2. Since this term is spin-independent, the distinction between transversely polarized targets and longitudinally polarized targets is meaningless to it.

In the limit $M \to 0$, the second structure reduces to $h_T \not{n} \gamma_5$ as in Section 2.3. However, its coefficient contains more than just the function we met there. The easiest way to see this is by examining its moments :

$$f_T(x) = -\frac{1}{M} S_\alpha \int \frac{d\lambda}{2\pi} e^{-i\lambda x} \langle PS| \overline{\psi}(\lambda n) \gamma^\alpha \gamma_5 \psi(0) |PS\rangle \quad , \tag{4.4}$$

$$\int dx\, x^{n-1} f_T(x) = -\frac{1}{M} S_\alpha \langle PS| \overline{\psi} \gamma^\alpha \gamma_5 \left(i\partial^+ \right)^{n-1} \psi |PS\rangle \quad . \tag{4.5}$$

The local operators associated to the moments of $f_T(x)$ are very similar to those related to $\tilde{f}(x)$. In fact, using Lorentz invariance we can write

$$\begin{aligned} \langle PS| \overline{\psi} \gamma^{(\mu_1} i\mathcal{D}^{\mu_2} \cdots i\mathcal{D}^{\mu_n)} \gamma_5 \psi |PS\rangle &= 2\tilde{a}_n M S^{(\mu_1} P^{\mu_2} \cdots P^{\mu_n)} \quad , \\ \langle PS| \overline{\psi} \gamma^+ i\mathcal{D}^+ \cdots i\mathcal{D}^+ \gamma_5 \psi |PS\rangle &= 2\tilde{a}_n M S^+ P^+ \cdots P^+ \quad . \end{aligned} \tag{4.6}$$

Here, we have symmetrized all of the indices and removed all possible traces to form a twist-2 operator. Working with the kinematics of Chapter 2, we can extract the value of $\tilde{a}_n$ in polarized DIS :

$$\tilde{a}_n = \int dx\, x^{n-1} \tilde{f}(x) \quad . \tag{4.7}$$

Hence only the twist-2 piece of $f_T(x)$ is known from DIS. The remaining twist-3 portion contains new physics. Let us see how f_T contributes to $T^{[\mu\nu]}$ at leading order.

[1] Here and for the remainder of this chapter, we work in the light-cone gauge $n \cdot \mathcal{A} = 0$.



Figure 4.1: Leading diagrams for 'naive' scattering with transverse polarization.

The relevant diagrams for quark scattering at leading order are shown in Figure 4.1. The result,

$$T^{[\mu\nu]} \sim i\epsilon^{\mu\nu\alpha\beta}(MS)_\beta \, \frac{2e_q^2}{\nu} \int dx \, f_T(x) \, \frac{1}{x - x_B} \, (xp+q)_\alpha \quad , \tag{4.8}$$

does not satisfy the QED Ward identity, $q_\mu T^{\mu\nu} = 0$, unless S and p are parallel. As usual, the failure of an amplitude to satisfy Ward identities indicates missing contributions. Here, we are being reprimanded for neglecting the contribution due to transverse quark momentum. Our quarks are massless, so their spins are aligned with their momenta. To support the transverse polarization of the nucleon, quarks must have a transverse component to their momenta.

Revisiting the diagram of Figure 4.1, we take the incoming quark to have momentum $xp+k_\perp$ and expand the amplitude about $k_\perp = 0$. The expansion of the intermediate propagator is

$$\frac{i}{x\not{p} + \not{q} + \not{k}_\perp} = \frac{i}{x\not{p} + \not{q}} + \frac{i}{x\not{p} + \not{q}} \, i\not{k}_\perp \, \frac{i}{x\not{p} + \not{q}} + \mathcal{O}(k_\perp^2/Q^2) \quad , \tag{4.9}$$

so the leading transverse momentum correction takes the form

$$ie_q^2 \int dx \int d^2\,k_\perp \, \mathrm{Tr}\left[\gamma^\mu \, \frac{i}{x\not{p} + \not{q}} \, i\not{k}_\perp \, \frac{i}{x\not{p} + \not{q}} \gamma^\nu \mathcal{M}(x, k_\perp)\right] \quad . \tag{4.10}$$

The transverse momentum distribution,

$$\begin{aligned} \mathcal{M}_{\alpha\beta}(x, k_\perp) &\equiv \sum_X \langle PS|\,\overline{\psi}_\beta\,|X\rangle\,\langle X|\,\psi_\alpha\,|PS\rangle\,\delta(n\cdot p_X + x - 1)\delta(p_{\perp X} + k_\perp) \\ &= \int \frac{d\lambda}{2\pi}\frac{d^2 z_\perp}{(2\pi)^2} e^{-i\lambda x} e^{iz_\perp\cdot k_\perp}\,\langle PS|\,\overline{\psi}_\beta(\lambda n)\psi_\alpha(z_\perp)\,|PS\rangle \quad , \end{aligned} \tag{4.11}$$

represents the probability of finding a quark with momentum $xp+k_\perp$ in a nucleon of momentum P and polarization S. Turning the factor of $k_\perp$ in our amplitude into a derivative via partial




Figure 4.2: The gluon correction required for QCD gauge-invariance in DIS with transverse polarization.

integration, we arrive at[2]

$$ie_q^2 \int dx \ \mathrm{Tr}\left[\gamma^\mu \ \frac{i}{x\not{p}+\not{q}} \ i\gamma_\alpha^\perp \ \frac{i}{x\not{p}+\not{q}}\gamma^\nu \mathcal{M}^\alpha(x)\right] \ . \tag{4.12}$$

$\mathcal{M}^\alpha(x)$ is a new distribution representing the momentum asymmetry induced by the hadron polarization.

In its present form, $\mathcal{M}^\alpha(x)$ is not gauge-invariant. Since transverse momentum cannot be separated from transversely polarized gauge fields, we must include these effects to achieve gauge invariance. This implies that the contribution represented by the diagram in Figure 4.2 is necessary. In terms of the distribution

$$\begin{aligned}
\mathcal{M}^\alpha(x,y) &\equiv \int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi} \, e^{-i\mu(x-y)} \, e^{-i\lambda y} \left\langle PS \left| \overline{\psi}(\lambda n) i\mathcal{D}^\alpha(\mu n)\psi(0)\right| PS\right\rangle \\
&= G_1(x,y) M S^\alpha \ \gamma_5 \not{p} + G_2(x,y) M i T^\alpha \ \not{p} \ ; \qquad (4.13) \\
G_1(x,y) &= -\frac{S_\sigma}{M}\int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi} \, e^{-i\mu(x-y)} \, e^{-i\lambda y} \left\langle PS \left| \overline{\psi}(\lambda n)\not{n}\gamma_5 i\mathcal{D}^\sigma(\mu n)\psi(0)\right| PS\right\rangle \qquad (4.14) \\
G_2(x,y) &= i\frac{T_\sigma}{M}\int \frac{d\lambda}{2\pi}\frac{d\mu}{2\pi} \, e^{-i\mu(x-y)} \, e^{-i\lambda y} \left\langle PS \left| \overline{\psi}(\lambda n)\not{n} i\mathcal{D}^\sigma(\mu n)\psi(0)\right| PS\right\rangle \ , \qquad (4.15)
\end{aligned}$$

where I have defined

$$T^\mu \equiv \epsilon^{\mu\nu\lambda\rho} S_\nu P_\lambda n_\rho \tag{4.16}$$

to introduce the other transverse direction, and

$$\mathcal{D}^\alpha(\mu n) \equiv \partial^\alpha + igA^\alpha(\mu n) \ , \tag{4.17}$$

our full correction can be neatly summed up as

$$ie_q^2 \int dx\, dy \ \mathrm{Tr}\left[\gamma^\mu \ \frac{i}{y\not{p}+\not{q}} \ i\gamma_\alpha^\perp \ \frac{i}{x\not{p}+\not{q}}\gamma^\nu \mathcal{M}^\alpha(x,y)\right] \ . \tag{4.18}$$

[2] The $\perp$ on $\gamma_\alpha^\perp$ is to remind us that the sum over α includes only transverse dimensions.

Using the fact that $p^2 = n^2 = 0$, this expression can be further simplified to

$$\frac{e_q^2}{4\nu}\int \frac{dx}{x-x_B}\,\mathrm{Tr}\left[\gamma^\mu \not p \gamma_\alpha^\perp \not n \gamma^\nu \int dy \mathcal{M}^\alpha(x,y) + \gamma^\mu \not n \gamma_\alpha^\perp \not p \gamma^\nu \int dy \mathcal{M}^\alpha(y,x)\right] \quad . \tag{4.19}$$

Substituting (4.13) and using the relations

$$G_1(x,y) = \quad G_1(y,x) \tag{4.20}$$

$$G_2(x,y) = -G_2(y,x) \ , \tag{4.21}$$

which follow from the reality of our distributions, leads to

$$-\,i\epsilon^{\mu\nu\alpha\beta}p_\alpha\,(MS)_\beta\,\frac{2e_q^2}{\nu}\int dx\left[\int dy\ (G_1(x,y)+G_2(x,y))\right]\ \frac{1}{x-x_B} \quad . \tag{4.22}$$

Apparently, only the first moments of $G_{1,2}(x,y)$ with respect to y enter our process. From their definitions, one can show that

$$\begin{aligned}&\int dy\,[G_1(x,y)+G_2(x,y)]\\ &\qquad = \frac{1}{M}\ \int\frac{d\lambda}{2\pi}\left\langle PS\left|\overline{\psi}(\lambda n) i\mathcal{D}_\perp^\sigma(\lambda n)\,(iT_\sigma \not n\ - S_\sigma \not n\,\gamma_5)\,\psi(0)\right|PS\right\rangle \quad .\end{aligned} \tag{4.23}$$

Since the expression in parentheses is just one of the many ways to say '$\gamma^\sigma \not n \not S \gamma_5$', the equation of motion in this gauge,

$$\not n \not p\, i\partial^+\psi = -\not n\, i\not{\mathcal{D}}_\perp \psi \tag{4.24}$$

allows us to write finally[3]

$$\int dy\,[G_1(x,y)+G_2(x,y)] = x\,f_T(x) \quad . \tag{4.25}$$

Adding (4.22) to (4.8), we see that this contribution does indeed cancel the gauge dependence of our initial result. The complete amplitude,[4]

$$T^{[\mu\nu]} = i\epsilon^{\mu\nu\alpha\beta}q_\alpha(MS)_\beta\,\frac{2e_q^2}{\nu}\ \int\,dx\,f_T(x)\,\left(\frac{1}{x-x_B}-\frac{1}{x+x_B}\right) \ , \tag{4.26}$$

is fully gauge-invariant. In fact, the only difference between transverse and longitudinal polarization appears to be in the relevant distribution function. From the point of view of the operator product expansion, this result is obvious. Since the OPE makes no reference to the

[3] In order to get rid of the structure $\not p \not n - \not n \not p$, we must assert the reality of our distributions one more time.

[4] we can't forget the crossing contribution!

polarization of the target, the coefficient of the twist-2 contribution must be the same for both processes. However, the twist-3 contributions are not constrained in this way. In fact, from the analysis above it seems like a miracle that only the simple distribution $f_T(x)$ appears in the final results rather than $G_1(x,y)$ and $G_2(x,y)$. We certainly cannot expect this behavior to persist in the next order.

The calculation above is extremely disconcerting. It seems very strange that we had to consider *gluon* scattering and transverse momentum effects to obtain *QED* gauge invariance. In a fundamental sense, it seems that this result should've been automatic. However, the Feynman diagram in Figure 4.1 implicitly assumes that we can interpret the scattering within the parton model. The trouble with a parton model interpretation of $f_T(x)$ can easily be seen by quantizing the theory on the light front. Rather than the treatment discussed in Appendix D.1, one can impose canonical commutation relations on the fields and their conjugate momenta at equal values of x^+ [83]. In this formalism, the '+' direction becomes the new 'time' variable. For massless field quanta moving in the + direction, this is a very logical choice for the time axis. In light-front quantization, the quark field contains two very different components. Defining

$$\psi_\pm \equiv \frac{1}{2}\gamma^\mp\gamma^\pm\psi \ , \tag{4.27}$$

we see that the equation of motion constrains ψ_- :

$$\psi_-(\lambda n) = -\frac{1}{2}\frac{1}{in\cdot\partial}\ \not{n} i\not{D}_\perp(\lambda n)\psi_+(\lambda n) \ . \tag{4.28}$$

Physically, this means that only ψ_+ contains freely propagating[5] quark quanta. The γ^- in the definition of ψ_- has annihilated all of the free portions of the quark field since these satisfy $\not{p}u(xp) = 0$. Looking at $f_T(x)$,

$$\begin{aligned} f_T(x) = -\frac{1}{M}S_\alpha \int \frac{d\lambda}{2\pi} e^{-i\lambda x}\Big[\quad & \langle PS|\,\overline{\psi}_-(\lambda n)\gamma^\alpha\gamma_5\psi_+(0)\,|PS\rangle \\ & + \langle PS|\,\overline{\psi}_+(\lambda n)\gamma^\alpha\gamma_5\psi_-(0)\,|PS\rangle\Big] \ , \end{aligned} \tag{4.29}$$

we see that it contains both the 'good' component ψ_+ and the 'bad' one ψ_-.

The external lines in our Feynman diagram represent *free* quark fields;[6] in order to accurately represent $f_T(x)$ with a diagram, we must first eliminate all of the bad quark components

[5] in the + direction

[6] Our Feynman rules have been derived in this way; see Section 1.2.



Figure 4.3: These one-particle reducible diagrams represent the true nature of figure 4.1. They are obtained by eliminating the 'bad' component of ψ via the light-cone equations of motion.

in favor of the good ones.[7] This process turns Figure 4.1 into Figure 4.3. An explicit calculation shows that the contribution from the graphs of Figure 4.3 is identical to (4.8).[8] However, from Fig. 4.3 it is quite clear that the contributions of Figure 4.2 are necessary to ensure QED gauge invariance. The point here is not that (4.8) is wrong, just incomplete. The form of (4.4) misleads us into thinking that it is the only contribution; the more honest form (4.29) reveals the true nature of transverse scattering to be inextricably coupled with multiparton correlation effects. We will see that the effect of this coupling has consequences far greater than the inclusion of a few more diagrams.

4.2 The Evolution Equation

Now that we have seen that transverse polarization cannot be decoupled from coherent partonic correlations, it is natural to ask what kind of effect this has on the analysis of the previous chapters. There, we were able to factorize the DIS cross-section into soft and hard sectors due to the *incoherence* of the dominant scattering mechanism. It is clear that the introduction of correlation effects will have some kind of an impact on this analysis, but its full consequences are not obvious.

One immediate effect of coherent scattering is a complication of the soft sector of our

[7] Note that this requirement is quite independent of QCD. We would have to perform this manipulation even in a theory without local gauge invariance, where it would lead only to the transverse momentum correction.

[8] Strictly speaking, these gluons must be combined with the related transverse momentum insertion to reproduce (4.8). For this reason, it is often advisable to begin with bad components and eliminate them via (4.28).

amplitude. If a factorization theorem does indeed exist at this level, the matrix elements representing the soft physics certainly must encode a great deal more information than those previously studied. The distribution in Equation (4.4) does not look much different from those we met in Section 2.3, but we have just seen that this form can be quite misleading.

A reliable way to see how much information a distribution contains is by studying its scale evolution. For a self-contained distribution which evolves autonomously, only one measurement is enough to determine the complete behavior. On the other hand, distributions which mix with each other require external inputs for a study of their scale dependence. As we have seen, mixing is an indication that a distribution does not contain all of the relevant physics.

Since operators of different twist do not belong to the same representation of the Lorentz group, we must separate the twist-2 and -3 contributions to $f_T(x,\mu^2)$ in order to study its evolution effectively. This is most easily done with the local operators that constitute its moments rather than the function itself. The $(n+1)$-st moment of $f_T(x,\mu^2)$ is written

$$\int dx\, x^n\, f_T(x,\mu^2) = \frac{1}{2M} n_{\mu_1}\cdots n_{\mu_n} \left\langle PS \left| \theta_n^{\perp(\mu_1\cdots\mu_n)} \right| PS \right\rangle \quad , \tag{4.30}$$

where

$$\theta_n^{\sigma(\mu_1\cdots\mu_n)} = \overline{\psi}\gamma^\sigma\gamma_5 i\mathcal{D}^{(\mu_1}\cdots i\mathcal{D}^{\mu_n)}\psi \quad , \tag{4.31}$$

with $(\mu_1\cdots\mu_n)$ indicating symmetrization of the indices and removal of the traces. The twist-2 $\theta^{(\sigma\mu_1\cdots\mu_n)}$ (totally symmetric and traceless) and twist-3 $\theta^{[\sigma(\mu_1]\mu_2\cdots\mu_n)}$ (mixed symmetric and traceless) contributions, where $[\sigma\mu_1]$ denotes antisymmetrization, can now be separated quite easily via

$$\theta^{\sigma(\mu_1\cdots\mu_n)} = \theta_{n2}^{(\sigma\mu_1\mu_2\cdots\mu_n)} + \frac{2n}{n+1}\theta_{n3}^{[\sigma(\mu_1]\mu_2\cdots\mu_n)} + \cdots \quad , \tag{4.32}$$

where the ellipses denote terms of higher twist.

As mentioned above, the twist-2 part of θ_n is identical to the operator related to the $(n+1)$st moment of $\tilde{f}(x)$. Its evolution has already been studied in Section 2.5. We argued in that section that the twist-2 nonsinglet quark operators must have autonomous evolution because there are simply no operators with which they can mix. This statement is not true at the twist-3 level. A complete basis of twist-3 quark operators was first identified in [84],

$$\begin{aligned} R_i^n &= \overline{\psi} i\mathcal{D}^{(\mu_1}\cdots i\mathcal{D}^{\mu_{i-1}}(-ig)\mathcal{F}^{\sigma\mu_i} i\mathcal{D}^{\mu_{i+1}}\cdots i\mathcal{D}^{\mu_{n-1}}\gamma^{\mu_n)}\gamma_5\psi \\ S_i^n &= \overline{\psi} i\mathcal{D}^{(\mu_1}\cdots i\mathcal{D}^{\mu_{i-1}} \qquad g\tilde{\mathcal{F}}^{\sigma\mu_i} i\mathcal{D}^{\mu_{i+1}}\cdots i\mathcal{D}^{\mu_{n-1}}\gamma^{\mu_n)} \quad \psi \quad , \end{aligned} \tag{4.33}$$

where $i = 1, ..., n-1$. The operator $\theta_{n3}^{[\sigma(\mu_1]\mu_2\cdots\mu_n)}$ is just a special linear combination of them:

$$\theta_{n3}^{[\sigma(\mu_1]\mu_2\cdots\mu_n)} = \frac{1}{4n}\sum_{i=1}^{n-1}(n-i)(R_i^n - R_{n-i}^n + S_i^n + S_{n-i}^n) \ . \tag{4.34}$$

The anomalous dimension matrix in the above operator basis was first worked out by Bukhvostov et al. and later reproduced by a number of authors with different methods [85]. The result is what one would generally expect. To evolve the matrix element of $\theta^{[\sigma(\mu_1]\mu_2\cdots\mu_n)}$, it is not enough just to know it at an initial scale - one must know all the matrix elements of $W_i^n = R_i^n - R_{n-i}^n + S_i^n + S_{n-i}^n$ there. Important physical information pertaining to the soft physics of our process is simply not probed by the experiment. This is actually quite clear from the discussion in the last section. There, we saw that the distributions $G_1(x,y)$ and $G_2(x,y)$ are in some sense more fundamental than $f_T(x)$. The local operators R_i^n and S_i^n are related to the moments of *these* distributions.

By studying the anomalous dimension matrix in the large-N_c limit,[9] Ali, Braun and Hiller found that the eigenvector corresponding to the lowest eigenvalue is just the linear combination of twist-3 operators on the right-hand side of Eq.(4.34). In other words, the twist-3 part of $f_T(x,\mu^2)$ evolves autonomously in this limit. To better understand ABH's result, we calculate the large-N_c evolution of $f_T(x,\mu^2)$ directly. Starting with the mixed-twist operator $\theta_n^{\sigma(\mu_1\mu_2\cdots\mu_n)}$ in Eq.(4.30), we look for possible divergences when inserted in multi-point Greens functions. To reduce the number of Feynman diagrams, we choose the light-cone gauge $\mathcal{A}^+ = 0$ and take the $\perp + \cdots +$ component of the θ-operator. The resulting operator,

$$\theta_n \equiv \theta_n^{\perp+\cdots+} = \overline{\psi}\gamma^\perp\gamma_5(i\partial^+)^n\psi \ , \tag{4.35}$$

has the simple Feynman rule $\gamma^\perp\gamma_5(k^+)^n$, where k is the momentum of the incoming quark.

By light-cone power counting, we need only consider two- and three-point functions. Since the external lines carry color, we must ask what type of diagrams dominate the large-N_c limit. As shown in Appendix C.3, the simple rule is that when all external lines are drawn to one point (infinity) the planer diagrams are leading. All one-particle-irreducible (1PI) leading two- and three-point diagrams with one θ insertion are shown in Fig. 4.4.

The ultraviolet divergences in the two point Greens function can obviously be subtracted with the matrix element of θ_n itself. The only diagram in which the divergences may not be

[9] Large-N_c physics is discussed in Appendix C.3.




Figure 4.4: The (a) two- and (b) three-point 1PI Feynman diagrams contributing to the evolution of θ_n in the large-N_c limit.

subtracted by θ_n is Fig. 4.4b. An explicit calculation shows that the ultraviolet divergences here require a subtraction of the local operator

$$-\frac{1}{2}C_A\frac{g^2}{8\pi^2}\frac{2}{\epsilon}\left(\frac{\mu^2 e^{\gamma_E}}{4\pi\mu_0^2}\right)^{-\epsilon/2}\left[-\frac{1}{(n+2)}\sum_{i=0}^{n-1}\overline{\psi}\;\not{n}\gamma_5(i\partial^+)^i i\mathcal{D}^\perp(i\partial^+)^{n-1-i}\psi\right.$$
$$\left.+\left(\sum_{i=1}^{n+1}\frac{1}{i}-\frac{1}{2(n+1)}\right)\left(\overline{\psi}\;i\not{\mathcal{D}}_\perp\not{n}\gamma^\perp\gamma_5(i\partial^+)^{n-1}\psi+\overline{\psi}(i\partial^+)^{n-1}\gamma^\perp\gamma_5\;\not{n}\;i\not{\mathcal{D}}_\perp\psi\right)\right]\quad, \tag{4.36}$$

where I have neglected the contributions of light-cone singularities which will be canceled eventually. Notice that the first term is present in the twist-2 operator

$$\theta_{n2}=\frac{1}{n+1}\left(\overline{\psi}\gamma^\perp\gamma_5(i\partial^+)^n\psi+\sum_{i=0}^{n-1}\overline{\psi}\gamma^+\gamma_5(i\partial^+)^i i\mathcal{D}^\perp(i\partial^+)^{n-i-1}\psi\right)\;, \tag{4.37}$$

and the remaining two terms can be converted to θ_n by using the equation of motion (4.28). Thus we easily arrive at the ABH conclusion that θ_{n3} evolves autonomously in the large-N_c limit.

Including the contribution from Fig. 4.4a as well as the wavefunction renormalization, we obtain the evolution equation

$$\mu^2\frac{d\theta_n}{d\mu^2}=\frac{\alpha_s(\mu^2)}{2\pi}\frac{N_c}{2}\left[\frac{n+1}{n+2}\theta_{n2}+\left(-2\sum_{i=1}^{n+1}\frac{1}{i}+\frac{1}{n+1}+\frac{1}{2}\right)\theta_n\right]\;. \tag{4.38}$$

Separating out the twist-2 and twist-3 parts, we not only recover the well-known twist-2 evolution,[10]

$$\mu^2\frac{d\theta_{n2}}{d\mu^2}=\frac{\alpha_s(\mu^2)}{2\pi}\;\frac{N_c}{2}\left(\frac{3}{2}+\frac{1}{(n+1)(n+2)}-2\sum_{i=1}^{n+1}\frac{1}{i}\right)\;, \tag{4.39}$$

[10] Note that this is the evolution of the $(n+1)$st moment of the distribution rather than the nth.

but also the twist-3 result

$$\mu^2 \frac{d\theta_{n3}}{d\mu^2} = \frac{\alpha_s(\mu^2)}{2\pi}\ \frac{N_c}{2}\left(-2\sum_{i=1}^{n+1}\frac{1}{i}+\frac{1}{n+1}+\frac{1}{2}\right)\theta_{n3}\ \ , \tag{4.40}$$

which is identical to the result in Ref. [36].

It is quite clear that the i-independence of the coefficients in the sum in Eq.(4.36) is the key for the autonomous evolution of θ_{n3}. This property is not totally unexpected if one inspects Fig. 4.4b more closely. Interpreting this diagram in coordinate space, we see that the internal gluon propagates *homogeneously* from one quark to the other. By homogeneously, we mean that at any point along the path of the propagation, the gluon behaves exactly the same way, except, of course, at the points where the gluon and quarks interact. The spatial location of the interaction with the external gluon determines the number of derivatives before and after the gluon field in the subtraction operator. Since the internal gluon propagation is homogeneous, different locations of the triple-gluon vertex produce similar physical effects. Therefore, the coefficients of the subtraction operators $\overline{\psi}\,\not{n}\gamma_5(i\partial^+)^i iD^\perp(i\partial^+)^{n-1-i}\psi$ will be independent of i. The two extra terms in Eq.(4.36) correspond to the triple-gluon vertex just next to the external quark lines, where the homogeneity is lost. However, these diagrams cannot break the symmetry since they can always be discarded in favor of the 'bad' quark component via the equation of motion (4.28).

Since the homogeneous property of the internal gluon line is independent of the γ-matrix structure of the operator inserted, we conclude that the other twist-3 distributions $e(x,\mu^2)$ and $h_L(x,\mu^2)$ [11] also evolve autonomously in the large-N_c limit. A quick calculation confirms the evolution equations found in Ref. [82].

Encouraged by the success of the above approach, we apply it to the analogous twist-4 evolution. In Ref. [86], the three one-variable distributions $f_4(x,\mu^2)$, $g_3(x,\mu^2)$ and $h_3(x,\mu^2)$ are shown to contain twist-4. For example, $f_4(x)$ is defined as

$$f_4(x) = \frac{1}{M^2}\int\frac{d\lambda}{2\pi}\langle P|\overline{\psi}(0)\gamma^-\psi(\lambda n)|P\rangle\ . \tag{4.41}$$

It was shown in Ref. [87] that $f_4(x,Q^2)$ contributes to the $1/Q^2$ term of the longitudinal

[11] These distributions are defined in [64].




Figure 4.5: Four-point 1PI Feynman diagrams contributing to the evolution of $\hat{O}$ in the large-N_c limit.

scaling function F_L of the nucleon

$$F_L(x_B, Q^2) = \frac{2x_B^2 M^2}{Q^2} \sum_a e_a^2 f_{4a}(x_B, Q^2) \ , \tag{4.42}$$

where we have neglected higher-order radiative corrections. Here, autonomous evolution of $f_4(x, \mu^2)$ would simplify the analysis of F_L data immensely.

In the large-N_c limit, we consider one insertion of the operator $\hat{O} = \overline{\psi}\gamma^-(i\partial^+)^n\psi$ into two-, three- and four-point Green's functions. At one-loop order, the 1PI two- and three-point diagrams are identical to those in Fig. 4.4 and the 1PI four-point diagrams are shown in Fig. 4.5. Only the three and four point diagrams can potentially destroy the autonomous evolution of $\hat{O}$. Let us start with Fig. 4.5a. One of its divergent contributions introduces the local subtraction

$$\sum_i \overline{\psi}\, i\not{\partial}_\perp \not{n} (i\partial^+)^i i\not{\partial}_\perp (i\partial^+)^{n-i-2}\psi + \text{h.c.} \ , \tag{4.43}$$

where the coefficients are independent of i because of the homogeneity of the gluon propagator. Using the equation of motion, we can write this as

$$\sum_i \overline{\psi}(i\partial^+)^i i\, \not{\partial}_\perp (i\partial^+)^{n-i-2}\psi + \text{h.c.} \tag{4.44}$$

Since this operator cannot be reduced to either the twist-2 or twist-4 part of $\hat{O}$, the evolution of the latter cannot be autonomous unless this contribution is canceled by other diagrams. The only other diagram containing the same divergence structure is Fig. 4.4b. Unfortunately, our explicit calculation did not produce this cancelation. Here, it is not the inhomogeneity of gluon propagation which destroys the autonomous evolution. In the twist-4 sector, there are simply more operators which do not distinguish between quark fields (are derived from homogeneous




Figure 4.6: Some two-loop 1PI Feynman diagrams that might break the autonomy of the θ_n in the large-N_c limit.

propagation) than appear in the decomposition of $\hat{O}$ into twist-2 and -4 contributions. The same phenomenon occurs for the twist-4 part of $g_3(x,\mu^2)$ and $h_3(x,\mu^2)$.

Thus, the large-N_c simplification seems to happen only for the evolution of the twist-3 part of $f_T(x,\mu^2)$, $h_L(x,\mu^2)$ and $e(x,\mu^2)$. Does it happen for them at two and higher loops? In Fig. 4.6, we see two examples of Feynman diagrams suspected to break the autonomy of the θ_{n3} evolution, i.e. they may contain divergences that cannot be subtracted by θ_{n2} and θ_{n3} only. The inhomogeneity of the gluon propagator leads to this suspicion. The internal gluon that propagates from one quark to another has different wavelengths in the different parts of the propagation. Its interaction with the external gluon will therefore be different at different spatial locations. Thus the subtraction operators have different coefficients depending on the number of derivatives before and after the external gluon field. An explicit calculation of Fig. 4.6a confirms this suspicion.

This leaves us with only one possibility for autonomous two-loop evolution of θ_{n3}: the unwanted structures cancel in the sum of all large-N_c two-loop diagrams. Calculating all those diagrams is a big task. However, even without an explicit calculation, we do not expect the cancelation to happen. The fundamental reason is that the large-N_c limit represents only a selection of a subset of Feynman diagrams, whereas the result of an individual diagram is independent of this limit. Cancelations of a structure do not generally happen among Feynman diagrams unless there is a symmetry.

4.3 Conclusions

Although deep-inelastic scattering with transversely polarized targets seems quite similar to that with longitudinal polarization, a closer look reveals it to be quite deeply entangled with coherent scattering effects. This entanglement stems from the fact that transverse polarization *induces* a transverse momentum distribution which *cannot* be decoupled from the effects of physical gluons.

A study of the evolution of the distribution function relevant to this process unearths a large number of coupled operators which have been obscured from view by the equations of motion. These operators produce an extremely complicated evolution for the twist-3 part of the measured structure function, with the effect that scale variation of the distribution function *cannot* be performed until one knows the full functional form of the more fundamental distributions $G_1(x, y)$ and $G_2(x, y)$. Unfortunately, these distributions are not readily extracted separately in experiment. Only the combination appearing in $f_T(x)$ shows up at leading order in DIS.

A calculation of Ali, Braun, and Hiller shows that the anomalous dimension matrix of these operators simplifies considerably in the limit of large number of colors. The result of this simplification is that the twist-3 part of $f_T(x)$ evolves autonomously in one-loop after all. Corrections to this evolution appear at $\mathcal{O}(1/N_c^2)$, so it should be trusted to approximately 10%. This amazing result represents an incredible simplification of the data analysis on polarized DIS and allows us to extract valuable information on gluon-quark correlations within the proton.

This simplification naturally leads us to ask if there are any other cases in which it occurs. Later calculations have shown that similar simplifications do indeed occur for the twist-3 parts of other distributions relevant to experiment. An extension of the method used by Ali, Braun, and Hiller to twist-4 and higher or to two or more loops would be a heroic undertaking. Here, we have shown how to avoid a calculation of the entire anomalous dimension matrix and get their result directly. Furthermore, using this method, one can see clearly *why* the simplification occurs.

We have seen that the large-N_c simplification happens because this limit, quite accidentally it seems, happens to throw out exactly the diagrams which induce mixing. In the light-cone gauge, the simplification can easily be traced to a special property of Fig. 4.4b. Unfortunately,

this special property is not enough to ensure autonomous evolution of the analogous twist-4 distributions at one loop, and does not persist to higher orders in the twist-3 case. Nonetheless, the discovery of Ali, Braun, and Hiller remains as a significant step forward in the study of the $f_T(x, \mu^2)$ structure function. Without the autonomous one-loop evolution, an analysis of experimental data on the twist-3 contribution would be severely constrained.

Appendix A

Standard Conventions and Formulae

This Appendix is meant to provide a table of standard formulae and conventions used in the text.

We work in 'God-given' units, where $c = \hbar = 1$.

Throughout, I use the metric

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \tag{A.1}$$

of signature -2.

The Levi-Cevita symbol is defined via $\epsilon^{0123} = +1$.

When written explicitly, I use the convention that the time component of a 4-vector goes first, followed by space. For example, $p^\mu = (p^0, p^1, p^2, p^3)$.

Indices in Greek at the beginning of the alphabet $(\alpha, \beta, \ldots)$ are usually Dirac indices. Exception : α and β are fair game for use as Lorentz indices.

Indices in Greek in the middle of the alphabet $(\mu, \nu, \ldots)$ are usually Lorentz indices.

Indices in Latin at the beginning of the alphabet $(a, b, \ldots)$ are usually in the adjoint representation of $SU(3)$.

Indices in Latin in the middle of the alphabet $(i, j, \ldots)$ are usually either spatial indices or in the fundamental or conjugate representation of $SU(3)$.

Context is really the final word here. If it *should be* a Lorentz index, it probably is...

The summation convention is everywhere in effect (unless specified otherwise). This means

that any repeated index in the same term of an expression is meant to be summed over all possible values of that index. In certain spaces, invariant summing requires one contravariant (up) and one covariant (down) index.

Suppression of indices implies contraction in the most obvious way. Either last index with first index or covariant index with contravariant (read from left to right). Exception : for double contraction on two indices my convention is $\theta_{\mu\nu}J^{\mu\nu} \equiv \theta J$.

The difference between the index i and the $\sqrt{-1} = i$ should be clear from context. I have tried to avoid grossly misleading presentations, but borderline cases may be present.

$[,]$ denotes the commutator, defined by $[A, B] \equiv AB - BA$.

$\{,\}$ denotes the anticommutator, defined by $\{A, B\} \equiv AB + BA$.

The symbol $\sim$ is used to indicate either approximate equality or equality in all important respects. In most cases, it can be read, "is essentially the same as".

The conjugation operators I employ are

T^* for complex

$T^\dagger$ for Hermitian

$\overline{T}$ for Dirac.

The notation $\not{a}$ is shorthand for $a_\mu\gamma^\mu$.

The absence of an explicit matrix structure where one should be indicates the presence of the identity.

The absence of integration limits implies integration over the *entire* relevant space. Sometimes, integration is arrested by the support of its integrand.

I do not differentiate explicitly between scalars and 4-vectors. This should be clear from the context. Spatial 3-vectors *always* appear with vector hats, i.e. $\vec{v}$.

I do not specify the number of δ-functions appearing in an expression explicitly unless this is not obvious from the context. For example, $\delta(\vec{p} - \vec{p}\,') \equiv \delta(p^1 - p'^1)\delta(p^2 - p'^2)\delta(p^3 - p'^3)$. One important exception is the volume of spacetime, $(2\pi)^3\,\delta^{(3)}(0)$.

A.1 Dirac Algebra

The γ-matrices are d $2^{d/2} \times 2^{d/2}$-dimensional operators on the $2^{d/2}$-dimensional Dirac spinor space defined by the *Clifford* algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}\mathbf{1} \ , \tag{A.2}$$

where $g^{\mu\nu}$ is the metric tensor of our space. In four dimensions, one defines

$$\gamma_5 \equiv i\gamma^0\gamma^1\gamma^2\gamma^3 = -\frac{i}{4!}\epsilon^{\mu\nu\lambda\rho}\gamma_\mu\gamma_\nu\gamma_\lambda\gamma_\rho \tag{A.3}$$

so that the projection operators

$$P_R \equiv \frac{1}{2}(1+\gamma_5) \qquad\qquad P_L \equiv \frac{1}{2}(1-\gamma_5) \tag{A.4}$$

project out positive (right-handed) and negative (left-handed) helicity states, respectively. The *Weyl* basis is defined by the choice

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \overline{\sigma}^\mu & 0 \end{pmatrix} \ , \tag{A.5}$$

where σ^i are the Pauli matrices,

$$\sigma^1 \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \ , \qquad \sigma^2 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \ , \qquad \sigma^3 \equiv \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \ , \tag{A.6}$$

σ^0 is the 2×2 identity, and

$$\overline{\sigma}^\mu \equiv \sigma_\mu \ . \tag{A.7}$$

In this basis, γ_5 takes the form

$$\gamma_5 = \begin{pmatrix} -\mathbf{1} & 0 \\ 0 & \mathbf{1} \end{pmatrix} \ , \tag{A.8}$$

where $\mathbf{1}$ represents the 2×2 identity. The generators of Lorentz transformations,

$$\sigma^{\mu\nu} \equiv \frac{i}{2}[\gamma^\mu, \gamma^\nu] \ , \tag{A.9}$$

take the form

$$\sigma^{0i} = -i\begin{pmatrix} \sigma^i & 0 \\ 0 & -\sigma^i \end{pmatrix} \tag{A.10}$$

$$\sigma^{ij} = \epsilon^{ijk}\begin{pmatrix} \sigma^k & 0 \\ 0 & \sigma^k \end{pmatrix} \ . \tag{A.11}$$

Under Hermitian conjugation, one has

$$(\gamma^\mu)^\dagger = \gamma^0\gamma^\mu\gamma^0 \tag{A.12}$$

$$(\sigma^{\mu\nu})^\dagger = \gamma^0\sigma^{\mu\nu}\gamma^0 \tag{A.13}$$

$$\gamma_5^\dagger = \gamma_5 = -\gamma^0\gamma_5\gamma^0 \ . \tag{A.14}$$

These matrices are linearly independent and, along with the identity and $\gamma^\mu\gamma_5$, form a closed set under multiplication in $d = 4$. The relations

$$\gamma_5^2 = 1 \tag{A.15}$$

$$\sigma^{\mu\nu}\gamma_5 = \frac{1}{2}\, i\, \epsilon^{\mu\nu\lambda\rho}\sigma_{\lambda\rho} \tag{A.16}$$

$$\gamma^\mu\gamma^\nu = g^{\mu\nu} - i\sigma^{\mu\nu} \tag{A.17}$$

$$\gamma^\mu\gamma^\nu\gamma^\lambda = g^{\mu\nu}\gamma^\lambda - g^{\mu\lambda}\gamma^\nu + g^{\nu\lambda}\gamma^\mu + i\epsilon^{\mu\nu\lambda\rho}\gamma_\rho\gamma_5 \tag{A.18}$$

$$\begin{aligned}\gamma^\mu\gamma^\nu\gamma^\lambda\gamma^\rho = {} & g^{\mu\nu}g^{\lambda\rho} - g^{\mu\lambda}g^{\nu\rho} + g^{\nu\lambda}g^{\mu\rho} \\ & -i\big(\ g^{\mu\nu}\sigma^{\lambda\rho} - g^{\mu\lambda}\sigma^{\nu\rho} + g^{\nu\lambda}\sigma^{\mu\rho} \\ & \quad + g^{\lambda\rho}\sigma^{\mu\nu} - g^{\nu\rho}\sigma^{\mu\lambda} + g^{\mu\rho}\sigma^{\nu\lambda}\big) - i\epsilon^{\mu\nu\lambda\rho}\gamma_5\end{aligned} \tag{A.19}$$

simplify the decomposition. In d dimensions, the identities

$$\gamma^\lambda\gamma^\mu\gamma_\lambda = (2-d)\gamma^\mu \tag{A.20}$$

$$\gamma^\lambda\gamma^\mu\gamma^\nu\gamma_\lambda = 2\gamma^\nu\gamma^\mu + (d-2)\gamma^\mu\gamma^\nu = 4g^{\mu\nu} + (d-4)\gamma^\mu\gamma^\nu \tag{A.21}$$

$$\gamma^\lambda\sigma^{\mu\nu}\gamma_\lambda = (d-4)\sigma^{\mu\nu} \tag{A.22}$$

$$\gamma^\lambda\gamma^\mu\gamma^\nu\gamma^\rho\gamma_\lambda = -2\gamma^\rho\gamma^\nu\gamma^\mu + (4-d)\gamma^\mu\gamma^\nu\gamma^\rho \tag{A.23}$$

$$\gamma^\lambda\sigma^{\mu\nu}\gamma^\alpha\gamma_\lambda = 2\gamma^\alpha\sigma^{\mu\nu} + (4-d)\sigma^{\mu\nu}\gamma^\alpha \tag{A.24}$$

$$\begin{aligned}\gamma^\lambda\gamma^{\mu_1}\gamma^{\mu_2}\cdots\gamma^{\mu_n}\gamma_\lambda = {} & 2\gamma^{\mu_n}\gamma^{\mu_1}\gamma^{\mu_2}\cdots\gamma^{\mu_{n-1}} \\ & +(-1)^n\left[2\gamma^{\mu_3}\gamma^{\mu_2}\gamma^{\mu_1}\gamma^{\mu_4}\cdots\gamma^{\mu_n} + (4-d)\gamma^{\mu_1}\gamma^{\mu_2}\cdots\gamma^{\mu_n}\right] \\ & +2\sum_{i=1}^{n-3}(-1)^i\gamma^{\mu_{n-i}}\gamma^{\mu_1}\cdots\gamma^{\mu_{n-i-1}}\gamma^{\mu_{n-i+1}}\cdots\gamma^{\mu_n} \ ; \qquad n > 3\end{aligned} \tag{A.25}$$

prove useful in reducing products of γ-matrices.

In Feynman diagrams, one is often faced with traces of γ-matrices. Due to the fact that the identity is the *only* matrix in our space whose trace is nonzero, a nonzero traces implies

that for each matrix γ^α in the trace there is another matrix γ^β such that $\gamma^\alpha\gamma^\beta$ is proportional to the identity. With this in mind, on can show that the trace formulae

$$\begin{aligned}
\mathrm{Tr}\left[\gamma^\mu\gamma^\nu\right] &= 4g^{\mu\nu} && \text{(A.26)}\\
\mathrm{Tr}\left[\gamma^\mu\gamma^\nu\gamma^\lambda\gamma^\rho\right] &= 4\left(g^{\mu\nu}g^{\lambda\rho} - g^{\mu\lambda}g^{\nu\rho} + g^{\nu\lambda}g^{\mu\rho}\right) && \text{(A.27)}\\
\mathrm{Tr}\left[\gamma^{\mu_1}\cdots\gamma^{\mu_n}\right] &= 0 \qquad\qquad n \text{ odd} && \text{(A.28)}\\
\mathrm{Tr}\left[\gamma^{\mu_1}\cdots\gamma^{\mu_{2n}}\right] &= \mathrm{Tr}\left[\gamma^{\mu_{2n}}\cdots\gamma^{\mu_1}\right] && \text{(A.29)}\\
\mathrm{Tr}\left[\not{a}\gamma^\alpha\not{b}\gamma^\beta\not{c}\gamma_\alpha\not{d}\gamma_\beta\right] &= (d-2)\left\{(4-d)\mathrm{Tr}\left[\not{a}\not{b}\not{c}\not{d}\right] - 16(a\cdot c)(b\cdot d)\right\} && \text{(A.30)}\\
&= 4(2-d)\left\{(8-d)(a\cdot c)(b\cdot d)\right. && \\
&\qquad +(d-4)[(a\cdot b)(c\cdot d) && \text{(A.31)}\\
&\qquad\left. +(a\cdot d)(b\cdot c)]\right\} && \\
\mathrm{Tr}\left[\not{a}\not{b}\not{c}\gamma^\alpha\not{p}\gamma^\beta\not{k}\gamma_\alpha\not{q}\gamma_\beta\right] &= (d-4)(6-d)\mathrm{Tr}\left[\not{a}\not{b}\not{c}\not{p}\not{k}\not{q}\right] && \\
&\qquad +4\mathrm{Tr}\left\{\not{a}\not{b}\not{c}\left[(4-d)(p\cdot k\ \not{q} + k\cdot q\ \not{p}\right)\right. && \text{(A.32)}\\
&\qquad\left. -2p\cdot q\ \not{k}\right]\right\} &&
\end{aligned}$$

are valid in d dimensions.[1] In general, the trace of an even number of γ-matrices is 4 times the sum over all different pairings of indices multiplied by the sign of the permutation. Written in this way, the trace of $2n$ γ-matrices has $(2n)!/2^n n!$ terms.

The trace formulae

$$\begin{aligned}
\mathrm{Tr}\left[\gamma_5\right] &= 0 && \text{(A.33)}\\
\mathrm{Tr}\left[\gamma_5\gamma^\mu\gamma^\nu\right] &= 0 && \text{(A.34)}\\
\mathrm{Tr}\left[\gamma_5\gamma^\mu\gamma^\nu\gamma^\lambda\gamma^\rho\right] &= -4i\epsilon^{\mu\nu\lambda\rho} && \text{(A.35)}
\end{aligned}$$

have meaning only in 4 dimensions.

A.2 $SU(N)$ Algebra

The (N^2-1) generators t^a of the special unitary group in N dimensions normalized to

$$\mathrm{Tr}\left(t^a t^b\right) = \frac{1}{2}\delta^{ab} \qquad \text{(A.36)}$$

[1] Since the trace of the identity is an overall factor, we have taken it as 4 in d dimensions.

satisfy the Lie algebra

$$\left[t^a, t^b\right] = i f^{abc} t^c \ , \tag{A.37}$$

which define the totally antisymmetric structure constants f^{abc} of the group. The generators also satisfy the Fiertz identity

$$(t^a)^i_j (t^a)^k_l = \frac{1}{2}\left(\delta^i_l \delta^k_j - \frac{1}{N}\delta^i_j \delta^k_l\right) \ . \tag{A.38}$$

For $N = 3$, we can use the representation

$$t^1 = \frac{1}{2}\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \ , \qquad t^2 = \frac{1}{2}\begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \ , \tag{A.39}$$

$$t^3 = \frac{1}{2}\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \ , \qquad t^4 = \frac{1}{2}\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \ , \tag{A.40}$$

$$t^5 = \frac{1}{2}\begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix} \ , \qquad t^6 = \frac{1}{2}\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \ , \tag{A.41}$$

$$t^7 = \frac{1}{2}\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} \ , \qquad t^8 = \frac{1}{2\sqrt{3}}\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \ , \tag{A.42}$$

in which the nonzero structure constants are

$\underline{a}$	$\underline{b}$	$\underline{c}$	$\underline{f^{abc}}$
1	2	3	1
1	4	7	$\frac{1}{2}$
1	5	6	$-\frac{1}{2}$
2	4	6	$\frac{1}{2}$
2	5	7	$\frac{1}{2}$
3	4	5	$\frac{1}{2}$
3	6	7	$-\frac{1}{2}$
4	5	8	$\frac{\sqrt{3}}{2}$
6	7	8	$\frac{\sqrt{3}}{2}$

(A.43)

The symmetric structure constants of the group, d^{abc}, are defined by

$$\{t^a, t^b\} = \frac{1}{N}\delta^{ab} + d^{abc}t^c \quad . \tag{A.44}$$

In terms of these constants, one can write

$$t^a t^b = \frac{1}{2}\left(\frac{1}{N}\,\delta^{ab} + if^{abc}t^c + d^{abc}t^c\right) \quad . \tag{A.45}$$

The Jacobi identities,

$$\left[\mathbf{F}^a, \mathbf{F}^b\right] = if^{abc}\mathbf{F}^c \quad , \tag{A.46}$$

$$\left[\mathbf{F}^a, \mathbf{D}^b\right] = if^{abc}\mathbf{D}^c \quad , \tag{A.47}$$

where $(\mathbf{F}^a)^{b\,c} \equiv -if^{abc}$ and $(\mathbf{D}^a)^{b\,c} \equiv d^{a\,b\,c}$, and the relation

$$f^{abe}f^{cde} = \frac{2}{N}\left(\delta^{ac}\delta^{bd} - \delta^{ad}\delta^{bc}\right) + \left(d^{ace}d^{bde} - d^{ade}d^{bce}\right) \tag{A.48}$$

follow from repeated use of (A.37) and (A.44). These expressions can be used to show the validity of

$$t^a\,t^a = C_F = \frac{N}{2} - \frac{1}{2N} \tag{A.49}$$

$$t^b\,t^a\,t^b = \left(C_F - \frac{C_A}{2}\right)t^a = -\frac{1}{2N}\,t^a \tag{A.50}$$

$$\mathrm{Tr}\left[\,t^a\,t^b\,t^c\,\right] = \frac{1}{4}\left(d^{abc} + if^{abc}\right) \tag{A.51}$$

$$\mathrm{Tr}\left[\mathbf{F}^a\mathbf{D}^b\right] = 0 \tag{A.52}$$

$$\mathrm{Tr}\left[\mathbf{F}^a\mathbf{F}^b\right] = C_A\delta^{ab} \tag{A.53}$$

$$\mathrm{Tr}\left[\mathbf{D}^a\mathbf{D}^b\right] = \left(N - \frac{4}{N}\right)\delta^{ab} \tag{A.54}$$

$$\mathrm{Tr}\left[\mathbf{F}^a\mathbf{D}^b\mathbf{D}^c\right] = \left(\frac{N}{2} - \frac{2}{N}\right)if^{abc} \tag{A.55}$$

$$\mathrm{Tr}\left[\mathbf{F}^a\mathbf{F}^b\mathbf{D}^c\right] = \frac{N}{2}\,d^{abc} \tag{A.56}$$

$$\mathrm{Tr}\left[\mathbf{F}^a\mathbf{F}^b\mathbf{F}^c\right] = \frac{N}{2}\,if^{abc} \tag{A.57}$$

$$\mathrm{Tr}\left[\mathbf{D}^a\mathbf{D}^b\mathbf{D}^c\right] = \left(\frac{N}{2} - \frac{6}{N}\right)d^{abc} \quad . \tag{A.58}$$

A.3 Covariant integration

The standard integrals one encounters in a dimensionally regularized quantum field theory with covariant gauge-fixing are of the form

$$\int \prod_{i=1}^{\ell} \frac{d^d l_i}{(2\pi)^d} N(\{l_i\},\{p_j\},\{m_k\}) \prod_{i=1}^{N_p} \frac{1}{k_i^2 - m_i^2} \ , \tag{A.59}$$

where $\{l_i\}$ are the loop momenta, $\{k_i\}$ the propagator momenta, $\{m_i\}$ the relevant masses, and $\{p_i\}$ external momenta. N is a definite function of the loop momenta (or, if one prefers, the propagator momenta), the external momenta, and the masses. The propagator momenta are necessarily linear in the loop momenta and can be chosen to have coefficient 1, so the technique of Feynman parameters (along with the use of Lorentz symmetry), can always reduce this integral to the form of (E.23) at the expense of introducing N_p new integration parameters. His technique is based on the identity

$$\frac{1}{AB} = \int_0^1 dx\, dy\, \delta(1-x-y)\, \frac{1}{[xA+yB]^2} \ . \tag{A.60}$$

Differentiating with respect to B gives

$$\frac{1}{AB^2} = \int_0^1 dx\, dy\, \delta(1-x-y)\, \frac{2y}{[xA+yB]^3} \ , \tag{A.61}$$

which immediately allows one to derive

$$\begin{aligned} \frac{1}{ABC} &= \int_0^1 dw\, dx\, dy\, dz\, \delta(1-x-y)\, \delta(1-z-w)\, \frac{2w}{[wxA+wyB+zC]^3} \\ &= \int_0^1 dx\, dy\, dz\, \delta(1-x-y-z)\, \frac{2}{[xA+yB+zC]^3} \ . \end{aligned} \tag{A.62}$$

This process continues; it is easy to show by induction that the formulae

$$\frac{1}{A_1 \cdots A_n} = \int_0^1 dx_1 \cdots dx_n\, \delta\left(\sum x_i - 1\right) \frac{\Gamma(n)}{[\sum x_i A_i]^n} \tag{A.63}$$

$$\frac{1}{A_1^{m_1} \cdots A_n^{m_n}} = \int_0^1 dx_1 \cdots dx_n\, \delta\left(\sum x_i - 1\right) \frac{\prod x_i^{m_i - 1}}{[\sum x_i A_i]^{\sum m_i}} \frac{\Gamma(\sum m_i)}{\prod \Gamma(m_i)} \tag{A.64}$$

hold for integer m_i. Furthermore, the right hand side is an analytic function of m_i that can be continued throughout the complex plane. I find the decoupled formulae

$$\frac{1}{A_1 \cdots A_{n+1}} = \int_0^1 dx_1 \cdots dx_n \frac{\Gamma(n+1) \prod x_i^{n-i}}{\left[\sum A_i (1-x_i) \prod^{i-1} x_j\right]^{n+1}} \tag{A.65}$$

$$\frac{1}{A_1^{m_1}\cdots A_{n+1}^{m_{n+1}}} = \int_0^1 dx_1\cdots dx_n \frac{\left[\prod x_i^{n-i}\right]\prod\left[(1-x_i)\prod^{i-1}x_j\right]^{m_i-1}}{\left[\sum A_i(1-x_i)\prod^{i-1}x_j\right]^{\sum m_i}}\frac{\Gamma\left(\sum m_i\right)}{\prod\Gamma\left(m_i\right)} \ , \qquad \text{(A.66)}$$

where I have defined $x_{n+1}\equiv 0$ and $\prod^0\equiv 1$, much more useful in practical calculations.

Once one has combined all of the denominators in a Feynman integral, one can shift the integration variable to get rid of all of the directional dependence of the resulting 'monster denominator' via[2]

$$l^2(l-p_1)^2(l-p_2)^2\cdots(l-p_n)^2 \to \left[\left(l-\sum x_ip_i\right)^2-\left(\sum x_ip_i\right)^2+\sum x_ip_i^2\right]^{n+1} \ . \qquad \text{(A.67)}$$

Any directional dependence in the numerator is illusory, as one can always remove external vectors from the integral, leaving a free index,[3] so we are left with a directionally independent integral. Using the symmetry of the problem, one can now eliminate all free indices via

$$\int\frac{d^dl}{(2\pi)^d}\frac{l^\mu(l^2)^\alpha}{(l^2+M^2)^\beta} = 0 \qquad \text{(A.68)}$$

$$\int\frac{d^dl}{(2\pi)^d}\frac{l^\mu l^\nu(l^2)^\alpha}{(l^2+M^2)^\beta} = \frac{g^{\mu\nu}}{d}\int\frac{d^dl}{(2\pi)^d}\frac{(l^2)^{\alpha+1}}{(l^2+M^2)^\beta} \ , \qquad \text{(A.69)}$$

or the higher-rank equivalent. The remaining d-dimensional integral is evaluated via

$$\int\frac{d^dl}{(2\pi)^d}\frac{(l^2)^\alpha}{(l^2+M^2)^\beta} = \frac{1}{(4\pi)^{d/2}}\left(M^2\right)^{d/2+\alpha-\beta}\frac{\Gamma(d/2+\alpha)\Gamma(\beta-\alpha-d/2)}{\Gamma(d/2)\Gamma(\beta)} \ . \qquad \text{(A.70)}$$

Since this always gives a power of M^2, which is invariably quadratic in the loop momenta, this procedure continues until all of the d-dimensional integrals have been taken. One is then left only with one-dimensional integrals, which are in principle straightforward.

The following formulae are useful in this context

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin\pi z} \qquad \text{(A.71)}$$

$$\Gamma\left(\frac{z+1}{2}\right) = 2^{-z}\sqrt{\pi}\frac{\Gamma(z+1)}{\Gamma(1+z/2)} \qquad \text{(A.72)}$$

[2]This expression assumes the use of the coupled parameter formula (A.63) and ignores masses. The sums run from 1 to n, with x_{n+1} determined by the δ-function in (A.63).

[3]In multi-loop calculations, one may find one's self with fractional powers in the numerator. These powers are to be considered part of the denominator and combined via the above approach. One must be careful in these situations since fractional powers can easily lead to ambiguities in the complex plane. To treat these integrals consistently, one always performs Wick rotations *before* combining denominators and integrating. In this way, there will always be an *integer* number of signs.

$$\Gamma(2z) = 2^{2z-1}\pi^{-1/2}\Gamma(z)\Gamma(z+1/2) \tag{A.73}$$

$$\log[\Gamma(1+\epsilon)] = -\gamma_E\epsilon + \sum_{n=2}^{\infty}\frac{(-1)^n}{n}\zeta(n)\epsilon^n \tag{A.74}$$

$$\int_0^1 dx\, x^{\alpha-1}(1-x)^{\beta-1} = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}\ . \tag{A.75}$$

For more complicated calculations,

$$\int_0^1 dx\, dy\, x^{\alpha-1}(1-x)^{\beta-1}y^{\gamma-1}(1-y)^{\alpha-\gamma-1}(1-xy)^{\mu} = \frac{\Gamma(\beta)\Gamma(\gamma)\Gamma(\alpha+\beta+\mu-\gamma)}{\Gamma(\alpha+\beta+\mu)\Gamma(\alpha+\beta-\gamma)} \tag{A.76}$$

may also be of use. In addition, the integrals

$$\int \frac{d^dk}{(2\pi)^d}\frac{1}{(k^2)^\alpha[(k-q)^2]^\beta} = \frac{(-q^2)^{d/2-\alpha-\beta}}{(4\pi)^{d/2}}\frac{\Gamma(d/2-\beta)\Gamma(d/2-\alpha)\Gamma(\alpha+\beta-d/2)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(d-\alpha-\beta)}\ , \tag{A.77}$$

$$\int \frac{d^dk}{(2\pi)^d}\frac{k^\mu}{(k^2)^\alpha[(k-q)^2]^\beta} = \frac{(-q^2)^{d/2-\alpha-\beta}}{(4\pi)^{d/2}}\frac{\Gamma(d/2-\beta)\Gamma(d/2+1-\alpha)\Gamma(\alpha+\beta-d/2)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(d+1-\alpha-\beta)}\,q^\mu\ , \tag{A.78}$$

$$\int \frac{d^dk}{(2\pi)^d}\frac{k^\mu k^\nu}{(k^2)^\alpha[(k-q)^2]^\beta} = \frac{(-q^2)^{d/2-\alpha-\beta}}{(4\pi)^{d/2}}\frac{\Gamma(d/2-\beta)\Gamma(d/2+1-\alpha)\Gamma(\alpha+\beta-d/2)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(d+2-\alpha-\beta)} \times\left[\frac{1}{2}\frac{d/2-\beta}{\alpha+\beta-d/2-1}\,q^2g^{\mu\nu}+\left(\frac{d}{2}+1-\alpha\right)q^\mu q^\nu\right]\ , \tag{A.79}$$

$$\int \frac{d^dk}{(2\pi)^d}\frac{k^\mu k^\nu k^\lambda}{(k^2)^\alpha[(k-q)^2]^\beta} = \frac{(-q^2)^{d/2-\alpha-\beta}}{(4\pi)^{d/2}}\frac{\Gamma(d/2-\beta)\Gamma(d/2+2-\alpha)\Gamma(\alpha+\beta-d/2)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(d+3-\alpha-\beta)} \times\left[\frac{1}{2}\frac{d/2-\beta}{\alpha+\beta-1-d/2}\,q^2\left(g^{\mu\nu}q^\lambda+g^{\nu\lambda}q^\mu+g^{\lambda\mu}q^\nu\right)+\left(\frac{d}{2}+2-\alpha\right)q^\mu q^\nu q^\lambda\right]\ , \tag{A.80}$$

can be done in general and have simple forms. These integrals have *already* been Wick rotated to Euclidean space. Loop momenta with free indices have been left alone, so one must be careful when using these formulae with internal contractions.

In all the above formulae, $\Gamma(n)$ is the Γ-function, defined via

$$\Gamma(n) \equiv \int_0^\infty dx x^{n-1} e^{-x} \tag{A.81}$$

for $\Re e\, n > 0$ and either the reflection identity (A.71) or the recursion relation

$$n\Gamma(n) = \Gamma(n+1) \tag{A.82}$$

elsewhere. It is a meromorphic function of n throughout the complex plane with simple poles at the negative integers and zero, with residues

$$\mathrm{Res}[\Gamma(z), -n] = \frac{(-1)^n}{n!} \ . \tag{A.83}$$

Note that

$$\Gamma(n+1) = n! \tag{A.84}$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \ . \tag{A.85}$$

Its expansion involves the Euler-Masceroni constant

$$\gamma_E \equiv \lim_{n\to\infty}\left[\sum_{i=1}^{n}\frac{1}{i} - \log n\right] = \sum_{n=2}^{\infty}\frac{(-1)^n}{n}\zeta(n) \sim 0.577\,215\,664\,901\,532\,861 \tag{A.86}$$

and the Riemann Zeta function,

$$\zeta(s) = \sum_{i=1}^{\infty}\frac{1}{i^s} \tag{A.87}$$

for $\Re e\, s > 1$. For other values in the complex plane of s, it is defined via

$$\zeta(s) = 2^s\pi^{s-1}\sin\frac{\pi s}{2}\,\Gamma(1-s)\zeta(1-s) \ , \tag{A.88}$$

and found to be a meromorphic function with a simple pole at $s = 1$ with residue

$$\mathrm{Res}[\zeta(s), 1] = 1 \ . \tag{A.89}$$

Values of $\zeta(s)$ at positive even integer s are known via the expansion

$$\frac{x}{e^x - 1} = 1 - \frac{x}{2} + \sum_{n=1}^{\infty}\frac{(-1)^{n+1}}{2^{2n-1}\pi^{2n}}\,\zeta(2n)x^{2n} \ . \tag{A.90}$$

In particular,

$$\zeta(2) = \frac{\pi^2}{6} \tag{A.91}$$

$$\zeta(4) = \frac{\pi^4}{90} \ . \tag{A.92}$$

Using these results in conjunction with (A.88), one can derive

$$\zeta(0) = -\frac{1}{2} \tag{A.93}$$

$$\zeta(-1) = -\frac{1}{12} \tag{A.94}$$

$$\zeta(-3) = \frac{1}{120} \ . \tag{A.95}$$

In addition, consistency between (A.83), (A.87), and (A.88) implies

$$\lim_{\epsilon \to 0} \frac{\zeta(-2n+\epsilon)}{\epsilon} = (-1)^n \, \frac{(2n)! \, \zeta(2n+1)}{2^{2n+1}\pi^{2n}} \tag{A.96}$$

for positive integer n.

For completeness, I also include the standard trancendentals,

$$\pi \sim 3.141\ 592\ 653\ 589\ 793\ 238 \tag{A.97}$$

$$e \sim 2.718\ 281\ 828\ 459\ 045\ 235 \ , \tag{A.98}$$

that appear in our relations.

A.4 QCD and Feynman's Rules of Perturbation Theory

The lagrangian density of massless QCD is written

$$\mathcal{L} = \overline{\psi} \, i \not{\mathcal{D}} \, \psi - \frac{1}{4} \mathcal{F}_a^{\mu\nu} \mathcal{F}^a_{\mu\nu} \ , \tag{A.99}$$

where ψ are the quark fields and $\mathcal{A}$ are the gluon fields. The covariant derivative $\mathcal{D}$ is given by

$$\overrightarrow{\mathcal{D}} \equiv \overrightarrow{\partial} + ig\mathcal{A} \ , \tag{A.100}$$

$$\overleftarrow{\mathcal{D}} \equiv \overleftarrow{\partial} - ig\mathcal{A} \ , \tag{A.101}$$

$$\mathcal{D}^{ab} \equiv \delta^{ab}\partial + g f^{abc} \mathcal{A}_c \ , \tag{A.102}$$

and the gluon field strength, $\mathcal{F}$, is

$$\mathcal{F}_a^{\mu\nu} \equiv -\frac{i}{g} \left[\mathcal{D}^\mu, \mathcal{D}^\nu \right]_a = \partial^\mu \mathcal{A}_a^\nu - \partial^\nu \mathcal{A}_a^\mu - g f^{abc} \mathcal{A}_b^\mu \mathcal{A}_c^\nu \ . \tag{A.103}$$

The Feynman rules for covariant gauge-fixing are shown in Figure A.1. For the use of these rules, see Section 1.3.




Figure A.1: Feynman's rules for a non-abelian gauge theory with covariant gauge-fixing. The straight line represents quark propagation, while the squiggly line represents gluons and the dashed ghosts. In order to avoid index pollution, most of the indices here have been suppressed. The proper pole-prescription is given by adding $i\varepsilon$ to each denominator. Along with each vertex, a δ-function ensuring energy-momentum conservation is implied.

This lagrangian leads to the equations of motion

$$\overrightarrow{\not{\mathcal{D}}}\,\psi = 0 \tag{A.104}$$

$$\overline{\psi}\,\overleftarrow{\not{\mathcal{D}}} = 0 \tag{A.105}$$

$$\mathcal{D}^{ab}_{\nu}\mathcal{F}^{\nu\mu}_{b} = g\overline{\psi}t^{a}\gamma^{\mu}\psi \tag{A.106}$$

and the energy-momentum tensor density[4]

$$\Theta^{\mu\nu} = \overline{\psi}\gamma^{(\mu}i\overleftrightarrow{\mathcal{D}}^{\nu)}\psi - \mathcal{F}^{\mu\alpha}_{a}\mathcal{F}^{\nu}_{a\ \alpha} + \frac{1}{4}g^{\mu\nu}\mathcal{F}^{\alpha\beta}_{a}\mathcal{F}^{a}_{\alpha\beta}\ , \tag{A.108}$$

where $\overleftrightarrow{\mathcal{D}} \equiv \frac{1}{2}\left(\overrightarrow{\mathcal{D}} - \overleftarrow{\mathcal{D}}\right)$.

The coupling of QCD enjoys the β-function

$$\mu^2\frac{d\alpha_s(\mu^2)}{d\mu^2} \equiv \beta(\alpha_s(\mu^2)) = -\frac{1}{4\pi}\,b_0\alpha_s(\mu^2)^2 - \frac{1}{(4\pi)^2}\,b_1\alpha_s(\mu^2)^3 - \cdots\ ; \tag{A.109}$$

$$b_0 = \frac{11}{3}\,C_A - \frac{4}{3}\,n_f T_F \tag{A.110}$$

$$b_1 = \frac{2}{3}\left[17C_A^2 - 2n_f T_F\,(3C_F + 5C_A)\right]\ , \tag{A.111}$$

where $\alpha_s = g^2/4\pi$, $C_A = 3$, $C_F = 4/3$, $T_F = 1/2$, and n_f is the number of active (light) quark flavors.

The fundamental Ward identity of QCD can be written

$$g\,\partial_\mu\left\langle \mathrm{T}\,j^{\mu}_{a}(x)\,\mathcal{O}(x_1,\ldots,x_n)\right\rangle = -i\partial_\mu\left\langle \mathrm{T}\,\frac{\delta\mathcal{O}(x_1,\ldots,x_n)}{\delta\mathcal{A}^{a}_{\mu}(x)}\right\rangle\ , \tag{A.112}$$

where

$$j^{\mu}_{a} = \overline{\psi}\gamma^{\mu}t^{a}\psi + f^{abc}\mathcal{F}^{\mu\alpha}_{b}\mathcal{A}^{c}_{\alpha} \tag{A.113}$$

is the chromodynamic current. This identity is valid for any gauge-invariant operator $\mathcal{O}$, though in covariant gauges one must add a ghost contribution to j^{μ}_{a}.

To compare fields that are separated in spacetime, it is necessary to introduce the gauge link

$$\mathcal{G}_C(x,y) = \mathrm{P}e^{-ig\int_0^1 A_\mu(z(\tau))(dz^\mu/d\tau)\,d\tau}\ , \tag{A.114}$$

[4]This expression is also valid for *massive* quarks, in which case I must specify that the symmetrization of indices in the first term does *not* indicate a removal of the trace; the trace of $\Theta^{\mu\nu}$ is given by

$$\Theta^{\mu}_{\ \mu} = m\overline{\psi}\psi + \frac{\beta(\alpha_s)}{8\pi\alpha_s^2}\,\mathcal{F}^{\mu\nu}_{a}\mathcal{F}^{a}_{\mu\nu}\ . \tag{A.107}$$

which transforms as

$$\mathcal{G}_C(x,y) \to U(x)\mathcal{G}(x,y)U^\dagger(y) \tag{A.115}$$

under a gauge transformation.

The one-loop splitting functions for unpolarized

$$\begin{aligned} P_{NS}(x) &= C_F\ \frac{1+x^2}{1-x} = P_s(x) && \text{(A.116)}\\ P_{sg}(x) &= T_F\ \left[x^2+(1-x)^2\right] && \text{(A.117)}\\ P_{gs}(x) &= C_F\ \frac{1+(1-x)^2}{x} && \text{(A.118)}\\ P_g(x) &= 2C_A\left[\frac{x}{1-x}+\frac{1-x}{x}+x(1-x)\right] && \text{(A.119)} \end{aligned}$$

and polarized

$$\begin{aligned} \tilde{P}_{NS}(x) &= \tilde{P}_s(x) = P_{NS}(x) && \text{(A.120)}\\ \tilde{P}_{sg}(x) &= T_F\ \left[x^2-(1-x)^2\right] && \text{(A.121)}\\ \tilde{P}_{gs}(x) &= C_F\ \frac{1-(1-x)^2}{x} && \text{(A.122)}\\ \tilde{P}_g(x) &= 2C_A\left[1-2x+\frac{1}{1-x}\right] && \text{(A.123)} \end{aligned}$$

parton densities, coupled with the endpoint contributions for quarks

$$A_q = C_F\int_0^x \frac{dy}{x^2}\ \frac{x^2+y^2}{x-y} \tag{A.124}$$

and gluons

$$A_g = \frac{4n_fT_F-11C_A}{6}+2C_A\int_0^x \frac{dy}{x-y} \tag{A.125}$$

generate the DGLAP evolution equation in the nonsinglet

$$\mu^2\frac{D_q}{D\mu^2}f_{NS/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi}\int_x^1\frac{dy}{y}P_{NS}\left(\frac{x}{y}\right)f_{NS/T}(y,\mu^2) \tag{A.126}$$

and singlet

$$\mu^2\frac{D_q}{D\mu^2}f_{s/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi}\int_x^1\frac{dy}{y}\left[P_s\left(\frac{x}{y}\right)f_{s/T}(y,\mu^2)+2n_fP_{sg}\left(\frac{x}{y}\right)f_{g/T}(y,\mu^2)\right] \tag{A.127}$$

$$\mu^2\frac{D_g}{D\mu^2}f_{g/T}(x,\mu^2) = \frac{\alpha_s(\mu^2)}{2\pi}\int_x^1\frac{dy}{y}\left[P_{gs}\left(\frac{x}{y}\right)f_{s/T}(y,\mu^2)+\quad P_g\left(\frac{x}{y}\right)f_{g/T}(y,\mu^2)\right] \tag{A.128}$$

sectors. The polarized evolution equations are obtained simply be adorning these with tildés.[5]

[5] As long as one is careful with the definitions of antiquark distributions. See Section 2.3.

A.5 Other Useful Relations

Here, I present some miscellaneous relations which prove useful in the context of quantum field theory :

$$
\begin{aligned}
[AB, CD] &= A[B, C]D + [A, C]BD + C[A, D]B + CA[B, D] && \text{(A.129)}\\
&= A\{B, C\}D - \{A, C\}BD - C\{A, D\}B + CA\{B, D\} && \text{(A.130)}\\
\epsilon^{\mu\nu\alpha\beta}\epsilon_{\mu\nu\alpha\beta} &= -24 && \text{(A.131)}\\
\epsilon^{\mu\nu\alpha\beta}\epsilon_{\lambda\nu\alpha\beta} &= -6\delta^{\mu}_{\lambda} && \text{(A.132)}\\
\epsilon^{\mu\nu\alpha\beta}\epsilon_{\lambda\rho\alpha\beta} &= -2\left(\delta^{\mu}_{\lambda}\delta^{\nu}_{\rho} - \delta^{\nu}_{\lambda}\delta^{\mu}_{\rho}\right) && \text{(A.133)}\\
\epsilon^{\mu\nu\alpha\beta}\epsilon_{\lambda\rho\sigma\beta} &= -\big(\quad \delta^{\mu}_{\lambda}\delta^{\nu}_{\rho}\delta^{\alpha}_{\sigma} - \delta^{\alpha}_{\lambda}\delta^{\nu}_{\rho}\delta^{\mu}_{\sigma} \\
&\qquad + \delta^{\nu}_{\lambda}\delta^{\alpha}_{\rho}\delta^{\mu}_{\sigma} - \delta^{\mu}_{\lambda}\delta^{\alpha}_{\rho}\delta^{\nu}_{\sigma} && \text{(A.134)}\\
&\qquad + \delta^{\alpha}_{\lambda}\delta^{\mu}_{\rho}\delta^{\nu}_{\sigma} - \delta^{\nu}_{\lambda}\delta^{\mu}_{\rho}\delta^{\alpha}_{\sigma}\big) \\
\epsilon^{\mu\nu\alpha\beta}\epsilon_{\lambda\rho\sigma\tau} &= -\left(\delta^{\mu}_{\lambda}\delta^{\nu}_{\rho}\delta^{\alpha}_{\sigma}\delta^{\beta}_{\tau} + 23 \text{ permutations}\right) && \text{(A.135)}\\
g^{\mu\nu}\epsilon^{\lambda\rho\sigma\tau} &= g^{\mu\lambda}\epsilon^{\nu\rho\sigma\tau} + g^{\mu\rho}\epsilon^{\lambda\nu\sigma\tau} + g^{\mu\sigma}\epsilon^{\lambda\rho\nu\tau} + g^{\mu\tau}\epsilon^{\lambda\rho\sigma\nu} && \text{(A.136)}\\
\frac{1}{(x + i\varepsilon)^n} &= \mathrm{P}\frac{1}{x^n} + (-1)^n \frac{i\pi}{\Gamma(n)} \left(\frac{d}{dx}\right)^{n-1} \delta(x) && \text{(A.137)}\\
\int dx\, e^{ikx} &= 2\pi\delta(k) && \text{(A.138)}
\end{aligned}
$$

and generally :

$$
\begin{aligned}
\sum_{i=1}^{n} &= n && \text{(A.139)}\\
\sum_{i=1}^{n} i &= \frac{n(n+1)}{2} && \text{(A.140)}\\
\sum_{i=1}^{n} i^2 &= \frac{n(n+1)(2n+1)}{6} && \text{(A.141)}\\
\sum_{i=1}^{n} i^3 &= \frac{n^2(n+1)^2}{4} && \text{(A.142)}\\
\sum_{i=0}^{n} i^k &= k!\lim_{t\to 0}\frac{d^k}{dt^k}\,\frac{e^{(n+1)t} - 1}{e^t - 1} && \text{(A.143)}\\
\sum_{i=1}^{n}(-1)^{n-i} &= \frac{1 - (-1)^n}{2} && \text{(A.144)}\\
\sum_{i=1}^{n}(-1)^{n-i}\, i &= \frac{(n+1)}{2} - \frac{1 + (-1)^n}{4} && \text{(A.145)}
\end{aligned}
$$

$$\sum_{i=1}^{n}(-1)^{n-i}\,i^2 = \frac{n(n+1)}{2} \tag{A.146}$$

$$\sum_{i=1}^{n}(-1)^{n-i}\,i^3 = \frac{(n+1)^2(2n-1)}{4} + \frac{1+(-1)^n}{8} \tag{A.147}$$

$$\sum_{i=0}^{n}(-1)^{n-i}\,i^k = \lim_{t\to 0}\frac{d^k}{dt^k}\,\frac{e^{(n+1)t}+(-1)^n}{e^t+1} \tag{A.148}$$

$$\sum_{i=0}^{n} x^i = \frac{1-x^{n+1}}{1-x} \tag{A.149}$$

$$\frac{x^{n-l}y^l - x^{l+j}y^{n-l-j}}{x-y} = \left[\Theta(n-2l-j-1)\sum_{i=l+j}^{n-l-1} -\Theta(2l+j-n-1)\sum_{i=n-l}^{l+j-1}\right] x^i y^{n-i-1} \tag{A.150}$$

$$\frac{1}{1-x} = \sum_{i=0}^{\infty} x^i \tag{A.151}$$

$$\log(1-x) = -\sum_{i=1}^{\infty}\frac{x^i}{i} \tag{A.152}$$

$$\frac{1}{1-x}\,\log(1-x) = -\sum_{i=1}^{\infty} x^i \sum_{j=1}^{i}\frac{1}{j} \tag{A.153}$$

$$\log^2(1-x) = 2\sum_{i=2}^{\infty}\frac{x^i}{i}\sum_{j=1}^{i-1}\frac{1}{j} \tag{A.154}$$

$$(x+y)^n = \sum_{i=0}^{n}\binom{n}{i}x^i y^{n-i} \tag{A.155}$$

$$\binom{n}{i} = \frac{n!}{i!(n-i)!} \tag{A.156}$$

$$\binom{n-1}{i-1} + \binom{n-1}{i} = \binom{n}{i} \tag{A.157}$$

$$\sum_{i=0}^{n}\binom{n}{i} = 2^n \tag{A.158}$$

$$\sum_{i=j}^{n}\binom{i}{j} = \binom{n+1}{j+1} \tag{A.159}$$

$$\sum_{i=0}^{j}\binom{n+i}{i} = \binom{n+j+1}{j} \tag{A.160}$$

$$\sum_{i=j}^{n}\binom{n}{i}\binom{i}{j}x^{n-i}y^{i-j} = \binom{n}{j}(x+y)^{n-j} \tag{A.161}$$

$$\sum_{i=0}^{n}(-1)^i\binom{n}{i} = \delta_{n0} \tag{A.162}$$

$$\sum_{i=0}^{j}(-1)^i\binom{n}{i} = (-1)^j\binom{n-1}{j} \tag{A.163}$$

$$\sum_{i=j}^{n}(-1)^i\binom{n}{i}\binom{i}{j} = (-1)^n\delta_{nj} \tag{A.164}$$

$$\sum_{i=j}^{n}(-1)^i\binom{n}{i-1}\binom{i}{j} = (-1)^n\binom{n+1}{j} \tag{A.165}$$

$$\sum_{i=j}^{n}(-1)^i\binom{n+1}{i+1}\binom{i}{j} = (-1)^j \tag{A.166}$$

$$\sum_{i=0}^{j}\frac{(-1)^i}{\binom{n}{i}} = \frac{n+1}{n+2}\left[1+\frac{(-1)^j}{\binom{n+1}{j+1}}\right] \quad . \tag{A.167}$$

A.6 General References

This section is meant to provide some standard reference material for further reading on the subjects presented in this thesis.

For a general overview of Quantum Field Theory, in order of sophistication :

I. J. R. Aitchison and A. J. G. Hey, *Gauge Theories in Particle Physics*, Institute of Physics Publishing, London, 1989.

M. E. Peskin and D. V. Schroder, *An Introduction to Quantum Field Theory*, Addison-Wesley, Reading, MA, 1995.

S. Weinberg, The Theory of Quantum Fields Vol. I & II, Cambridge University, Cambridge, 1995.

C. Itzykson and J. B. Zuber, *Quantum Field Theory*, McGraw-Hill Inc., New York, NY, 1980.

For a detailed introduction to perturbative QCD :

R. D. Field, *Applications of Perturbative QCD*, Addison-Wesley, Reading, MA, 1989.

For general renormalizability requirements and renormalization theorems :

J. C. Collins, *Renormalization*, Cambridge University, Cambridge, 1984.

For a rigorous study of infrared divergences and factorization theorems :

G. Sterman, *An Introduction Quantum Field Theory,* Cambridge University, Cambridge, 1993.

A. H. Mueller, ed., *Perturbative Quantum Chromodynamics*, World Scientific, Singapore, 1989.

For path integral techniques :

R. J. Rivers, *Path Integral Methods in Quantum Field Theory*, Cambridge University, Cambridge, MA, 1987.

For spinors :

V. B. Berestetskii, E. M. Lifshitz and L. P. Pitaevskii, *Quantum Electrodynamics*, Pergamon Press, New York, NY, 1980.

H. J. W. Müller-Kirsten and A Wiedemann, *Supersymmetry : An Introduction with Conceptual and Calculational Details*, World Scientific, Singapore, 1987.

For group theory :

M. Hamermesh, *Group Theory and Its Application to Physical Problems*, Addison-Wesley Publishing Co., Reading, MA, 1962.

H. Georgi, *Lie Algebras in Particle Physics*, Perseus Books, Reading, MA, 1982.

For integral tables and special mathematical functions :

G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists*, Academic Press, San Diego, CA, 1995.

I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, San Diego, CA, 1994.

M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions*, Dover Publications, New York, NY, 1965.

Appendix B

The Lorentz Group and its Representations

In this appendix, we will discuss some of the more mathematical properties of the spacetime translation and rotation group. First, I define some terms.

A *group* is a collection of objects which satisfy properties

1) Any two elements, x, y, in the group may be composed with each other to from another element of the group, xy or yx. This operation is called *multiplication*.

2) Group multiplication must be *associative*, so that $\forall x, y, z \in g$

$$x(yz) = (xy)z \quad . \tag{B.1}$$

3) g also must possess an *identity*, $\mathbf{1}$, such that $\forall x \in g$

$$x\mathbf{1} = \mathbf{1}x = x \quad . \tag{B.2}$$

4) Each element of the group must have an *inverse*, x^{-1}, associated with it such that

$$xx^{-1} = x^{-1}x = \mathbf{1} \quad . \tag{B.3}$$

Group multiplication does *not* have to be commutative. In fact, it usually isn't. A group whose multiplication operation is commutative is called *abelian*.

A *representation* of a group is a collection of objects, one for each element of the group, which satisfy the group multiplication law. It is called a representation because it represents the abstract group in some concrete way. For example, the matrices

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \qquad \mathbf{2} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad , \tag{B.4}$$

with ordinary matrix multiplication, represent the group $\mathbf{Z}_2$. Here, $\mathbf{2} = \mathbf{2}^{-1}$. The objects $\{1, 0\}$, with the multiplication operation identified as addition and $1 + 1$ defined to be 0, also form a representation of $\mathbf{Z}_2$. In this representation, $\mathbf{1} = 0$.

A representation $\mathbf{R}$ is called *irreducible* if for every $x, y \in \mathbf{R}$ there exists an element $z \in \mathbf{R}$ such that $yz = x$. For example, complex 2-vectors with unit norm from an irreducible representation of $SU(2)$. Choosing a complex 2-vector with unit norm at random to associate with the identity, say $(1, 0)$, we see that any other vector in the representation can be obtained via an $SU(2)$ transformation :

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} \alpha & -\beta^* \\ \beta & \alpha^* \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad . \tag{B.5}$$

Associating the vector (α, β) with the transformation necessary here gives us the representation. On the other hand, the representation of $SU(2)$ formed by unitary hermitian 2×2 matrices is *not* irreducible. Indeed, performing unitary $SU(2)$ transformations on the unitary hermitian matrix

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \tag{B.6}$$

will certainly not get us to any other element of our representation. This element forms a representation all by itself. However, once we remove it, we are left with an irreducible representation which any of the three Pauli matrices will be happy to span for us.

This appendix is concerned with the irreducible representations of the spacetime symmetry group, $SO(3, 1)$. We will find it to be a continuous group that relates different inertial frames of reference to each other. Since it is a fundamental symmetry of nature, physical particles and states will form representations of it. This is the foothold which we use to formalize a theory of objects that we cannot directly interact with (on their level), and whose interactions do not always follow our intuition.

Much of the analysis of Section B.2 is based on that found in S. Weinberg [88].

B.1 Relativity

The contemporary understanding of special relativity is based on the assumption that the speed of light *in a vacuum*, c, is the same in all reference frames. To see what this implies, let

us suppose that a beam of light travels in vacuum from spacetime point 0 to spacetime point P. The coordinates of P can be written in a compact notation as x^μ, where μ runs from 0 to 3. x^0 is the time coordinate and x^i, $i = 1 - 3$, are the spatial components of this '4-vector'. Since light goes from 0 to P in vacua,

$$c^2(x^0)^2 = (x^1)^2 + (x^2)^2 + (x^3)^2 \ , \tag{B.7}$$

or[1]

$$c^2(x^0)^2 - (x^1)^2 - (x^2)^2 - (x^3)^2 \equiv x^2 = 0 \ . \tag{B.8}$$

In another reference frame whose origin coincides with this one, the final destination of the light beam will be P'. However, since c is the same in all reference frames, the new coordinates must also satisfy (B.8). Hence the quantity x^2 will be zero either in *all* reference frames or none.

Let us study an example where $x^2 \neq 0$. We consider a laser on a train pointed at a mirror on the ceiling. The train has height h and is moving at a speed v with respect to the ground in the 1-direction. Taking the origin to be the time and place the laser beam is turned on, we study the final destination of the 'front' of the laser beam in two different frames. Observer A seated in the train measures the final destination $x^\mu_A = (2h/c, 0, 0, 0)$.[2] Observer B standing on the ground measures $x^\mu_B = 2h/c\,\gamma(1, v, 0, 0)$, where $\gamma^2 = 1/(1 - v^2/c^2)$. Hence $x^\mu_A \neq x^\mu_B$, but $x^2_A = x^2_B$. It can be shown that this is the case for any process in special relativity. The assumption that the speed of light is the same in all reference frames then implies that the line element

$$ds^2 = -g_{\mu\nu} dx^\mu \, dx^\nu \ , \tag{B.9}$$

where

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \ , \tag{B.10}$$

[1]The component x^2 and the 4-vector square $x^2 = c^2(x^0)^2 - (x^1)^2 - (x^2)^2 - (x^3)^2$ are obviously quite different. However, since the 2-component of a vector is seldom singled out, confusion is rare.

[2]Here, as always, component listing is read $x^\mu = (x^0, x^1, x^2, x^2)$.

is invariant under all changes of frame. In this expression we have implied summation over μ and ν. Any time an index is repeated summation over that index is implied unless explicitly stated otherwise.[3] For simplicity, we have chosen units in which[4]

$$c = 1 \quad . \tag{B.11}$$

With these conventions, we may write $x^2 = g_{\mu\nu}x^\mu x^\nu$. The matrix $g_{\mu\nu}$ is called the *metric tensor*. It defines the geometry of the spacetime we work on. The specific form (B.10) is called the Minkowskii metric and is relevant for the flat spacetime considered in this work.

Physical constraints require our changes in frame to be linear[5], so we can write a general transformation as

$$x^\mu \to x'^\mu = \Lambda^\mu_\nu \, x^\nu \ . \tag{B.12}$$

Λ is our transformation matrix, known in flat space as a Lorentz transform.[6] Invariance of the line element implies

$$g_{\alpha\beta} = \Lambda^\mu_\alpha \Lambda^\nu_\beta g_{\mu\nu} \quad . \tag{B.13}$$

Since the metric behaves in this way, we see that the product of any two vectors x and p, $x \cdot p \equiv g_{\mu\nu}x^\mu p^\nu$, is also invariant under changes of frame.[7]

We can think of Eq.(B.12) as the way an upper index transforms. Defining $x_\mu \equiv g_{\mu\nu}x^\nu$, we see that a vector with a lowered index transforms like $x'_\mu = g_{\mu\alpha}\Lambda^\alpha_\nu x^\nu$. Defining $g^{\mu\nu}$ such that $g^{\mu\alpha}g_{\alpha\nu} = \delta^\mu_\nu$, with $\delta^\mu_\nu = 0$ if $\mu \neq \nu$ and $= 1$ if $\mu = \nu$, we can put this transformation into the form

$$x_\mu \to x'_\mu = (\Lambda^{-1})^\nu_\mu x_\nu \quad . \tag{B.14}$$

[3]This is known as Einstein's 'summation convention'. Note that in order to sum properly, one index must be 'up' and the other 'down'. We will see why this is the case below.

[4] This means if you ask how far away something is, I simply tell you how long it takes light to get there.

[5]If you don't believe me, imagine how a walk to the grocery store then to the post office looks in two reference frames. The grocery store has the two coordinates G_A and G_B while the post office has the two coordinates P_A and P_B. The vector from the grocery store to the post office is by definition $P_A - G_A$ in the first frame and $P_B - G_B$ in the second. If the transformation were nonlinear, then this vector would not transform into its image in the other frame.

[6]By convention, the *first* index of Λ is 'up' while the second index is 'down'. It really should be written $\Lambda^\mu{}_\nu$. Indices on matrices are *always* understood to be this way, unless otherwise emphasized.

[7]Now we see why the notation x^2 was used for the strange object in Eq.(B.8): $g_{\mu\nu}x^\mu x^\nu = x \cdot x = x^2$.

It is now obvious why $g_{\mu\nu}x^\mu p^\nu = x_\mu p^\mu$ is invariant under transformations - one transforms like Λ and the other like Λ^{-1}.

Normal vectors, like position and momentum, naturally have an upstairs index. These are called *contravariant* vectors since they transform in a different way than the axes (rotating the axes one way produces the same effect as rotating vectors the opposite way). Vectors which naturally have a downstairs index, like the derivative, are called *covariant* vectors since they transform in the same way as the axes. To see this, let us examine the derivative operator :

$$\frac{\partial}{\partial x^\mu} f(x) \to \frac{\partial}{\partial x'^\mu} f(x) = \frac{\partial}{\partial x'^\mu} f(\Lambda^{-1}x') = (\Lambda^{-1})^\nu_\mu \frac{\partial}{\partial x^\nu} f(x) \quad . \tag{B.15}$$

Hence, $\partial/\partial x^\mu \equiv \partial_\mu$ does indeed transform opposite to x^μ (or in the same way as the axes).[8]

Tensors behave in exactly the same way. A tensor with n contravariant and m covariant indices transforms like

$$T^{\mu_1\cdots\mu_n}_{\nu_1\cdots\nu_m} \to (\Lambda)^{\mu_1}_{\alpha_1}\cdots(\Lambda)^{\mu_n}_{\alpha_n}(\Lambda^{-1})^{\beta_1}_{\nu_1}\cdots(\Lambda^{-1})^{\beta_m}_{\nu_m} T^{\alpha_1\cdots\alpha_n}_{\beta_1\cdots\beta_m} \quad . \tag{B.16}$$

Often, we wish to emphasize certain symmetries possessed by tensors. This is done via parentheses and brackets in the following way [9]

$$\begin{aligned} T^{(\alpha\beta)\mu_1\cdots\mu_n} &\equiv \frac{1}{2}\left(T^{\alpha\beta\mu_1\cdots\mu_n} + T^{\beta\alpha\mu_1\cdots\mu_n}\right) \\ T^{[\alpha\beta]\mu_1\cdots\mu_n} &\equiv \frac{1}{2}\left(T^{\alpha\beta\mu_1\cdots\mu_n} - T^{\beta\alpha\mu_1\cdots\mu_n}\right) \quad . \end{aligned} \tag{B.17}$$

Invariance under coordinate transformations plays an important role in both General Relativity and the Standard Model. In General Relativity, this invariance plays a central role and severely constrains the form of theory. In the SM, we are concerned mainly with flat space so the constraints imposed are not as harsh, but the theory is still constrained in much the same way as rotational invariance constrains classical mechanics. For example, suppose we wish to

[8] When indices are suppressed in this kind of calculation, confusion as to the specific way they are contracted can arise. The standard convention is that a proper sum goes from covariant to contravariant. For example, $(\Lambda x)^\mu = \Lambda^\mu{}_\nu x^\nu$, but $\Lambda_\mu{}^\nu x_\nu = (\Lambda^{-1})^\nu_\mu x_\nu = (x\Lambda^{-1})_\mu$. Note that in the second example the placement of the indices is crucial to a correct understanding of the product. The first expression is not a proper sum since it goes from contravariant to covariant. In order to write it in an unambiguous way, we had to emphasize the order of the indices.

[9] Sometimes, as we will see below, parentheses also imply tracelessness. It should be clear from the context which is meant.

calculate a vector quantity which depends only on a certain momentum, p. A priori, we must calculate 4 different quantities. However, the requirement of Lorentz covariance forces this object to transform like a vector under Lorentz transformations. Since p is the only vector in the problem, our quantity must be proportional to p. We need only calculate the constant of proportionality (which will in general depend on all invariants in the problem, i.e. masses and p^2). With two indices, the problem is a little more complicated. Suppose we have a two-index tensor which depends on no momenta. The argument above would lead to 0 for an answer since there is 'nothing to carry the indices'. What this means is that there is no reference to frame in this problem. Our quantity must be frame-independent. However, this does not imply that it is zero since the tensor $g^{\mu\nu}$ satisfies our requirement, as exemplified by (B.13). In this case, Lorentz invariance forces our tensor to be proportional to $g^{\mu\nu}$.

There is only one other tensor-like object which is invariant under Lorentz transformations. The Levi-Cevita[10] tensor is defined as a 4-index tensor which is completely antisymmetric in all four indices. Since there are only four possible values for each of the indices, the Levi-Cevita tensor has only one independent component. This component is chosen to be 1, i.e.

$$\epsilon^{0123} = +1 \quad . \tag{B.18}$$

All other components are obtained from the requirement of total antisymmetry. To see that the Levi-Cevita tensor is invariant, we transform it :

$$\epsilon^{\mu_1\mu_2\mu_3\mu_4} \rightarrow \Lambda^{\mu_1}_{\nu_1}\Lambda^{\mu_2}_{\nu_2}\Lambda^{\mu_3}_{\nu_3}\Lambda^{\mu_4}_{\nu_4}\epsilon^{\nu_1\nu_2\nu_3\nu_4} = \det(\Lambda)\epsilon^{\mu_1\mu_2\mu_3\mu_4} \quad , \tag{B.19}$$

where the last equality comes from the definition of the determinant,

$$\det(\Lambda) \equiv \frac{1}{4!}\epsilon_{\mu_1\mu_2\mu_3\mu_4}\epsilon^{\nu_1\nu_2\nu_3\nu_4}\Lambda^{\mu_1}_{\nu_1}\Lambda^{\mu_2}_{\nu_2}\Lambda^{\mu_3}_{\nu_3}\Lambda^{\mu_4}_{\nu_4} \quad . \tag{B.20}$$

(B.13) implies that the determinant of Λ is either $+1$ or -1. The structure of the Lorentz group is such that all transformations with determinant -1 can be obtained from determinant $+1$ transformations and the two discreet operations of parity and time-reversal. We see from (B.19) that these two operations change the sign of the Levi-Cevita symbol. This fact causes ϵ to play an important role in the spin structure of hadrons in QCD.

[10] Cevita is pronounced 'chävētă', while Levi is like 'Drove my Chevy to the *levi...*'.

Up to now, our analysis has been quite general. None of the above manipulations have required the use of the Minkowskii metric (B.10). At this point, we specialize to flat-space and discuss the Lorentz transformations explicitly. (B.13) implies that Λ is orthogonal, i.e. $\Lambda^{\mathrm{T}}\Lambda = 1$. In view of the above discussion on the operations of parity and time-reversal, we will consider only those transformations with $\det\Lambda = +1$. The group of orthogonal transformations in Minkowskii space with unit determinant is called $SO(3,1)$ (the *special orthogonal* group in 3 spacelike and 1 timelike dimensions). Any real orthogonal matrix can be written as e^M, with $M = -M^{\mathrm{T}}$ a real skew-symmetric matrix. A general skew-symmetric real 4×4 matrix has six independent components, and hence the space of these objects can be spanned by six linearly independent matrices. The matrices we choose to form this basis are called the *generators* of $SO(3,1)$ since they *generate* the group. The algebra satisfied by the generators is called the *Lie*[11] algebra of the group. Any member of the group is uniquely determined by the six coefficients of the generators. The fact that there is a one-to-one mapping between these coefficients and the elements of the group means that the coefficients form a representation of the group, the *adjoint* representation.

The six generators of $SO(3,1)$ can be conveniently placed into an antisymmetric 4-tensor, i.e.[12]

$$M^{\alpha}_{\beta} = \frac{1}{2}\theta_{\mu\nu}(J^{\mu\nu})^{\alpha}_{\beta} \quad . \tag{B.21}$$

In principle, we are now free to choose whatever form we like for the generators $J^{\mu\nu}$, as long as we end up with six linearly independent skew-symmetric real matrices. The most simple choice is also the most natural :

$$(J^{\mu\nu})_{\alpha\beta} = \delta^{\mu}_{\alpha}\delta^{\nu}_{\beta} - \delta^{\nu}_{\alpha}\delta^{\mu}_{\beta} \quad . \tag{B.22}$$

Working out the commutator of the generators,

$$\left[J^{\mu\nu}, J^{\alpha\beta}\right] = g^{\mu\beta}J^{\nu\alpha} - g^{\nu\beta}J^{\mu\alpha} - g^{\mu\alpha}J^{\nu\beta} + g^{\nu\alpha}J^{\mu\beta} \quad , \tag{B.23}$$

we find that $SO(3,1)$ is a non-abelian group. This means that we must be careful with the order of transformations. In particular, θ's for successive transformations do *not* add.

[11] As in '... the *lie* of the stone.'

[12] The factor of 1/2 corrects the overcounting of the implied sum on $\mu\nu$.

Since three-dimensional rotations are among the possible Lorentz transformations, our generators should include those for ordinary $SO(3)$ rotations. This is indeed the case, as we can see by examining any generator with both μ and ν spacelike. For example,

$$(J^{12})^{\alpha}_{\beta} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad . \tag{B.24}$$

It is easy to see that this matrix has four different powers, only two of which are linearly independent. Hence the $SO(3,1)$ rotation it generates is

$$(e^{\theta J^{12}})^{\alpha}_{\beta} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta & 0 \\ 0 & \sin\theta & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad , \tag{B.25}$$

which is a rotation of θ radians about the 3-axis. Hence we identify $J^{12} = L^3$. For general rotations, the identification[13] $\epsilon^{ijk} J^{ij} = 2L^k$ is appropriate. (B.23) then implies

$$\left[L^i, L^j\right] = \epsilon^{ijk} L^k \quad , \tag{B.26}$$

as expected for generators of $SO(3)$ rotations. Note that pure rotations form a subgroup of $SO(3,1)$, that is $SO(3) \subset SO(3,1)$.

The other three generators have one time index and one spatial index. These objects will generate Lorentz boosts, as can be seen by examining J^{01} :

$$(J^{01})^{\alpha}_{\beta} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad . \tag{B.27}$$

[13] It is a standard convention to use Greek letters for general indices and Roman letters when one wants to specify a spatial index. The ϵ tensor here is the Levi-Cevita tensor for $SO(3)$, with the convention $\epsilon^{123} = +1$. Since the metric relevant to $SO(3)$ is trivial, we make no distinction between contravariant and covariant indices. However, a few well-placed minus signs are required when going from a general expression to one which specifies spatial indices since $g_{ij} = -\delta_{ij}$.

Once again, the powers of this matrix are cyclic so exponentiation is trivial :

$$(e^{\theta J^{01}})^{\alpha}_{\beta} = \begin{pmatrix} \cosh\theta & \sinh\theta & 0 & 0 \\ \sinh\theta & \cosh\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad . \tag{B.28}$$

Under the action of this transformation,[14]

$$\begin{aligned} t &\rightarrow t\cosh\theta + x\sinh\theta \\ x &\rightarrow x\cosh\theta + t\sinh\theta \\ y &\rightarrow y \\ z &\rightarrow z \ . \end{aligned} \tag{B.29}$$

It is obvious from the infinitesimal form of this transformation that we have given the vector x a velocity θ along the 1-direction. For asymptotically large values of θ, x and t approach the same value $(t+x)\cosh\theta$, making the vector x approximately lightlike.[15] Hence asymptotically large θ boost the vector x approximately to the speed of light. This argument promotes the identification $v = \tanh\theta$. We can now eliminate θ in favor of v to obtain

$$\begin{aligned} t &\rightarrow \gamma(t + vx) \\ x &\rightarrow \gamma(x + vt) \\ y &\rightarrow y \\ z &\rightarrow z \ , \end{aligned} \tag{B.30}$$

the new coordinates of a vector x which has been endowed with a speed v along the 1-direction. Again, we have used $\gamma \equiv (1-v^2)^{-1/2}$. This transformation reduces to the familiar Galilean transformation if we reinstate c and take $v \ll c$ (unless we take xv/c^2 comparable to t).

The transformations $J^{0i} \equiv K^i$ do not form a subgroup of $SO(3,1)$ since (B.23) implies

$$\left[K^i, K^j\right] = -\epsilon^{ijk} L^k \quad . \tag{B.31}$$

[14]We will use the notation $x^0 = t$, $x^1 = x$, $x^2 = y$, and $x^3 = z$ interchangeably. Confusion between $x^1 = x$ and $x^\mu = x$ should be rare.

[15]Of course, this vector does not *actually* become lightlike since its invariant square is invariant. However, in our new frame x^2 will in fact be much smaller than the square of its t and x coordinates. This makes the vector look lightlike in this frame.

The commutator of L with K is

$$\left[L^i, K^j\right] = \epsilon^{ijk} K^k \ , \tag{B.32}$$

which closes the algebra of the group.

B.2 Unitary Representations of $SO(3,1)$

In this section, we will study how the Lorentz group acts on fields. The first thing we must do is expand our group to include translations. Under a translation, the vector x is shifted to $x + a$. This transformation is *not* linear since $(x + y) + a \neq (x + a) + (y + a)$. However, a *function* of the vector x will transform linearly under a translation :

$$T(a)f(x) = f(x - a) \ . \tag{B.33}$$

Note that we have implicitly defined the transformation $T(a)$ in such a way that it shifts the *function* it acts on forward an amount a rather than the axes. Hence the value of the unshifted function at x is the same as the value of the shifted function at $x - a$. This extension of the Lorentz group is called the Poincaré[16] group, and its representations form the basis for quantum field theory.

Since we do not want to alter the normalizations of our states through these transformations, we will be concerned with *unitary* representations of the Poincaré group. Elements of these representations will be related to the coordinate transformations Λ and a, i.e. $U = U(\Lambda, a)$. The group multiplication rule,

$$U(\Lambda_2, a_2)U(\Lambda_1, a_1) = U(\Lambda_2\Lambda_1, a_2 + \Lambda_2 a_1) \ , \tag{B.34}$$

is governed by these transformations as well. This rule implies $U^{-1}(\Lambda, a) = U^\dagger(\Lambda, a) = U(\Lambda^{-1}, -\Lambda^{-1}a)$. U is unitary, so it can be represented as the exponential of an anti-hermitian matrix. The translations add four generators to the Lorentz group, one for each dimension of spacetime, so we have the form

$$U(1 + \frac{1}{2}\theta_{\alpha\beta}J^{\alpha\beta}, \epsilon^\mu) = 1 - i\frac{1}{2}\theta_{\alpha\beta}\hat{J}^{\alpha\beta} + i\epsilon^\mu \hat{P}_\mu \tag{B.35}$$

[16] like *s'il vous plait*

for arbitrary infinitesimal ϵ and θ. Note that J are the matrices from the last section; they act on physical spacetime vectors. $\hat{J}$ and $\hat{P}$ are hermitian operators that act on the space in which the fields we are concerned with live. Consider the transformation

$$U(\Lambda,a)U(1+\frac{1}{2}\theta J,\epsilon)U^{-1}(\Lambda,a)=U(1+\frac{1}{2}\Lambda\theta J\Lambda^{-1},\Lambda\epsilon-\frac{1}{2}\Lambda\theta J\Lambda^{-1}a)\ . \tag{B.36}$$

Expanding in θ and ϵ, substituting the explicit expression for J, and noting that each component of θ and ϵ may be varied independently, we arrive at

$$U(\Lambda,a)\hat{J}^{\mu\nu}U^{-1}(\Lambda,a) = (\Lambda^{-1})^{\mu}_{\alpha}(\Lambda^{-1})^{\nu}_{\beta}\left(\hat{J}^{\alpha\beta}-a^{\alpha}\hat{P}^{\beta}+a^{\beta}\hat{P}^{\alpha}\right)\ , \tag{B.37}$$

$$U(\Lambda,a)\hat{P}^{\mu}U^{-1}(\Lambda,a) = (\Lambda^{-1})^{\mu}_{\nu}\hat{P}^{\nu}\ . \tag{B.38}$$

Equation (B.37) states that for pure rotations ($a=0$), transforming the operator $\hat{J}$ is the same as performing the *inverse* transformation on the coordinate indices of $\hat{J}$. The inverse transformation appears here for the same reason that $x-a$ appears as the argument of $T(a)f(x)$. When translations are included, we see that $\hat{J}$ transforms in exactly the way we expect an angular momentum to transform under a change of origin. (B.38) tells us that the operator $\hat{P}$ transforms as a vector under Lorentz transformations, but is invariant under translation. We can obtain the Lie algebra of this group by considering (B.37) and (B.38) for infinitesimal transformations Λ and a. Performing the same manipulations as above, we see that

$$-i\left[\hat{J}^{\mu\nu},\hat{J}^{\alpha\beta}\right] = g^{\mu\beta}\hat{J}^{\nu\alpha}-g^{\nu\beta}\hat{J}^{\mu\alpha}-g^{\mu\alpha}\hat{J}^{\nu\beta}+g^{\nu\alpha}\hat{J}^{\mu\beta}\ , \tag{B.39}$$

$$-i\left[\hat{P}^{\mu},\hat{J}^{\alpha\beta}\right] = g^{\mu\alpha}\hat{P}^{\beta}-g^{\mu\beta}\hat{P}^{\alpha}\ , \tag{B.40}$$

$$-i\left[\hat{P}^{\mu},\hat{P}^{\nu}\right] = 0\ . \tag{B.41}$$

The last identity implies that the subgroup of pure translations is an abelian group. The algebra of the Lorentz subgroup is the same as we found above, as it should be. However, it must be emphasized that there we were discussing 4×4 matrices rather than the infinite dimensional quantum mechanical operators here.

$\hat{P}$ and $\hat{J}$ have a very concrete physical meaning in quantum mechanics. To see this, consider a system which is translationally invariant. Since the phase of an isolated[17] quantum

[17]If it wasn't isolated, it could not be translationally invariant. If state vectors interact, we must translate them all simultaneously. This is exactly the same as in classical mechanics, where we must consider the *total* momentum of a system to obtain conservation.

mechanical system is not observable, the state may change at most by the overall phase

$$T(a)\,|S\rangle = e^{i\delta(a)}\,|S\rangle \quad , \tag{B.42}$$

where $\delta(a)$ is some real function of a. In light of the above discussion, we see that $\delta(a)$ should be identified with $\hat{P}\cdot a$. This implies that $|S\rangle$ is an eigenstate of $\hat{P}$. According to (B.41), this eigenvalue is conserved under translation. In classical mechanics, the conserved quantity related to translation invariance is the momentum.[18] Hence, we identify the conserved quantity here with the momentum of the system.[19] Since the momentum of the system is an eigenvalue of $\hat{P}$, we call $\hat{P}$ the momentum operator.

Attempting to apply the same analysis to the operator $\hat{J}$, we run into a little snag. The fact that the $\hat{J}$'s do not commute with each other implies that $|S\rangle$ cannot simultaneously be an eigenstate of all of them unless most of its eigenvalues are zero.[20] When discussing $\hat{J}$, it is convenient to define $\hat{L}^i \equiv 1/2\epsilon^{ijk}\hat{J}^{jk}$ and $\hat{K}^i \equiv J^{0i}$ in analogy with the last section. The commutation relations (B.39) imply

$$[\hat{L}^i, \hat{L}^j] = i\epsilon^{ijk}\hat{L}^k \quad , \tag{B.43}$$

$$[\hat{K}^i, \hat{K}^j] = -i\epsilon^{ijk}\hat{L}^k \quad , \tag{B.44}$$

$$[\hat{L}^i, \hat{K}^j] = i\epsilon^{ijk}\hat{K}^k \quad . \tag{B.45}$$

These relations have the interesting consequence that the operators

$$\begin{aligned} \hat{A}^i &= \frac{1}{2}\left(\hat{L}^i + i\hat{K}^i\right) \\ \hat{B}^i &= \frac{1}{2}\left(\hat{L}^i - i\hat{K}^i\right) \end{aligned} \tag{B.46}$$

[18] Remember, momentum is conserved in the absence of external forces, i.e. forces which do not belong to the system we consider. These forces would break translational invariance, and thus momentum conservation.

[19] This really implies that the momentum of the system is *proportional* to the eigenvalue. If the momentum of the system is p and the eigenvalue is k, we say $p = \hbar k$. p has the dimensions of momentum (Energy/Speed=Energy since we work in units where speed is dimensionless) and k has the dimensions of an inverse time (since the exponential of $\hat{P}\cdot a$ makes sense; remember that since speed is dimensionless length and time are the same unit), so $\hbar$ is Energy×Time. Once again, we can adjust our units so that $\hbar = 1$. In other units, we can actually measure its value; in SI units, $\hbar = 1.05457266(63) \times 10^{-34}\, J\cdot s$.

[20] We can see this as follows. Suppose $\hat{J}^{\mu\nu}|S\rangle = j^{\mu\nu}|S\rangle$. Now, $[\hat{J}^{\mu\nu}, \hat{J}^{\lambda\rho}]|S\rangle = 0$ since the numbers $j^{\mu\nu}$ and $j^{\lambda\rho}$ commute. This implies that the commutator has eigenvalue zero.

generate commuting subgroups of $SO(3,1)$,

$$\begin{aligned} [\hat{A}^i, \hat{A}^j] &= i\epsilon^{ijk}\hat{A}^k \\ [\hat{B}^i, \hat{B}^j] &= i\epsilon^{ijk}\hat{B}^k \\ [\hat{A}^i, \hat{B}^j] &= 0 \ . \end{aligned} \tag{B.47}$$

Furthermore, the commutation relations satisfied by these generators identify the subgroups with $SU(2)$. Hence the group of Lorentz transformations, $SO(3,1)$, is seen to be nothing more than the direct product of two commuting $SU(2)$'s. This will be seen even more explicitly when we study the fundamental representation of $SL(2,C)$ and its relation to $SO(3,1)$ in the next section.

In non-relativistic physics, we concern ourselves with time evolution and consider a quantity to be *conserved* if it does not change with time. The Schrödinger approach to quantum mechanics considers state vectors that change with time,

$$|S(t)\rangle = e^{-it\hat{H}}\,|S(0)\rangle \quad . \tag{B.48}$$

The time dependence is governed by the time-translation operator, or *Hamiltonian*, $\hat{H} \equiv \hat{P}^0$. The action of an operator on a state will change with time because the state itself changes with time. In any physical matrix element, we can use this relation to display the time dependence explicitly :

$$\langle S'(t)|\,\mathcal{O}\,|S(t)\rangle = \langle S'(0)|\,e^{it\hat{H}}\mathcal{O}e^{-it\hat{H}}\,|S(0)\rangle \quad . \tag{B.49}$$

This promotes the identification $\mathcal{O}(t) \equiv e^{it\hat{H}}\mathcal{O}e^{-it\hat{H}}$, which puts the time-dependence of the matrix element into the operator. This is Heisenberg's view of quantum mechanics. Here, the action of an operator on a state will change with time because the operator changes with time. If the operator we are concerned with commutes with the Hamiltonian, it will not depend on time. Hence *all* of its matrix elements will be conserved. Looking at Eqs.(B.40) and (B.41), we see that $\hat{P}^i$ and $\hat{L}^i$ are conserved operators.[21] Since $\hat{L}^i$ is a conserved operator associated with spatial rotation, we identify it with the angular momentum operator. If a system is invariant

[21] In non-relativistic quantum mechanics, one usually has some sort of space-dependent potential that completely destroys translation invariance. These potentials act as static momentum sources which effectively generate nonzero commutators of the Hamiltonian with momentum and angular momentum operators.

under rotation about a certain axis, say $\hat{n}$, the argument given above for $\hat{P}$ implies that it is and eigenstate of $\hat{n} \cdot \hat{L}$. On the other hand, the commutator

$$[\hat{H}, \hat{K}^i] = i\hat{P}^i \tag{B.50}$$

implies that $\hat{K}^i$ will depend on time.

In relativistic quantum mechanics, we are concerned mostly with operators; the external states can all be formed from the vacuum by the action of various operators. For this reason, it is Heisenberg's approach that is most useful here. The fact that time and space mix under Poincaré transformations makes it natural for us to consider them together. In a translationally invariant theory, the space dependence of the external states can be removed in exactly the same way as the time dependence. For any spatial 4-vector x^μ and any quantum operator $\mathcal{O}$, we have the operator identity

$$\mathcal{O}(x) = e^{ix\cdot\hat{P}}\mathcal{O}(0)e^{-ix\cdot\hat{P}} \quad . \tag{B.51}$$

Operators that commute with all of the $\hat{P}^\mu$ do not depend on coordinates at all.[22]

We now turn to a classification of the states in our theory. Suppose we have a state which represents an electron with momentum p^μ and spin polarization s^μ. Poincaré transformations can change the electron's momentum and spin, but certainly cannot change the fact that it is an electron. This means that general properties attributed to electrons must be associated with operators that commute with *all* of the generators of the group. Operators with this property are called *Casimirs* of the group. It is easily shown that the mass operator[23] $\hat{P} \cdot \hat{P} = \hat{P}^2$ is a casimir of the Poincaré group. The other casimir has to do with spin. We know that an electron has spin-1/2 regardless of where it is or how we rotate it. There must be an operator that expresses that. By inspection, one can see that the Pauli-Ljubanskii vector

$$\hat{W}^\mu \equiv -\frac{1}{2}\epsilon^{\mu\nu\alpha\beta}\hat{P}_\nu\hat{J}_{\alpha\beta} \tag{B.52}$$

is exactly what is needed. Under parity, $\hat{W}^\mu$ behaves exactly as a spin should since ϵ changes sign. $\hat{W}$ is also orthogonal to $\hat{P}$ [24] and reduces to a spatial angular momentum operator in

[22] This is what we mean by 'translationally invariant'!

[23] Remember that in relativity the invariant square of the momentum of a system is the square of the mass of that system.

[24] Under the influence of the Levi-Cevita symbol, the operators $\hat{P}$ and $\hat{J}$ commute.

the rest frame, $\hat{P} = (M, 0, 0, 0)$. The commutation relations of this vector with the generators of the Poincaré group are identical with those of $\hat{P}^\mu$:

$$\begin{aligned} -i[\hat{W}^\mu, \hat{P}^\nu] &= 0 \\ -i[\hat{W}^\mu, \hat{J}^{\alpha\beta}] &= g^{\mu\alpha}\hat{W}^\beta - g^{\mu\beta}\hat{W}^\alpha \quad . \end{aligned} \tag{B.53}$$

This makes it obvious that its square

$$\hat{W}^2 = -\frac{1}{2}\hat{J}_{\mu\nu}\hat{J}^{\mu\nu}\hat{P}^2 + \hat{J}_{\lambda\rho}\hat{J}^{\sigma\rho}\hat{P}^\lambda\hat{P}_\sigma \tag{B.54}$$

is a casimir.[25]

To classify the irreducible representations of the Poincaré group, we appeal to the idea of Wigner's 'little group' [89]. To begin with, we diagonalize $\hat{P}^\mu$:

$$\hat{P}^\mu |p\rangle = p^\mu |p\rangle \quad . \tag{B.55}$$

Since p^μ is a vector, it will transform under Lorentz transformations as

$$p^\mu \to \Lambda^\mu_\nu p^\nu \quad . \tag{B.56}$$

We can use this to transform our state to a convenient frame with a representative momentum $\tilde{p}^\mu$. In this frame, there will be residual transformations which do not change $\tilde{p}^\mu$, but may act on our state. This group is called the little group associated with the reference momentum $\tilde{p}^\mu$. The representations of the little group will tell us how many different states of momentum $\tilde{p}^\mu$ are related to our state through the Poincaré group. Since $\hat{W}$ commutes with $\hat{P}$, we can diagonalize one of its components to break the degeneracy associated with this subgroup. We will consider two examples of reference momentum $\tilde{p}^\mu$ since these are the only two which actually occur in nature (to our knowledge).

As our first example, we take states for which $p^0 > 0$ and $p^2 = M^2 > 0$. A look at the Lorentz transformations tells us that if $p^0 > 0$ in one frame, it will be in all frames, provided $p^2 \geq 0$.[26] Furthermore, there is always a *rest frame*, defined by $\tilde{p}^\mu = (M, 0, 0, 0)$. The little

[25] This is obvious once one knows that $\hat{W}$ is a vector that commutes with $\hat{P}$. Any Lorentz scalar will commute with the rotation generators. It is the fact the $\hat{W}$ is translation invariant that makes it special.

[26] We consider here only the proper Lorentz group. Parity and time-reversal can be taken into account separately later.

group associated with this momentum is the group of spatial rotations, $SO(3)$. In this frame, we have $\hat{W}^0 = 0$ and $\hat{W}^i = M\hat{L}_i$. Since $\hat{L}^2 = \hat{L}_i\hat{L}_i$ commutes with $\hat{L}_i$, we can simultaneously diagonalize $\hat{L}^2$ and one of the $\hat{L}_i$:

$$\begin{aligned} \hat{L}^2 \left|\ell m\right\rangle &= \ell(\ell+1)\left|\ell m\right\rangle \\ \hat{L}_3 \left|\ell m\right\rangle &= m\left|\ell m\right\rangle \quad . \end{aligned} \tag{B.57}$$

Since $\hat{L}^2$ is positive definite, we can take $\ell \geq 0$. The reason for the special form of its eigenvalue will become clear shortly. The hermiticity of $\hat{L}^2$ and $\hat{L}_3$ allows us to choose orthonormal eigenstates. Defining

$$\hat{L}_\pm \equiv \hat{L}_1 \pm i\hat{L}_2 \quad , \tag{B.58}$$

we have

$$[\hat{L}_3, \hat{L}_\pm] = \pm\hat{L}_\pm \quad . \tag{B.59}$$

Hence

$$\hat{L}_3\hat{L}_\pm \left|\ell m\right\rangle = (m \pm 1)\hat{L}_\pm \left|\ell m\right\rangle \quad , \tag{B.60}$$

making $\hat{L}_\pm$ the *raising* and *lowering* operators associated with $\hat{L}_3$. If this process were to go on indefinitely, we would at some point have a state for which $m > \ell$. However,

$$\ell(\ell+1) = \left\langle \ell m\right| \hat{L}^2 \left|\ell m\right\rangle = \left\langle \ell m\right| \hat{L}_1^2 + \hat{L}_2^2 \left|\ell m\right\rangle + m^2 \tag{B.61}$$

implies that $m^2 \leq \ell(\ell+1)$. This means that there must be some state $\left|\ell m_\pm\right\rangle$ that truncates the process,

$$\hat{L}_\pm \left|\ell m_\pm\right\rangle = 0 \quad . \tag{B.62}$$

Writing $\hat{L}^2 = \hat{L}_\pm\hat{L}_\mp + \hat{L}_3^2 \mp \hat{L}_3$, it is easy to show that $m_\pm = \pm\ell$. Starting with the 'top' state, $\left|\ell\ell\right\rangle$, we can apply $\hat{L}_-$ to successively lower m by one unit each time. At some point, we must reach the 'bottom' state $\left|\ell(-\ell)\right\rangle$ else suffer the consequences of inconsistency. Since we cannot apply L_- a fraction of a time, this implies that $\ell - n = -\ell$ for some whole number n. Hence ℓ must be either integer or half integer.[27] For each ℓ, we can define all of the states in our representation from the top state via successive application of $\hat{L}_-$. The appropriate normalization is

$$\hat{L}_\pm \left|\ell m\right\rangle = \sqrt{(\ell \mp m)(\ell \pm m + 1)}\left|\ell \;\; m \pm 1\right\rangle \quad , \tag{B.63}$$

[27] This is the reason we assumed the form $\ell(\ell+1)$ for the eigenvalue of $\hat{L}^2$.

as can be obtained from the requirement that $\ell(\ell+1) = \langle \ell m | \hat{L}^2 | \ell m \rangle$.[28]

In the rest frame of a massive system, we have shown that the spectrum of $\hat{W}^2$ is $(-M^2\ell(\ell+1))$, where $\ell \geq 0$ is either integer or half-integer. Since this object is a casimir of the Poincaré group, different values of ℓ correspond to different irreducible representations. For fixed ℓ, there are $2\ell+1$ different states in the representation. Under the action of the rotation group, these states behave as though they have different angular momentum, so we say that in the rest frame the state has angular momentum (spin) ℓ. The spin vector is defined by $\hat{S}^\mu \equiv \hat{W}^\mu/M$. States in other frames can be obtained by boosting the rest frame states.

The other kind of states which appear in nature have $p^2 = 0$, $p^0 > 0$. In this case, we take $\tilde{p}^\mu = (\omega, 0, 0, \omega)$ for our reference momentum. The little group for this momentum is generated by the components of $\hat{W}$:

$$\begin{aligned} \hat{W}^0 &= \omega \hat{L}_3 \\ \hat{W}^1 &= \omega(\hat{L}_1 + \hat{K}_2) \\ \hat{W}^2 &= \omega(\hat{L}_2 - \hat{K}_1) \\ \hat{W}^3 &= \omega \hat{L}_3 \ . \end{aligned} \tag{B.64}$$

These generators satisfy the commutation relations

$$-i[\hat{W}^1, \hat{W}^3] = -\omega \hat{W}^2$$

$$-i[\hat{W}^2, \hat{W}^3] = \omega \hat{W}^1 \tag{B.65}$$

$$-i[\hat{W}^1, \hat{W}^2] = 0 \ , \tag{B.66}$$

so we can simultaneously diagonalize $\hat{W}^1$ and $\hat{W}^2$. However, using the commutation relations, one can show that the existence of one state with finite eigenvalues w_1 and w_2 for $\hat{W}^1$ and $\hat{W}^2$ implies the existence of a continuous infinity of such states :

$$e^{i\theta \hat{L}_3} |w_1, w_2\rangle = |w_1 \cos\theta + w_2 \sin\theta, w_2 \cos\theta - w_1 \sin\theta\rangle \ . \tag{B.67}$$

Since we observe no massless states with a continuous degree of freedom like this, we have no choice but to conclude that $\hat{W}^1$ and $\hat{W}^2$ annihilate our states. In this subspace, $\hat{W}^1$ and $\hat{W}^2$ commute with $\hat{W}^3$ so we are free to diagonalize it. This leaves us with no generators to

[28] We also use the fact that $(\hat{L}_\pm |\ell m\rangle)^\dagger = \langle \ell m | \hat{L}_\mp$.

connect different eigenstates of $\hat{W}^3$. If $\hat{W}^3$ has more than one eigenvalue, these states must be in different irreducible representations of the Poincaré group since they are not linked by any of its generators.[29]

Massless states of 3-momentum $\vec{p}$ are characterized by the action of the *helicity operator*, $\vec{p}\cdot\hat{\vec{L}}/p^0$. The eigenvalue of this operator, λ, is called the *helicity* of the state. Written in this way, the helicity of a system is invariant under the action of the Poincaré group. Although the proper Lorentz group cannot connect massless states with different helicity, the action of parity relates helicity $+\lambda$ to helicity $-\lambda$.[30] Topological considerations require[31]

$$e^{4\pi i\hat{L}_3} = e^{4\pi i\lambda} = 1 \quad , \tag{B.68}$$

or λ is either integer or half-integer. The fact that λ is the eigenvalue of an angular momentum operator tempts us to call it the spin projection of the system. However, since the square of the Pauli-Ljubanskii vector is 0 regardless of the value of λ, we must be careful with such terms.

Explicit forms for the quantum mechanical operators $\hat{P}$ and $\hat{J}$ will in general depend on the specific theory. These operators can be obtained from the lagrangian using general techniques outlined in Section D.2.

B.3 The Spinor Representation of $SO(3,1)$

In this section, we consider the smallest representation of $SO(3,1)$. From ordinary quantum mechanics, we know that the group $SO(3)$ is locally isomorphic to $SU(2)$ (it satisfies the same Lie algebra). This led us to the fact that the fundamental representation of $SO(3)$ can be realized in terms of quantum mechanical 2-spinors. Now, we are faced with the problem of

[29] The other generators are of no use to us here since they will change the momentum of the state.

[30] This is because parity changes the sign of $\vec{p}$ while leaving $\hat{\vec{L}}$ invariant.

[31] The Lorentz group is not simply connected, so the identification of a 2π rotation with the identity is misleading. However, going around the group *twice* will always get us back to the same element. One way to see this is through the rotation subgroup, $SO(3)$. Rotation through an angle θ off the z-axis and ϕ around it (which is represented (θ,ϕ)) is physically identical to $(2\pi-\theta,\pi+\phi)$. These are two *distinct* group elements which are *identified* in this representation. This degeneracy is only twofold, so going around twice gets one back to the same group element.

finding the fundamental representation of the somewhat larger group, $SO(3,1)$. A simple way of going from a certain group to a larger one is to relax some of the constraints defining the smaller group. Relaxing the determinant condition in $SU(2)$ only yields a trivial phase which commutes with everything. One look at Eq.(B.39) tells us that this is not good enough. Let us see what happens if we relax the uniterity condition.

The group of 2×2 complex matrices with unit determinant is a representation of $SL(2,C)$, the special linear group in two complex dimensions. A general element of $SL(2,C)$ has four complex entries with one complex condition. Hence it is composed of six independent elements. This certainly is a good sign. Let us take ξ^α to be a two component object which transforms under $SL(2,C)$ as

$$\xi^\alpha \to (T\xi)^\alpha \ , \tag{B.69}$$

and η_α to be one which transforms as

$$\eta_\alpha \to (\eta T^{-1})_\alpha \ . \tag{B.70}$$

Obviously, because of the way we have defined ξ and η, $\eta_\alpha \xi^\alpha$ is invariant under $SL(2,C)$. Our experience with $SO(3,1)$ has taught us that conditions on matrices lead to invariants of the group. There, we saw that the condition $\det\Lambda = +1$ led to the invariance of the Levi-Cevita symbol. Here, we define $\epsilon_{\alpha\beta}$ to be the totally antisymmetric tensor of $SL(2,C)$. We use the convention $\epsilon_{12} = +1$. Since ϵ is invariant, the product $\epsilon_{\alpha\beta}\xi^\alpha\eta^\beta$ is also invariant. This prompts us to identify

$$\eta_\alpha \equiv \epsilon_{\alpha\beta}\eta^\beta \ , \tag{B.71}$$

making ϵ the metric tensor in this space. Note that our metric is *anti*symmetric. This means that we must be extremely careful with the order of indices and the proper way to sum here. For example, $\epsilon_{\alpha\beta}\eta^\alpha = -\eta_\beta \neq \eta_\beta$. We can define a raising operator such that

$$\epsilon^{\alpha\beta}\epsilon_{\beta\gamma} = \delta^\alpha_\gamma \ . \tag{B.72}$$

Note that this implies that $\epsilon^{\alpha\beta} = -\epsilon_{\alpha\beta}$.

The complex conjugate of ξ^α transforms under $SL(2,C)$ like $T^* \neq T$. Hence it forms a separate space of transformations. Denoting $(\xi^\alpha)^* \equiv \xi^{\dot\alpha}$, we can play the whole game over again. The index $\dot\alpha$ is called a 'dotted' index and lives in a different space than α. Since the two indices transform in completely separate ways, there is no metric which connects them.

As with any group representation, products of spinors give us higher representations. For example, the product

$$\zeta^{\alpha\beta} = \xi^{\alpha}\eta^{\beta} \tag{B.73}$$

is a spinor tensor of rank two (or a *bispinor*). This representation is reducible, as we can see by separating it into two independent pieces :

$$\zeta^{\alpha\beta} = \frac{1}{2}\left(\zeta^{\alpha\beta} + \zeta^{\beta\alpha}\right) + \frac{1}{2}\left(\zeta^{\alpha\beta} - \zeta^{\beta\alpha}\right) \quad . \tag{B.74}$$

The second term is proportional to the metric $\epsilon^{\alpha\beta}$ since it is antisymmetric in α and β. Hence it is invariant under $SL(2,C)$. The first term is symmetric in α and β. We denote this as $\zeta^{(\alpha\beta)}$. Since it is symmetric, it only has three independent components. Note that the total number of components of $\zeta^{\alpha\beta}$, 4, is conserved in the process of breaking it up into irreducible representations. This is a good check that we have done the separation correctly. This equation may be written as $\mathbf{2} \times \mathbf{2} = \mathbf{3} + \mathbf{1}$ since we began with the direct product of two spinor representations ($\mathbf{2}$) and ended with the sum of a singlet ($\mathbf{1}$) and a triplet ($\mathbf{3}$) representation.

In general, a spinor tensor with $2s$ undotted indices may be completely symmetrized by a systematic removal of all of its 'traces' (contractions with the metric tensor). The remaining object transforms as spin s and the traces we removed transform as spin $s-1, s-2, \ldots, 0$ or $1/2$, depending on whether $2s$ is even or odd. These statements can be used as a definition of 'spin' if one so chooses. The motivation comes from the eigenstates of the angular momentum operator, which we have not yet discussed in this context. However, the spectrum of eigenvalues of J^2 and J_z depends only on the commutation relations of the group $SO(3,1)$. Since these relations are the same as those of $SL(2,C)$, as we will show below, the angular momentum operators in our spinor space will have the same spectrum we discussed above. An irreducible representation of $SL(2,C)$ with three independent components, like $\zeta^{(\alpha\beta)}$, will have eigenvalues of $\hat{J}_z = 0, \pm 1$ and $\hat{J}^2 = 2$. Thus is behaves like it has angular momentum, or spin, 1.

Mixed combinations of spinors are of a completely another form entirely. Since dotted and undotted indices live in two different spaces, it makes no sense to symmetrize or antisymmetrize them. The product of a spinor ($\mathbf{2}$) and an antispinor ($\overline{\mathbf{2}}$) representation is itself irreducible

($\mathbf{2} \times \overline{\mathbf{2}} = \mathbf{4}$). A mixed bispinor will also be real in the sense that

$$(\zeta^{\alpha\dot{\beta}})^* = \zeta^{\dot{\alpha}\beta} = \zeta^{\beta\dot{\alpha}} \quad , \tag{B.75}$$

where the last equality comes from the fact that the order of dotted and undotted indices is unimportant; these indices are in two different spaces. Mixed bispinors form an irreducible representation of $SL(2,C)$ with four real entries. If representations of $SL(2,C)$ can indeed be identified with representations of $SO(3,1)$, then a relation between this irreducible 4-dimensional representation and the representation x^μ from Section B.1 is possible.

Writing $\zeta^{\alpha\dot{\beta}}$ as a 2×2 matrix, we see that the relation (B.75) translates into a hermiticity condition on ζ.[32] Any hermitian 2×2 matrix may be expanded in terms of the identity and the Pauli matrices,

$$\sigma^1 \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma^2 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma^3 \equiv \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad . \tag{B.76}$$

Writing $\overline{\sigma}^\mu \equiv (1, -\vec{\sigma}\,)$, we make the identification

$$\zeta^{\alpha\dot{\beta}} = (x^\mu \overline{\sigma}_\mu)^{\alpha\dot{\beta}} \quad . \tag{B.77}$$

We would like for ζ to behave under $SL(2,C)$ rotations in such a way that this relation is preserved. An infinitesimal $SL(2,C)$ transformation has six independent parameters, and so can be expressed

$$(T)^\alpha{}_\beta = \delta^\alpha_\beta - \frac{i}{2}\theta_{\mu\nu}\,(\Sigma^{\mu\nu})^\alpha{}_\beta \quad , \tag{B.78}$$

where the Σ are linearly independent 2×2 matrices. Note that we have emphasized the fact that α is the first index on T and β the second. The transformation for ζ is

$$\begin{aligned} (T\zeta)^{\alpha\dot{\beta}} &= \zeta^{\alpha\dot{\beta}} - \frac{i}{2}\theta_{\mu\nu}\,(\Sigma^{\mu\nu})^\alpha{}_\gamma\,\zeta^{\gamma\dot{\beta}} + \frac{i}{2}\theta_{\mu\nu}\,(\Sigma^{\mu\nu\,*})^{\dot{\beta}}{}_{\dot{\gamma}}\,\zeta^{\alpha\dot{\gamma}} \\ &= \zeta^{\alpha\dot{\beta}} - \frac{i}{2}\theta_{\mu\nu}\left(\Sigma^{\mu\nu}\zeta - \zeta(\Sigma^{\mu\nu})^\dagger\right)^{\alpha\dot{\beta}} \quad , \end{aligned} \tag{B.79}$$

where in the second line we have suppressed the indices. The transposition comes from the convention of matrix multiplication. In another frame, the new ζ should be related to the

[32] The transposition comes from the convention that undotted indices should label rows and the dotted ones columns. Of course, this convention is arbitrary, but one must be consistent with whatever convention is chosen.

new x in the same way as the old ones for *any* choice of x^μ. According to (B.77), (B.79), and (B.22), we require

$$-i\Sigma^{\mu\nu}\overline{\sigma}^\lambda + i\overline{\sigma}^\lambda\left(\Sigma^{\mu\nu}\right)^\dagger = g^{\nu\lambda}\overline{\sigma}^\mu - g^{\mu\lambda}\overline{\sigma}^\nu \quad . \tag{B.80}$$

If we can solve this equation for Σ, we have found a covariant relationship between the **4** of $SL(2,C)$ and the **4** of $SO(3,1)$.

This equation is not as complicated as it seems; the easiest way to solve it is to recognize that each element is just a 2×2 matrix. *Any* 2×2 complex matrix can be written as a linear combination of the Pauli matrices and the identity, if we allow complex coefficients. Taking special values for μ, ν, and λ then allows us to deduce the coefficients. Defining

$$\left(\sigma^\mu\right)_{\dot{\alpha}\beta} \equiv \epsilon_{\dot{\alpha}\dot{\beta}}\epsilon_{\beta\alpha}\left(\overline{\sigma}^\mu\right)^{\alpha\dot{\beta}} \quad , \tag{B.81}$$

we have

$$\left(\Sigma^{\mu\nu}\right)^\alpha{}_\beta = \left(\overline{\sigma}^{\mu\nu}\right)^\alpha{}_\beta \equiv \frac{i}{4}\left(\overline{\sigma}^\mu\sigma^\nu - \overline{\sigma}^\nu\sigma^\mu\right)^\alpha{}_\beta \quad . \tag{B.82}$$

It is easy to show that these objects satisfy the commutation relations of $SO(3,1)$, Eq.(B.39).

The relationship between vector indices and spinor indices allows us to understand the construction of $SO(3,1)$'s irreducible tensor representations. According to Eq.(B.77), we have the correspondence

$$T^{\mu_1\cdots\mu_n} \sim \tau^{\alpha_1\cdots\alpha_n\dot{\beta}_1\cdots\dot{\beta}_n} \tag{B.83}$$

for an arbitrary Lorentz tensor T. We can form irreducible representations of $SL(2,C)$ by systematically removing the traces of τ with respect to like indices. Studying the symmetry of different representations under the interchange of two *Lorentz* indices, i.e. the simultaneous replacement $(\alpha_i\dot{\beta}_i, \alpha_j\dot{\beta}_j) \to (\alpha_j\dot{\beta}_j, \alpha_i\dot{\beta}_i)$, we see that they can be constructed to be either symmetric or antisymmetric.[33]

If antisymmetric, one can either have

$$\epsilon^{\alpha_i\alpha_j}\zeta^{(\dot{\beta}_i\dot{\beta}_j)} \tag{B.84}$$

or

$$\epsilon^{\dot{\beta}_i\dot{\beta}_j}\zeta^{(\alpha_i\alpha_j)} \quad . \tag{B.85}$$

[33] In reality, the construction is not quite this simple. The natural irreducible representations of $SL(2,C)$ do not lend themselves to interpretation in $SO(3,1)$. However, certain linear combinations of them have symmetry properties which are readily understood within $SO(3,1)$.

Neither of these is hermitian, but the sum and the difference (divided by i) both are. These combinations will be related to irreducible tensor representations of the Lorentz group which are antisymmetric under the interchange of two indices.

If symmetric, we have two possibilities. First, all four indices could be contained within metric tensors :

$$\epsilon^{\alpha_i\alpha_j}\epsilon^{\dot{\beta}_i\dot{\beta}_j} \quad . \tag{B.86}$$

Inspection of the relationship (B.77) between $SO(3,1)$ and $SL(2,C)$ reveals that this object represents the metric tensor $g^{\mu\nu}$ of $SO(3,1)$. Hence this possibility leads to a trace in Lorentz space. Alternatively, the structure

$$\zeta^{(\alpha_i\alpha_j)(\dot{\beta}_i\dot{\beta}_j)} \tag{B.87}$$

satisfies our requirement. This object is obviously related to a symmetric Lorentz tensor which does not have a trace. This simple argument can be used to construct the irreducible breakdowns of *any* tensor in $SO(3,1)$. In particular, it implies that the largest representation possible is the fully symmetric and traceless combination

$$T^{(\mu_1\cdots\mu_n)} \quad . \tag{B.88}$$

Since this tensor corresponds to a spinor tensor with n symmetrized dotted indices and n symmetrized undotted indices, its spin is $s = 2n/2 = n$.

To see how this works explicitly, let us study the decomposition of the tensor

$$T^{\mu_1\mu_2\mu_3} = \tau^{\alpha_1\alpha_2\alpha_3\dot{\beta}_1\dot{\beta}_2\dot{\beta}_3} \quad . \tag{B.89}$$

In order to avoid overcounting, we must proceed as systematically as possible. With this in mind, we leave the dotted indices alone for the moment and work only with the undotted. The possible structures are

$$\tau_s = \tau^{(\alpha_1\alpha_2\alpha_3)\dot{\beta}_1\dot{\beta}_2\dot{\beta}_3} \tag{B.90}$$

$$\tau_{t2} = \epsilon^{\alpha_1\alpha_2}\zeta^{\alpha_3\dot{\beta}_1\dot{\beta}_2\dot{\beta}_3} \tag{B.91}$$

$$\tau_{t3} = \epsilon^{\alpha_1\alpha_3}\zeta^{\alpha_2\dot{\beta}_1\dot{\beta}_2\dot{\beta}_3} \quad . \tag{B.92}$$

Note that we did not take all three possible traces as they are not all independent. Traces in $SL(2,C)$ are related to antisymmetric structures, so the symmetric combination

$$\epsilon^{\alpha\beta}\zeta^{\gamma} + \epsilon^{\beta\gamma}\zeta^{\alpha} + \epsilon^{\gamma\alpha}\zeta^{\beta} = 0 \tag{B.93}$$

is constrained to vanish. This can be seen by explicitly checking all three components. In general, the procedure is to *choose one index* and systematically remove all of the traces involving *that index*. Other traces are constrained by consistency relations like (B.93). The dotted indices decompose for *each* of the above structures in the same way.

Equation (B.90) gives the irreducible representations

$$\tau_{ss} = \tau^{(\alpha_1\alpha_2\alpha_3)(\dot\beta_1\dot\beta_2\dot\beta_3)} \tag{B.94}$$

$$\tau_{st2} = \epsilon^{\dot\beta_1\dot\beta_2}\zeta^{(\alpha_1\alpha_2\alpha_3)\dot\beta_3} \tag{B.95}$$

$$\tau_{st3} = \epsilon^{\dot\beta_1\dot\beta_3}\zeta^{(\alpha_1\alpha_2\alpha_3)\dot\beta_2} \quad . \tag{B.96}$$

The first of these is the fully symmetric spin-3 contribution. It corresponds to the fully symmetric and traceless

$$T^{(\mu_1\mu_2\mu_3)} \quad . \tag{B.97}$$

Since three fully symmetrized $SL(2,C)$ indices have four components : all the same (2), one different (2), this representation has $4 \cdot 4 = 16$ components.

Turning to the trace terms, we see that neither (B.95) nor (B.96) is hermitian, so neither can directly be related to representations of $SO(3,1)$. However, their complex conjugates are contained in[34] τ_{t2s} and τ_{t3s}. Representations of $SO(3,1)$ can only be formed with these 'mixed' spinor tensors via hermitian combinations. Since we cannot separate these from each other while at the same time retaining hermiticity and completeness of the representation, τ_{st2} and τ_{t2s} must *combine* to form *one* irreducible representation of $SO(3,1)$,

$$T^{[\mu_1(\mu_2]\mu_3)} \quad . \tag{B.98}$$

The fact that $SO(3,1)$ possesses these two invariant subgroups is already obvious from the behavior of its generators in Eqs.(B.46) and (B.47). The number of components the tensor in (B.95) contains is $4 \cdot 2 = 8$; (B.98) contains 16 components, as does its friend

$$T^{[\mu_1(\mu_3]\mu_2)} \quad . \tag{B.99}$$

Note that in order to actually form representations with the indicated symmetry structure, we

[34] My notation reads 'symmetric', 'trace with α_2', and 'trace with α_3'. The order refers undotted versus dotted indices.

would need to involve the tensor

$$\epsilon^{\alpha_2\alpha_3}\zeta^{\alpha_1(\dot\beta_1\dot\beta_2\dot\beta_3)} \tag{B.100}$$

and its hermitian conjugate. With these six tensors, we can form three hermitian Lorentz structures. However, in light of the above discussion, only two of them are independent.

We are left with

$$\tau_{t2t3} = \epsilon^{\alpha_1\alpha_2}\epsilon^{\dot\beta_1\dot\beta_3}\zeta^{\alpha_3\dot\beta_2} \tag{B.101}$$

$$\tau_{t3t2} = \epsilon^{\alpha_1\alpha_3}\epsilon^{\dot\beta_1\dot\beta_2}\zeta^{\alpha_2\dot\beta_3} \tag{B.102}$$

$$\tau_{t2t2} = \epsilon^{\alpha_1\alpha_2}\epsilon^{\dot\beta_1\dot\beta_2}\zeta^{\alpha_3\dot\beta_3} \tag{B.103}$$

$$\tau_{t3t3} = \epsilon^{\alpha_1\alpha_3}\epsilon^{\dot\beta_1\dot\beta_3}\zeta^{\alpha_2\dot\beta_2} \quad . \tag{B.104}$$

The first two of these expressions must be combined to form a representation of $SO(3,1)$, as before. In this case, the presence of two metric tensors allows us to construct *separate* hermitian representations. Involving the other four mixed-symmetric two-trace tensor structures allows us to construct a fully symmetric[35] tensor which can be identified as the trace we removed from (B.97) and a fully antisymmetric tensor which is identified with

$$T^{[\mu_1\mu_2\mu_3]} \quad . \tag{B.105}$$

Both of these tensors have 4 components. The remaining two structures, τ_{t2t2} and τ_{T3T3}, are readily identified in $SO(3,1)$ as the other two traces :

$$\tau_{t2t2} \sim g^{\mu_1\mu_2}P^{\mu_3} \tag{B.106}$$

$$\tau_{t3t3} \sim g^{\mu_1\mu_3}P^{\mu_2} \quad . \tag{B.107}$$

Hence in $SL(2,C)$ we have the decomposition

$$\mathbf{2}\times\overline{\mathbf{2}}\times\mathbf{2}\times\overline{\mathbf{2}}\times\mathbf{2}\times\overline{\mathbf{2}} = \mathbf{16}+(\mathbf{8}+\overline{\mathbf{8}})+(\mathbf{8}+\overline{\mathbf{8}})+(\mathbf{4}+\overline{\mathbf{4}})+\mathbf{4}+\mathbf{4} \quad , \tag{B.108}$$

while in $SO(3,1)$,

$$\mathbf{4}\times\mathbf{4}\times\mathbf{4} = \mathbf{16}_S+\mathbf{16}_M+\mathbf{16}_M+\mathbf{4}_S+\mathbf{4}_M+\mathbf{4}_M+\mathbf{4}_A \tag{B.109}$$

is induced. Note that in $SO(3,1)$ all three possible traces show up in the final decomposition, the third being formed from the combination of $SL(2,C)$'s $\mathbf{4}+\overline{\mathbf{4}}$ which is orthogonal to the fully antisymmetric tensor. This is because traces in $SO(3,1)$ are unconstrained.

[35] under the exchange of *pairs* of indices, i.e. $(\alpha_i\dot\beta_i,\alpha_j\dot\beta_j)\to(\alpha_j\dot\beta_j,\alpha_i\dot\beta_i)$.

This procedure can be extended to tensors of higher rank without incident. While somewhat tedious, it certainly beats counting components and enforcing constraint relations to avoid overcounting. An equivalent procedure one can use to directly obtain the representations of $SO(3,1)$ is offered by Young's Tableaux. The most useful reference that I have come across on this method is [90].

At some point, we would like to use all of this formalism to do physics. Consider a classical spinor field, $\xi^\alpha(x)$. Following the discussion of the last section on irreducible representations of the Poincaré group, we wish to diagonalize the momentum operator $\hat{P}^\mu$. This operator has two spinor representations, $(\hat{P}\cdot\sigma)_{\dot{\alpha}\beta}$ and $(\hat{P}\cdot\overline{\sigma})^{\alpha\dot{\beta}}$. Each representation necessarily has one dotted and one undotted index in $SL(2,C)$, so the object $\hat{P}\xi$ cannot transform in the same way as ξ itself. This leaves us with two options : we can either require $\hat{P}$ to have eigenvalue zero or allow it to change our spinor into another kind of spinor. The former option leads to the massless Weyl spinors that describe neutrinos, and the latter leads to the massive Dirac spinors appropriate for electrons.

Weyl spinors are obtained simply from the requirement

$$(\hat{P}\cdot\sigma)_{\dot{\alpha}\beta}\xi^\beta = 0 \quad . \tag{B.110}$$

Contracting with $(\hat{P}\cdot\overline{\sigma})^{\gamma\dot{\alpha}}$ and using the identity

$$(\overline{\sigma}^\mu\sigma^\nu)^\alpha{}_\beta = g^{\mu\nu}\delta^\alpha_\beta - 2i\,(\overline{\sigma}^{\mu\nu})^\alpha{}_\beta \quad , \tag{B.111}$$

it is easily seen that the casimir $\hat{P}^2$ has eigenvalue zero so these spinors can only represent massless states. Within the proper Lorentz group, this spinor stands alone; it forms its own irreducible representation. However, under parity our spinor field transforms as[36]

$$\mathcal{P}\xi^\alpha = \eta_{\dot{\alpha}} \quad . \tag{B.112}$$

Hence if we wish to consider representations of the full Lorentz group, we must include the

[36] This can be seen in the following way : $\mathcal{P}\xi^\alpha$ cannot transform like ξ^α because $\mathcal{P}$ does not commute with all Lorentz transformations. However, since the spatial part of $\mathcal{P}$ is proportional to the identity, it will commute with spatial rotations. For spatial rotations, our $SL(2,C)$ matrix is unitary so $(T^{-1})^\dagger = T$. This implies that $\mathcal{P}\xi^\alpha$ transforms like $\eta_{\dot{\alpha}}$.

field $\eta_{\dot\alpha}$ with ξ^α in our massless representation.[37] This field satisfies

$$(\hat{P}\cdot\overline{\sigma})^{\alpha\dot\beta}\eta_{\dot\beta}=0 \ . \tag{B.113}$$

Since we wish to consider these fields together, we write

$$\psi=\begin{pmatrix}\eta_{\dot\alpha}\\ \xi^\alpha\end{pmatrix} \ , \tag{B.114}$$

for which the statement of translation invariance becomes

$$\hat{P}_\mu\begin{pmatrix}0 & (\sigma^\mu)_{\dot\alpha\beta}\\ (\overline{\sigma}^\mu)^{\alpha\dot\beta} & 0\end{pmatrix}\psi=0 \ . \tag{B.115}$$

The 4-component spinor ψ is a very strange object. It mixes dotted and undotted indices in a way we have not yet encountered. This mixing has been forced upon us by parity. Systems in which parity is not conserved, like the weak interactions, do not require us to consider both spinors together. In fact, it has long been thought that the analogue of ξ^α for neutrinos simply does not exist. Here we are concerned with the strong and electromagnetic interactions, both of which conserve parity, so the existence of one helicity necessarily implies the existence of the other. The 4×4 matrices in the parentheses come up over and over again. We will call them γ^μ. We also introduce $\hat{\not{P}}$ as shorthand for $\hat{P}_\mu\gamma^\mu$. Using this compact notation, we write the 'equation of motion' for massless spinor fields as

$$\hat{\not{P}}\psi=0 \ . \tag{B.116}$$

For a massless spinor moving in the 3-direction, our equation of motion reduces to

$$2p^0\begin{pmatrix}0&0&0&0\\0&0&0&1\\1&0&0&0\\0&0&0&0\end{pmatrix}\psi=0 \ , \tag{B.117}$$

which has two linearly independent solutions,

$$u(p,+)=\sqrt{2p^0}\begin{pmatrix}0\\0\\1\\0\end{pmatrix} \quad \text{and} \quad u(p,-)=\sqrt{2p^0}\begin{pmatrix}0\\1\\0\\0\end{pmatrix} \ , \tag{B.118}$$

[37] This is a specific example of the general argument given in the last section that parity relates the two helicities $\pm\lambda$ to each other. We will see below that ξ^α corresponds to massless states with helicity $+1/2$ while $\eta_{\dot\alpha}$ describes states with helicity $-1/2$.

which we normalize in such a way that the following relations hold

$$u^\dagger(p,+)u(p,+) = u^\dagger(p,-)u(p,-) = 2p^0 \tag{B.119}$$

$$\overline{u}(p,+)u(p,+) = \overline{u}(p,-)u(p,-) = 0 \tag{B.120}$$

$$\overline{u}(p,+)u(p,-) = \overline{u}(p,-)u(p,+) = 0 \tag{B.121}$$

$$\sum_{h=\pm} u(p,h)\overline{u}(p,h) = \not{p} \ , \tag{B.122}$$

where I have defined $\overline{u} \equiv u^\dagger\gamma^0$. Note that (B.119) is not Lorentz invariant since $\psi^\dagger\psi$ involves a sum over dotted and undotted indices. However,

$$\overline{\psi}\psi = \eta_\alpha\xi^\alpha + \xi^{\dot\alpha}\eta_{\dot\alpha} \tag{B.123}$$

is invariant since it involves contractions with like indices only. (B.120) is simply the statement that for massless spinors the upper and lower components are uncoupled. This is not the case for massive spinors, where the momentum operator itself couples dotted and undotted indices.

Since $\overline{\psi}\psi$ is invariant and $\gamma^\mu\psi$ transforms in $SL(2,C)$ like ψ itself, the combination $\overline{\psi}\gamma^\mu\psi$ transforms under the action of the Lorentz group as a vector (it is invariant under $SL(2,C)$). This means that $\psi^\dagger\psi = \overline{\psi}\gamma^0\psi$ is just the time component of a 4-vector. The last relation is a statement about the completeness of our solutions. It is covariant in the sense that both sides transform like the $SL(2,C)$ indices on a γ-matrix.

Under Lorentz transformations, the lower components of ψ transform according to (B.82). The upper components transform like

$$(\Sigma^{\mu\nu})_{\dot\alpha}^{\ \dot\beta} = \left(\overline{\sigma}^{\mu\nu\,\dagger}\right)_{\dot\alpha}^{\ \dot\beta} = (\sigma^{\mu\nu})_{\dot\alpha}^{\ \dot\beta} \equiv \frac{i}{4}\left(\sigma^\mu\overline{\sigma}^\nu - \sigma^\nu\overline{\sigma}^\mu\right)_{\dot\alpha}^{\ \dot\beta} \ , \tag{B.124}$$

so ψ transforms according to[38]

$$\Sigma^{\mu\nu} = \begin{pmatrix} \sigma^{\mu\nu} & 0 \\ 0 & \overline{\sigma}^{\mu\nu} \end{pmatrix} = \frac{i}{4}\left[\gamma^\mu,\gamma^\nu\right] \equiv \frac{1}{2}\sigma^{\mu\nu} \ . \tag{B.125}$$

Although this is not a hermitian operator and certainly cannot be identified with $\hat{J}$,[39] it does describe the action of the Lorentz group on classical fields ψ. For a *classical* massless particle

[38] Since the (2×2) $\sigma^{\mu\nu}$ has dotted and undotted $SL(2,C)$ indices and this new (4×4) $\sigma^{\mu\nu}$ has only the mixed 'Dirac' indices, confusion should be rare. Throughout the text, the (2×2) $\sigma^{\mu\nu}$ does not make an appearence.

[39] We will see how this object is related to $\hat{J}$ in Section D.2.

moving in the 3-direction, the relevant helicity operator is

$$\Sigma^{12} = \begin{pmatrix} \sigma^3/2 & 0 \\ 0 & \sigma^3/2 \end{pmatrix} \quad . \tag{B.126}$$

The action of this operator on our u's tells us that $u(p,+)$ represents a particle with helicity $+1/2$ and $u(p,-)$ represents a particle with helicity $-1/2$. In this sense, massless spinors can be said to have spin-1/2.

Dirac spinors are obtained by writing

$$(\hat{P}\cdot\sigma)_{\dot{\alpha}\beta}\xi^{\beta} = m\,\eta_{\dot{\alpha}} \quad , \tag{B.127}$$

with m some real constant. Since $\hat{P}^2$ is a casimir of the Poincaré group, consistency requires the relation[40]

$$(\hat{P}\cdot\overline{\sigma})^{\alpha\dot{\beta}}\eta_{\dot{\beta}} = m\xi^{\alpha} \quad . \tag{B.128}$$

Once again, we combine the two spinor fields into one 4-component spinor. Using the same notation, we write the equation of motion for massive spinors as

$$(\hat{\not{P}} - m)\psi = 0 \quad . \tag{B.129}$$

For a spinor at rest, this equation reduces to

$$\begin{pmatrix} -m & 0 & m & 0 \\ 0 & -m & 0 & m \\ m & 0 & -m & 0 \\ 0 & m & 0 & -m \end{pmatrix} \psi = 0 \quad . \tag{B.130}$$

Once again, we have two linearly independent solutions :

$$u(m,\uparrow) = \sqrt{m}\begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad u(m,\downarrow) = \sqrt{m}\begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix} \quad . \tag{B.131}$$

The analogues of (B.119), (B.120), (B.121), and (B.122) are

$$u^{\dagger}(p,\uparrow)u(p,\uparrow) \quad = \quad u^{\dagger}(p,\downarrow)u(p,\downarrow) = 2p^0 \tag{B.132}$$

[40] We could have a different eigenvalue. In that case, the mass of the particle we are describing would be $\sqrt{mm'}$. However, we can always re-scale the fields to obtain the same eigenvalue.

$$\overline{u}(p,\uparrow)u(p,\uparrow) = \overline{u}(p,\downarrow)u(p,\downarrow) = 2m \tag{B.133}$$

$$\overline{u}(p,\uparrow)u(p,\downarrow) = \overline{u}(p,\downarrow)u(p,\uparrow) = 0 \tag{B.134}$$

$$\sum_{s=\updownarrow} u(p,s)\overline{u}(p,s) = \not{p} + m \ . \tag{B.135}$$

Here, we have used the fact that (B.132) is actually the time component of a vector and (B.133) is invariant under the Lorentz group to write the appropriate expressions on the right-hand-sides of these relations. Substitution of our expressions into (B.135) actually gives $m\gamma^0 + m1$, where 1 is the identity in Dirac space. Since the left-hand-side has the $SL(2,C)$ indices of a γ-matrix as before, we deduce that the correct extension of $m\gamma^0 + m$ to other frames is $\not{p} + m$. This can be checked by transforming the solutions to other frames and explicitly calculating the completeness sum.

Under Lorentz transformations, massive spinors transform via the same generators as massless spinors. The spin vector introduced in Section B.2 is given by the spatial components of Σ. Application of these matrices reveals that $u(m,\uparrow)$ and $u(m,\downarrow)$ represent particles with spin-1/2 up and down along the z-axis, respectively. Spinors representing particles with spin along other axes can be written as linear combinations of these solutions.

The γ-matrices have many nice properties which make them easy to work with. The anticommutation relation,

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} \ , \tag{B.136}$$

can easily be proven using (B.111). This relation tells us that to span the space of possible Dirac matrices we need only consider antisymmetric products. Our explicit expression for the antisymmetric combination of two γ-matrices, $\sigma^{\mu\nu}$, tells us that this object represents six Dirac matrices that are linearly independent of each other. Since there are only four gamma matrices, the antisymmetric combination of three will have one missing. Multiplying by the square of the missing matrix (which is proportional to the identity), we see that this combination can be written $\gamma^\mu\gamma^0\gamma^1\gamma^2\gamma^3$. The antisymmetric product of four matrices must include all of them. Defining

$$\gamma_5 \equiv i\gamma^0\gamma^1\gamma^2\gamma^3 = -\frac{i}{4!}\epsilon^{\mu\nu\alpha\beta}\gamma_\mu\gamma_\nu\gamma_\alpha\gamma_\beta \ , \tag{B.137}$$

we see that the space of Dirac matrices is spanned by the linearly independent set

1	Scalar

$$\begin{array}{rl} \gamma^\mu & \text{Vector} \\ \sigma^{\mu\nu} & \text{Tensor} \\ \gamma^\mu\gamma_5 & \text{Pseudovector} \\ \gamma_5 & \text{Pseudoscalar} \ . \end{array} \tag{B.138}$$

After each matrix, I have noted the Lorentz transformation properties of the bilinear $\overline{\psi}\,\Gamma\,\psi$. The prefix 'pseudo' refers to the transformation properties under parity. The presence of ϵ in the definition of γ_5 means that it has an extra sign change under parity. The properties

$$(\gamma^\mu)^\dagger = \gamma^0\gamma^\mu\gamma^0 \tag{B.139}$$

$$(\sigma^{\mu\nu})^\dagger = \gamma^0\sigma^{\mu\nu}\gamma^0 \tag{B.140}$$

$$(\gamma_5)^\dagger = \gamma_5 \tag{B.141}$$

$$\{\gamma^\mu, \gamma_5\} = 0 \tag{B.142}$$

imply that all these bilinears are real except the pseudoscalar, which is purely imaginary. More relations between the γ-matrices can be found in Appendix A.1.

The matrix γ_5 is very useful in forming spin projection operators. The fact that $\gamma_5^2 = 1$ along with its explicit form in the Weyl basis,

$$\gamma_5 = \begin{pmatrix} -\mathbf{1} & 0 \\ 0 & \mathbf{1} \end{pmatrix} \ , \tag{B.143}$$

imply that the operators

$$P_R \equiv \frac{1}{2}\,(1+\gamma_5) \qquad\qquad P_L \equiv \frac{1}{2}\,(1-\gamma_5) \tag{B.144}$$

project out the undotted and dotted, or positive and negative helicity states, respectively. Furthermore, it can easily be checked that the bilinear

$$\overline{u}(p,s)\gamma^\mu\gamma_5 u(p,s) = 4ms^\mu \tag{B.145}$$

returns the spin vector for massive particles. This formula is also valid for massless fermions, provided the replacement $ms^\mu \to hp^\mu$ is understood. For massless particles, the projection operators are extremely useful. According to (B.122), the bilinear

$$u(p,h)\overline{u}(p,h) = \frac{1}{2}\,(1+2h\gamma_5)\ \not{p} \ . \tag{B.146}$$

Here, h is the *true* helicity of the fermion, taking the values $\pm 1/2$.

Up until this point, we have focused on *positive* energy solutions, i.e. $p^0 > 0$. However, for reasons which will become clear in Appendix D.1, we must also consider the case in which $p^0 < 0$. Here, we define $k^\mu \equiv -p^\mu$ and write

$$\hat{\not{P}}\, v(k,h) = \not{p}\, v(k,h) = -\not{k}\, v(k,h) = 0 \tag{B.147}$$

for massless fermions and

$$\hat{\not{P}}\, v(k,s) = \not{p}\, v(k,s) = -\not{k}\, v(k,s) = m v(k,s) \tag{B.148}$$

for massive. From Equation (B.147), it is clear that the solutions $v(k,h)$ for massless fermions correspond exactly to the positive energy solutions :

$$v(\vec{p},h) = u(-\vec{p},-h) \ . \tag{B.149}$$

Calling the objects represented by u fermions and those represented by v antifermions, this relation implies that an antifermion with positive helicity moving along the positive z-axis is equivalent to a fermion with negative helicity moving along the negative z-axis.

For massive fermions, the correspondence is not so simple. In this case, the Dirac equation becomes

$$\begin{pmatrix} m & 0 & m & 0 \\ 0 & m & 0 & m \\ m & 0 & m & 0 \\ 0 & m & 0 & m \end{pmatrix} v(k,s) = 0 \tag{B.150}$$

in the antifermion rest frame. Once again, we have two linearly independent solutions which we can take as

$$v(k,\downarrow) = \sqrt{m}\begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix} \quad \text{and} \quad v(k,\uparrow) = \sqrt{m}\begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix} \ . \tag{B.151}$$

These solutions also refer to a spin-1/2 object, but the spin-up and spin-down solutions have changed roles owing to an extra sign in the Pauli-Ljubanskii vector. Here, we have the relations

$$v^\dagger(k,\uparrow)v(k,\uparrow) \quad = \quad v^\dagger(k,\downarrow)v(k,\downarrow) = 2k^0 \tag{B.152}$$

$$\overline{v}(k,\uparrow)v(k,\uparrow) = \overline{v}(k,\downarrow)v(k,\downarrow) = -2m \tag{B.153}$$

$$\overline{v}(k,\uparrow)v(k,\downarrow) = \overline{v}(k,\downarrow)v(k,\uparrow) = 0 \tag{B.154}$$

$$\overline{v}(k,s)u(p,s') = \overline{u}(p,s)v(k,s') = 0 \tag{B.155}$$

$$\sum_{s=\updownarrow} v(k,s)\overline{v}(k,s) = -(\not{p}+m) = \not{k} - m \ . \tag{B.156}$$

Explicit forms can be found for both $u(p,s)$ and $v(p,s)$ by boosting the rest frame solutions via (B.125). The result is

$$u(p,s) = \begin{pmatrix} \sqrt{p\cdot\sigma}\,\xi_s \\ \sqrt{p\cdot\overline{\sigma}}\,\xi_s \end{pmatrix} \quad \text{and} \quad v(k,s) = \begin{pmatrix} \sqrt{k\cdot\sigma}\,\xi_{-s} \\ -\sqrt{k\cdot\overline{\sigma}}\,\xi_{-s} \end{pmatrix} \ , \tag{B.157}$$

where ξ_s is a 2-component spinor aligned with the direction $\vec{s}$,

$$\vec{s}\cdot\vec{\sigma}\,\xi_s = \xi_s \ . \tag{B.158}$$

These solutions are constructed so that in the rest frame $\vec{s}\cdot\vec{W}$ is diagonal. The square-root of a matrix is understood as an instruction to take the *positive* square root of each eigenvalue.

Appendix C

$SU(N)$

The special unitary group in N dimensions, $SU(N)$, has many applications throughout physics. For $N = 2$, we have a group structure which is homomorphic to the rotation group, $SO(3)$. Its representations house non-relativistic fields with intrinsic angular momentum, or spin, as well as the orbital excitations of molecules. Taking N as the number of quark flavors allows us to represent the approximate flavor invariance of the QCD lagrangian. This symmetry was the first true indication that hadrons are composite. $SU(2n_f)$ describes the approximate spin-flavor symmetry of a theory with n_f extremely massive quarks. Most recently, $SU(2)$ and $SU(3)$ along with the sister group $U(1)$ have been useful in describing the full symmetry group of the Standard Model. Although this thesis is most concerned with the group $SU(3)$ and its role in strong interaction physics, we will find it useful to familiarize ourselves with the general structure of this family of groups.

C.1 General Properties

In this section, we will discuss the definition and some of the mathematical properties of $SU(N)$. Since we are concerned mainly with the fundamental representation of $SU(N)$, we begin with a discussion in this context. The fundamental representation of $SU(N)$ can be realized by the group of unitary $N \times N$ matrices with determinant $+1$. Suppose the vector q^i, $i = 1$ to N, transforms in the fundamental representation of $SU(N)$. Then

$$q^i \to (Uq)^i \quad . \tag{C.1}$$

This implies that the complex conjugate transforms as

$$(q^i)^* \to ((Uq)^i)^* = (U^* q^*)^i = ((U^\dagger)^{\mathrm{T}} q^*)^i = (q^* U^\dagger)^i \quad . \tag{C.2}$$

Since U is unitary, the product $q^* q$ is invariant under the transformation. In analogy with the Lorentz group, let us denote $(q^i)^* \equiv q_i$. Objects which transform as U are said to be in the $\mathbf{N}$ of $SU(N)$, while their conjugates are in the $\overline{\mathbf{N}}$. Since $q^* q = q_i q^i = q_i q^j \delta^i_j$ is invariant, we have an invariant Krönekker delta. The invariance of this object is directly related to the uniterity of the elements of $SU(N)$, which prompts us to look for an invariant related to the determinant of U. Just as in the case with the Lorentz group, the totally antisymmetric tensor $\epsilon^{i_1 \cdots i_N}$ does the trick. These are the only two invariants in the fundamental representation of $SU(N)$.

Now that we have discussed the objects these transformations act on, let us parameterize a general element of the group in this representation. Any unitary matrix can be written

$$U = e^{iH} \quad , \tag{C.3}$$

where H is a hermitian matrix. Using the matrix identity [1] $\log \det U = \mathrm{Tr} \log U$, we find that $\det U = 1 \leftrightarrow \mathrm{Tr}\, H = 2n\pi$, with $n \in \mathbf{Z}$. Any $N \times N$ matrix can be decomposed into traceless part and a a trace :

$$H = \left(H - \frac{1}{N} \mathrm{Tr}\, H \right) + \frac{1}{N} \mathrm{Tr}\, H \quad . \tag{C.6}$$

Hence our unitary matrix U is the product of the exponential of a traceless matrix and a phase, $e^{2in\pi/N}$. Since this phase can only attain certain discreet values, one cannot move continuously from one value of n to another. Furthermore, only the set with $n = 0$ is a

[1] This identity can be proven for any matrix which is the exponential of a complex multiple of a hermitian matrix as follows. Given

$$\begin{aligned} \mathcal{M} &= e^{\alpha H} \ ; \\ U^\dagger H U &= \tilde{H} \ , \end{aligned} \tag{C.4}$$

with $\tilde{H}$ diagonal and U unitary, we see immediately that

$$\det \left(U^\dagger \mathcal{M} U \right) = \det \mathcal{M} = \det e^{\alpha \tilde{H}} = e^{\alpha\, \mathrm{Tr}\, \tilde{H}} = e^{\alpha\, \mathrm{Tr}\, H} = e^{\mathrm{Tr} \log \mathcal{M}} \quad . \tag{C.5}$$

The validity of this identity rests mainly on two facts : that hermitian matrices can be diagonalized by a unitary transformation and that the relation is trivially true for diagonal matrices.

subgroup since it alone contains its inverses and the identity. I will call this subgroup the proper $SU(N)$ transformations and subsequently omit the 'proper'. The whole group can be considered (when it is absolutely necessary to do so) as the product of this $SU(N)$ with the discreet $U(1)$ group $e^{2in\pi/N}$, $n \in \mathbf{Z}$. Some of the consequences of this phase to the global properties of the SM are discussed in [91].

We are now left with the task of parameterizing a traceless $N \times N$ hermitian matrix. Such a matrix has $(N^2 - N)$ complex off-diagonal elements constrained by $(N^2 - N)/2$ complex hermiticity constraints and N real diagonal elements constrained by 1 condition on the trace. In other words, they have $(N^2 - 1)$ independent elements. Any set of $(N^2 - 1)$ linearly independent traceless hermitian $N \times N$ matrices will act as a basis for this space. Since they exhaust the space of H, they generate all of the elements of $SU(N)$ through (C.3). The most natural way to choose these generators is[2]

$$\begin{aligned} (H)^i_j &= A^l_k (M^k_l)^i_j \ ; \\ (M^k_l)^i_j &= \delta^k_j \delta^i_l - \frac{1}{N} \delta^k_l \delta^i_j \ , \end{aligned} \tag{C.7}$$

since this choice does not introduce any new objects. All we are using to construct our generators are the invariants we have from the above discussion. Different choices of the parameters A^l_k will lead to different elements of the group.

Although there are in principle N^2 matrices M, they are not all linearly independent; they are related by $\delta^i_j (M^k_l)^j_i = 0$. This little inconvenience leads to several problems, not the least of which is that the A's contain more degrees of freedom than required. Another difficulty is that our generators, as it stands, are not hermitian. Consider the hermitian conjugate to (M^k_l) [3]

$$((M^k_l)^\dagger)^i_j \sim (M^k_l)^j_i \ . \tag{C.8}$$

This matrix has the same entries (if we use the convention that 'upper' indices label rows while 'lower' indices label columns) as $(M^l_k)^i_j$. Hence the hermitian matrices $(M^k_l + M^l_k)$

[2]Since $q^i \to (Uq)^i = U^i_j q^j$, our matrices must have one index in the fundamental and one in the conjugate representation.

[3]We use $\sim$ rather than $=$ here because the indices i and j are in different spaces on either side of the equation.

and $(M^k_l - M^l_k)/i$ should in principle be used as generators rather than M^k_l and M^l_k. These generators will have coefficients which are strictly real, while our original 'natural' choice will have complex coefficients constrained by the condition $(A^i_j)^* = A^j_i$. Although this is just the statement that complex conjugation interchanges the roles of the fundamental and complex representations, it is not easy to implement in practice. Since the sum $M^k_l + M^l_k$ makes sense only for specific values of k and l,[4] these indices can no longer be thought of as truly in *any* representation of $SU(N)$. This fact leads to many complications if one wishes to consider N large. However, in practice we seldom need to consider $N > 3$ so the hermitian choice is often made.

Let us consider generators which are all hermitian and which do not overcount the relevant degrees of freedom. Taking our new generators to be t^a, $a = 1$ to $(N^2 - 1)$, we write

$$U = e^{i\theta^a t^a} \quad . \tag{C.9}$$

The θ^a's are in a new $(N^2 - 1)$-dimensional space. As with the Lorentz group, these objects transform in the *adjoint* representation of $SU(N)$. Note that in the adjoint representation there are no 'upper' and 'lower' indices since the θ's are real.

Since we are working within a group, products of transformations are themselves transformations. In particular,[5]

$$\begin{aligned} UVU^\dagger V^\dagger &= e^{i\lambda\theta} e^{i\lambda\omega} e^{-i\lambda\theta} e^{-i\lambda\omega} \\ &= 1 + \lambda^2[\omega, \theta] + \mathcal{O}(\lambda^3) \end{aligned} \tag{C.10}$$

is an element of the group and hence is expressible in terms of the generators. This implies for small λ that

$$\omega^a \theta^b [t^a, t^b] = i\eta^c t^c \tag{C.11}$$

for some η. Differentiating with respect to ω and θ, we obtain

$$[t^a, t^b] = i f^{a\,b\,c} t^c \quad , \tag{C.12}$$

where I have introduced the $f^{a\,b\,c}$'s as the relevant derivatives of η. The $f^{a\,b\,c}$'s are called the *structure constants* of the group, and Equation (C.12) is said to define the *Lie algebra* of the

[4] We cannot have k or l in both the fundamental *and* the conjugate representation!

[5] I have suppressed the adjoint indices via $\theta \equiv \theta^a t^a$.

group. The structure constants are obviously antisymmetric in their first two indices, but we can make them totally antisymmetric by choosing the t^a's to be orthogonal in the sense that [6]

$$\mathrm{Tr}\, t^a t^b = T_F \delta^{a\,b} \ , \tag{C.13}$$

where $\delta^{a\,b}$ is the Krönekker delta in adjoint space and T_F is some positive normalization. With this choice,

$$f^{a\,b\,c} = -\frac{i}{T_F}\mathrm{Tr}\,[t^a, t^b]t^c \ . \tag{C.14}$$

Total antisymmetry follows from this identity and the cyclicity of the trace operation.

Since the Jacobi identity

$$[t^a,[t^b,t^c]] + [t^b,[t^c,t^a]] + [t^c,[t^a,t^b]] = 0 \tag{C.15}$$

is trivially true, we have a consistency relation for the $f^{a\,b\,c}$'s

$$f^{a\,b\,e}f^{c\,d\,e} + f^{b\,c\,e}f^{a\,d\,e} + f^{c\,a\,e}f^{b\,d\,e} = 0 \ . \tag{C.16}$$

Let us calculate the transformation law in the adjoint representation. Since the θ^a's and the A^i_j's play the same role, under an $SU(N)$ transformation $\theta \to U\theta U^\dagger$. For infinitesimal transformations, we have

$$\theta \to \theta + i[\omega, \theta] \ , \tag{C.17}$$

or

$$\theta^a \to \theta^a - f^{a\,b\,c}\omega^b\theta^c \ . \tag{C.18}$$

Since this is an $SU(N)$ transformation, it can be written $\theta^a \to U^{a\,b}\theta^b$, with U given by $e^{i\omega^a T^a_A}$, exactly in the same way as before. The only difference is that now the generators are $(N^2-1)\times(N^2-1)$ dimensional matrices. This implies the identification

$$(T^a_A)^{b\,c} = -if^{a\,b\,c} \ , \tag{C.19}$$

from which standpoint the Jacobi identity, (C.16), is simply a re-expression of (C.12) in the adjoint representation.

[6] The normalization is not important here. It need only be the same for all generators to induce fully antisymmetric structure constants. In some sense, choosing the same normalization for all of our generators restores the symmetry we had to break to obtain a basis which does not overcount the relevant degrees of freedom. Any other choice would have a favored direction in adjoint space.

Since the Krönekker delta and the structure constants seem to be properties of the group itself rather than a specific transformation, we expect them to be invariant under transformations. We can check this by explicit calculation :

$$\begin{aligned} \delta^{a\,b} &\rightarrow \delta^{a\,b} + f^{a\,c\,d}\theta^d\delta^{c\,b} + f^{b\,c\,d}\theta^d\delta^{a\,c} \\ &= \delta^{a\,b} \\ f^{a\,b\,c} &\rightarrow f^{a\,b\,c} - (f^{b\,c\,e}f^{a\,d\,e} + f^{c\,a\,e}f^{b\,d\,e} + f^{a\,b\,e}f^{c\,d\,e})\theta^d \qquad \text{(C.20)} \\ &= f^{a\,b\,c}\ , \qquad \text{(C.21)} \end{aligned}$$

where the last equality follows from the Jacobi identity, (C.16). In the adjoint representation the generators are the structure constants, implying that the generators are invariant under the action of the group if we rotate *all* of their indices. This is obvious with the natural choice for the generators, (C.7), since there they are expressed solely in terms of the invariant δ^i_j.

There is one other invariant in the adjoint representation which will be useful to us. This object appears because of the fact that the anticommutator of two generators is a hermitian $(N \times N)$ matrix and so can be represented as a linear combination of the generators themselves and the identity. We write

$$\left\{t^a, t^b\right\} = \alpha^{a\,b} + d^{a\,b\,c}t^c\ . \qquad \text{(C.22)}$$

Taking the trace of both sides, we find that $\alpha^{a\,b} = 2T_F/N\,\delta^{a\,b}$. The $d^{a\,b\,c}$'s are, with our choice of normalization, fully symmetric since

$$d^{a\,b\,c} = \frac{1}{T_F}\mathrm{Tr}\,\left\{t^a, t^b\right\}t^c\ . \qquad \text{(C.23)}$$

These structure constants transform according to

$$d^{a\,b\,c} \rightarrow d^{a\,b\,c} - (f^{a\,d\,e}d^{b\,c\,e} + f^{b\,d\,e}d^{c\,a\,e} + f^{c\,d\,e}d^{a\,b\,e})\theta^d\ . \qquad \text{(C.24)}$$

An identity similar to Jacobi's,

$$[t^a, \{t^b, t^c\}] + [t^c, \{t^a, t^b\}] + [t^b, \{t^c, t^a\}] = 0\ , \qquad \text{(C.25)}$$

can be employed to show that the quantity in parentheses is in fact zero. These definitions allow us to write

$$t^a t^b = \frac{1}{2}\left([t^a, t^b] + \{t^a, t^b\}\right) = \frac{1}{2}\left(\frac{2T_F}{N}\delta^{a\,b} + d^{a\,b\,c}t^c + if^{a\,b\,c}t^c\right)\ , \qquad \text{(C.26)}$$

a very useful result.

Since we have chosen the same normalization for all of our generators, there is no preferred direction in adjoint space. Any quantity which does not depend on an external adjoint quantity must be expressed in terms of invariants, in exactly the same way that Lorentz covariant quantities which do not depend on external vectors must be expressed in terms of the metric and the Levi-Cevita tensor. This is a *representation independent* result since it has only to do with the way we have normalized our generators. Since all other representations can be built from the fundamental representation, normalizing the t^a induces a normalization on T_R^a for any representation R. This normalization is given by

$$\mathrm{Tr}_R\, T_R^a T_R^b = T_R \delta^{a\,b} \quad , \tag{C.27}$$

where we have used the fact that the left-hand-side is independent of any external adjoints and hence must be expressed in terms of the invariant δ.

The value T_R depends only on the representation the trace is performed in. For example, let us calculate T_A[7]

$$\mathrm{Tr}_A\, T_A^a T_A^b = f^{a\,c\,d} f^{b\,c\,d} = T_A \delta^{a\,b} \quad . \tag{C.28}$$

The sum on the f's can be expressed in terms of the t's using (C.14) :

$$f^{a\,c\,d} f^{b\,c\,d} = -\frac{1}{T_F^2} \mathrm{Tr}[t^a, t^c] t^d \; \mathrm{Tr}[t^b, t^c] t^d \quad . \tag{C.29}$$

$(t^a)^i_j (t^a)^k_l$ is an object with two fundamental indices and two conjugate indices which does not depend on an external vector, hence it can be expressed in terms of the available invariants, $\delta^i_j \delta^k_l$ and $\delta^i_l \delta^k_j$. Since both matrices are traceless, we write

$$(t^a)^i_j (t^a)^k_l \propto \delta^i_l \delta^k_j - \frac{1}{N} \delta^i_j \delta^k_l \quad . \tag{C.30}$$

Contracting with $\delta^j_k \delta^l_i$ gives $\mathrm{Tr}\; t^a t^a = T_F \delta^{a\,b} \delta^{a\,b} = (N^2-1) T_F$, so

$$(t^a)^i_j (t^a)^k_l = T_F \left(\delta^i_l \delta^k_j - \frac{1}{N} \delta^i_j \delta^k_l \right) \quad . \tag{C.31}$$

This useful relation is known as a *Fiertz*[8] identity. Using it we can expand the trace and do the sum, obtaining

$$T_A = 2 N T_F \quad . \tag{C.32}$$

[7] Confusion between the generators T_R^a and their normalization T_R should be rare.

[8] pronounced almost like *fierce*

The operator $(T_R^a T_R^a)_{\alpha\beta}$ has two indices in the representation R and no adjoint indices. Hence it must be proportional to the identity in R, i.e.[9]

$$(T_R^a T_R^a)_{\alpha\beta} = C_R \delta_{\alpha\beta} \quad . \tag{C.33}$$

It is obvious from the Fiertz identity that $C_F = T_F(N^2-1)/N$ and $C_A = T_A$. As with the operators $\hat{P}^2$ and $\hat{W}^2$ in $SO(3,1)$, this operator is a casimir for $SU(N)$. From here, one can calculate many group structures. I list a few useful relations in Appendix A.2 for reference.

Explicit forms for the structure constants can be obtained for general $SU(N)$ if one uses the 'natural' choice for the generators, M_j^i. It is straightforward to check that

$$(M_j^i)_p^k (M_m^n)_l^p = \delta_j^k \delta_m^i \delta_l^n - \frac{1}{N}\delta_j^i \delta_m^k \delta_l^n - \frac{1}{N}\delta_j^k \delta_m^n \delta_l^i + \frac{1}{N^2}\delta_j^i \delta_m^n \delta_l^k \quad . \tag{C.34}$$

The normalization of these generators is obtained via

$$(M_j^i)_p^k (M_m^n)_l^p \delta_k^l = (M_j^i)_m^n \quad . \tag{C.35}$$

The interpretation of this relation is obvious once one considers the symmetry of the situation. Above, we normalized our generators according to $\mathrm{Tr}\, t^a t^b = T_F \delta^{ab}$. Here, it is not yet clear what form δ^{ab} will take. In the most fundamental sense, δ^{ab} is simply an invariant tensor with two indices in the adjoint representation. By inspection, one can see that in this representation for the generators adjoint indices consist of traceless pairs of fundamental and anti-fundamental indices. The *only* such invariant object at our disposal is $(M_j^i)_m^n$ itself. Indeed, these two conditions were the reason this form was chosen for M in the first place! Hence we are led to the identification

$$\delta^{ab} \to \left(M_j^i\right)_m^n \quad . \tag{C.36}$$

Proper normalization is obtained by the requirement that

$$N^2 - 1 = (M_j^i)_m^n (M_i^j)_n^m \quad , \tag{C.37}$$

which can be checked explicitly. With this in mind, (C.35) implies that are generators are normalized according to $T_F = 1$.

[9]The fact that I have written both indices downstairs here has no meaning. What is required is the same index structure as the generators have in the representation R.

From here, we are free to calculate any structures we wish. For example,

$$C_F \delta^i_j = \left(M^k_l\right)^i_p \left(M^n_m\right)^p_j \left(M^l_k\right)^m_n = \left(N - \frac{1}{N}\right) \delta^i_j \ . \tag{C.38}$$

Furthermore, the explicit form of the product (C.34) allows a direct calculation of the anti-symmetric

$$\begin{aligned} \left[M^i_l, M^j_m\right] &= i f^{ijk}_{lmn} M^n_k \ ; \\ f^{ijk}_{lmn} &= -i \left(\delta^i_m \delta^j_n \delta^k_l - \delta^k_m \delta^j_l \delta^i_n\right) \end{aligned} \tag{C.39}$$

and symmetric

$$\begin{aligned} \left\{M^i_l, M^j_m\right\} &= \frac{2}{N} \left(M^i_l\right)^j_m + d^{ijk}_{lmn} M^n_k \ ; \\ d^{ijk}_{lmn} &= \delta^i_m \delta^j_n \delta^k_l + \delta^k_m \delta^j_l \delta^i_n \\ &\quad - \frac{2}{N} \left(\delta^i_l \delta^j_n \delta^k_m + \delta^j_m \delta^k_l \delta^i_n + \delta^k_n \delta^i_m \delta^j_l\right) \\ &\quad + \frac{4}{N^2} \delta^i_l \delta^j_m \delta^k_n \ . \end{aligned} \tag{C.40}$$

structure constants. Note that these constants have the stated symmetry properties only under the exchange of *pairs* of indices. Note also that d^{ijk}_{lmn} was *chosen* to have this symmetry. Two of its terms cannot contribute to (C.40). Armed with these expressions for general N, we can directly verify any of the expressions in Appendix A.2. Proper normalization to any value of T_F is trivially performed by rescaling the generators.[10]

As a last topic for this section, let us consider the problem of finding irreducible representations of $SU(N)$. Suppose we wish to consider $(\mathbf{N} \times \mathbf{N})$,

$$T^{ij} \ . \tag{C.41}$$

Since there are no invariants with two fundamental indices, the decomposition of this tensor is trivial. We simply symmetrize and antisymmetrize its indices :

$$T^{ij} = \frac{1}{2}\left(T^{ij} + T^{ji}\right) + \frac{1}{2}\left(T^{ij} - T^{ji}\right) \ . \tag{C.42}$$

The symmetric representation has $N(N-1)/2 + N = N(N+1)/2$ components while the antisymmetric has $N(N-1)/2$. In this way, we arrive at

$$\mathbf{N} \times \mathbf{N} = \frac{\mathbf{N(N+1)}}{\mathbf{2}} + \frac{\mathbf{N(N-1)}}{\mathbf{2}} \ . \tag{C.43}$$

[10] As far as normalization is concerned, f and d count as generators.

Note that we don't need to subtract a trace here; this concept has no meaning in our space. Larger numbers of indices in one representation are handled in an analogous way. One simply adds indices successively to the decomposition in (C.42), symmetrizing and antisymmetrizing as one goes. For example, $(\mathbf{N} \times \mathbf{N} \times \mathbf{N})$ is decomposed as

$$\begin{aligned} T^{ijk} &\sim T^{(ij)k} + T^{[ij]k} \\ &\sim T^{(ijk)} + T^{(i[j]k)} + T^{[i(j]k)} + T^{[ijk]} \ . \end{aligned} \tag{C.44}$$

For $SU(3)$, this decomposition corresponds to

$$\mathbf{3} \times \mathbf{3} \times \mathbf{3} = \mathbf{10} + \mathbf{8} + \mathbf{8} + \mathbf{1} \ . \tag{C.45}$$

This result is relevant to the study of baryons both in $SU(3)_f$, where the **8** corresponds to the octet of spin-1/2 and the **10** corresponds to the decouplet of spin-3/2, and the color $SU(3)$ gauge group, where the **1** is the colorless singlet required for a physical state. In $SU(3)$, the totally antisymmetric combination $T^{[ijk]}$ is obviously invariant since it can be written in terms of ϵ^{ijk}. This correspondence allows us to clearly see the fact that

$$T^{[i_1 \cdots i_{N-1}]} = \epsilon^{i_1 \cdots i_N} \tilde{T}_{i_N} \tag{C.46}$$

transforms as an $\overline{\mathbf{N}}$ rather than an $\mathbf{N}$.

The decomposition of $\mathbf{N} \times \overline{\mathbf{N}}$ is quite different. Here, we have the invariant δ^i_j which leads to a trace subtraction. In this case, it is the concept of symmetrization and antisymmetrization which does not make sense. This leads to the decomposition

$$T^i_j = \left(T^i_j - \frac{1}{N} \delta^i_j \, T^k_k \right) + \frac{1}{N} \delta^i_j \, T^k_k \ . \tag{C.47}$$

The trace is invariant and forms the representation **1** of $SU(N)$, while the remainder transforms in the *adjoint* and forms the $(\mathbf{N^2 - 1})$. Hence

$$\mathbf{N} \times \overline{\mathbf{N}} = \left(\mathbf{N^2 - 1} \right) + \mathbf{1} \ . \tag{C.48}$$

This decomposition is relevant to meson physics, where the $SU(3)_f$ group tells us we will have an octet (the **8**) and a singlet (the **1**) and the $SU(3)$ color group tells us that the combination of a quark and an antiquark can produce a physical color singlet state.

C.2 Local $SU(N)$

As detailed in Chapter 1, QCD is a quantum field theory based on a local $SU(3)$ symmetry. One considers several flavors of quark triplets in the **3** and antiquark triplets in the $\overline{\mathbf{3}}$, and requires the lagrangian to be invariant under local $SU(3)$ transformations. This leads to the introduction of the *covariant* derivative, $\mathcal{D}$. If we just compared the field at two different points (using the ordinary derivative), differences in convention between the two points would lead to physically unimportant contributions. The covariant derivative takes all of these changes in convention into account and asks for the difference between *physical* configurations. However, it can only do this for two infinitesimally close points.

Suppose I want to compare field values at two points which are a finite distance apart. Since these objects are defined with different conventions, we must first change the convention of one of them to match the other convention. This operation is called *parallel transport*. The changes in convention occur locally, so we will need to know the convention at every point on a path between the two points to perform this transformation. The covariant derivative tells us that local changes in convention are accounted for by the gauge field $\mathcal{A}$. Hence we expect parallel transport to involve some sort of nonlocal combination of gauge fields. The infinite number of paths connecting two points in 4-dimensional spacetime make it necessary for us to choose a curve, C, to define a unique parallel transport operator. Defining C by points $z(\tau)$ such that $z(0) = y$ and $z(1) = x$, it can be shown that the operator

$$\mathcal{G}_C(x,y) = \mathrm{P} e^{-ig\int_0^1 A_\mu(z(\tau))(dz^\mu/d\tau)\,d\tau} \tag{C.49}$$

satisfies the requirements for parallel transport from y to x along C. The P-symbol orders the $\mathcal{A}$ fields such that those corresponding to larger τ come first, i.e.[11]

$$\begin{aligned}\mathrm{P}\left\{e^{\int_0^1 M(\tau)d\tau}\right\} \;\equiv\; & 1 + \int_0^1 d\tau M(\tau) + \int_0^1 d\tau M(\tau)\int_0^\tau d\tau' M(\tau') \\ & + \int_0^1 d\tau M(\tau)\int_0^\tau d\tau' M(\tau')\int_0^{\tau'} d\tau'' M(\tau'') + \cdots \quad .\end{aligned} \tag{C.50}$$

It may be checked explicitly for infinitesimal gauge transformations that

$$\mathcal{G}_C(x,y) \to U(x)\mathcal{G}_C(x,y)U^\dagger(y) \tag{C.51}$$

[11] The $1/n!$ in the expansion of an exponential appears naturally here if the path-ordering is trivial since $\int_0^1 d\tau \int_0^\tau d\tau' f(\tau)f(\tau') = 1/2\left(\int_0^1 d\tau f(\tau)\right)^2$ if $f(\tau)$ commutes with itself for different values of τ.

for any choice of the path C. Since all gauge transformations we consider can be obtained from repeated application of infinitesimal ones, this is all we need.[12] Qualitatively, we can think of (C.49) as iteratively taking changes in convention into account. The first term corresponds to no change in convention, the second takes a change in convention into account at one point along C and sums over all possible places the convention could be changed, the third changes conventions twice and sums over the positions of those changes, etc. The path ordering assures us that changes in convention are made in the correct order as we traverse C.

The comparison of field values at two different points in spacetime may now be written

$$(\Delta_C \psi)(x,y) = \psi(x) - \mathcal{G}_C(x,y)\psi(y) \ . \tag{C.52}$$

From the transformation properties of $\mathcal{G}$, it is obvious that $(\Delta\psi)(x,y)$ transforms like $\psi(x)$. Of course, we may define this difference at any point in spacetime by appropriate use of $\mathcal{G}$:

$$(\Delta_{C,C'}\psi)(x,y)\,|_z = \mathcal{G}_{C'}(z,x)\psi(x) - \mathcal{G}_{C+C'}(z,y)\psi(y) \ , \tag{C.53}$$

where C' is some path connecting x and z. The parallel transport operator $\mathcal{G}$ is commonly called a *gauge link* and is very useful in defining nonlocal distributions, as in Section 2.3.

Different paths used to define $\mathcal{G}_C$ are *not* equivalent. The difference between them can be characterized by the *Wilson loop*,

$$\mathcal{U}_\ell(x) \equiv \mathcal{G}_{C'}(x,y)\mathcal{G}_C(y,x) \ . \tag{C.54}$$

The loop ℓ forms a closed curve containing the points x and y and defined by the paths C and C'. Expanding this object about $g = 0$, we see that its terms arrange themselves into path-ordered loop integrals :

$$\begin{aligned}\mathcal{U}_\ell(x) \sim 1 &- ig \oint_\ell \mathcal{A}(z)\cdot dz \\ &-ig^2 f^{abc}t^c \int_1^2 \mathcal{A}^a(z(\tau))\cdot(dz/d\tau)d\tau \int_0^1 \mathcal{A}^b(z(\tau'))\cdot(dz/d\tau')d\tau' \\ &-g^2 \int_0^2 \mathcal{A}(z(\tau))\cdot(dz/d\tau)d\tau \int_0^\tau \mathcal{A}(z(\tau'))\cdot(dz/d\tau')d\tau' + \mathcal{O}(g^3) \ .\end{aligned} \tag{C.55}$$

[12] As stated before, other transformations can be accounted for by rotations in the other sectors of the SM. Topologically nontrivial gauge configurations are associated with behavior at infinity and do not concern us here.

Here, $z^\mu(\tau)$ traces out C for $\tau \in [0,1)$ and C' for $\tau \in (1,2]$.

In abelian theories, one can use Stokes' theorem to convert these integrals into surface integrals over the area ℓ bounds. The same spirit is present in non-abelian theories, although the difficulties associated with noncommutativity make it less transparent. To see this, we consider an infinitesimal loop. Taking $y = x + dx^\mu + dx^\nu$, where μ and ν are for the moment *fixed* directions, we can evaluate the difference in choice between a path that goes first along dx^μ then along dx^ν and a path that does the opposite by approximating the integrals :

$$\begin{aligned} \oint_\ell \mathcal{A}(z)\cdot dz \quad &\sim \quad \mathcal{A}^\mu(x)dx_\mu + \mathcal{A}^\nu(x+dx^\mu)dx_\nu \\ &\quad -\mathcal{A}^\nu(x)dx_\nu - \mathcal{A}^\mu(x+dx^\nu)dx_\mu \qquad \text{(C.56)} \\ &= \quad (\partial^\mu \mathcal{A}^\nu - \partial^\nu \mathcal{A}^\mu)\, dx_\mu dx_\nu \ , \qquad \text{(C.57)} \end{aligned}$$

where I have suspended the Einstein summation convention for the moment. Similarly, the non-abelian commutator term in (C.55) is approximated as

$$\begin{aligned} -ig^2 f^{abc} t^c \int_1^2 \mathcal{A}^a(z(\tau))\cdot(dz/d\tau)d\tau \int_0^1 \mathcal{A}^b(z(\tau'))\cdot(dz/d\tau')d\tau' \\ \sim ig^2 f^{abc} t^c \mathcal{A}^\nu_a(x)\mathcal{A}^\mu_b(x)\, dx^\nu dx^\mu \ . \qquad \text{(C.58)} \end{aligned}$$

Combining these two contributions, the integral is expressed

$$-ig\mathcal{F}^{\mu\nu}(x)\, dx_\mu dx_\nu \ . \qquad \text{(C.59)}$$

The antisymmetric combination of the differentials gives an infinitesimal area element - the area of the bounded region. This statement is valid for any choices of μ and ν, allowing us to apply it to the discretization of *any* spacetime region bounded by curves C and C'. Furthermore, this analysis can be extended to higher orders in the coupling.

The fact that $\mathcal{F}$ itself is not gauge invariant makes it difficult to compare infinitesimal contributions from different regions. However, the gauge-invariant *trace* of $\mathcal{U}$ admits a full generalization of Stokes' theorem :

$$\mathrm{Tr}\, \mathcal{U}_\ell = \mathrm{Tr}\, e^{-ig\int_S \mathcal{F}^{\mu\nu}\, d\sigma_{\mu\nu}} \ , \qquad \text{(C.60)}$$

where S is any surface bounded by ℓ and $d\sigma^{\mu\nu}$ is the four-dimensional area element. This object can be used as a *gauge-invariant* measure of the strength of the gauge field in a region. It is also one of the fundamental building blocks of the discretized theory of lattice QCD.

C.3 The Limit of Large N_c

Some very useful results have come from considering a local gauge theory based on the group $SU(N_c)$ in the limit of large N_c. In this section, I will briefly overview some of the main results and simplifying features of this limit. Some useful references on this subject are [92].

The lagrangian of large-N_c QCD is identical to that of normal QCD, except that here there are N_c quark colors and $(N_c^2 - 1)$ gluons. As we have seen above, adjoint objects such as gluons can be represented as the traceless combination of fundamental and an anti-fundamental indices. Since we intend to take N_c to infinity, the fact that the trace degree of freedom should be subtracted out is of no great importance. Hence, with corrections of order $1/N_c^2$, we can consider a gluon as a quark-antiquark combination.[13] The replacement of all gluon propagators with double quark propagators, with fermion number flow in opposite directions, makes it easy to count powers of N_c. Each unconstrained loop gives a factor of N_c when its colors are summed over. For example, Figure C.1a comes with the factor N_c^2 because there are two unconstrained 'fermion' loops. Unless we can regulate this behavior in some way, it quickly becomes obvious that each higher order of perturbation theory will contain diagrams with more powers of N_c. This is certainly not the hallmark of a theory with a well-defined limit as $N_c \to \infty$.


By inspection, one can see that factors of N_c are *always* accompanied with factors of $g_{N_c}^2$, where g_{N_c} is the coupling constant of our theory. Since this is *our* theory, we can force g to scale with N_c in any way we want. Comparison with the real world requires only that g_3 has the value measured in experiment. With this in mind, we see that our problem can be solved by taking

$$g_{N_c} \equiv \frac{g}{\sqrt{N_c}} \quad . \tag{C.61}$$


This definition causes all of the *leading* powers of N_c at each order in perturbation theory to contribute at the *same* power of N_c to the final amplitude under consideration.

Now that we have a well-defined limit, it behooves us to find out which diagrams contribute in this limit and which do not. Since each power of the coupling contributes a $1/\sqrt{N_c}$ to the amplitude, leading diagrams have the property that each coupling generates 'half' of an


[13] This is to be understood as taking place exclusively in the $SU(N_c)$ sector of the theory.



(a)



(b)



(c)

Figure C.1: An illustration of the large-N_c counting rules. (a) is enhanced by a power of N_c^2 because of the unconstrained color loops within the bubble. Since it is a two-loop graph, it will contribute to the leading large-N_c behavior. On the other hand, (b) contains only one unconstrained color loop. As a three-loop diagram, its contribution comes only at $\mathcal{O}(1/N_c^2)$. Notice that (a) is a planar diagram, while (b) must contain a self-intersection when forced into a plane. As it stands, the counting rules for more complicated correlation functions, like that represented in (c), are ambiguous. At first, we may think that it is planar. However, drawing all external lines *directly* to the open dot (representing infinity), reveals its true nature.

unconstrained color loop. Real quarks have only one color line, so quark loops cannot give rise to factors of N_c even though they *are* associated with factors of $g^2_{N_c}$. This implies that *all* diagrams that involve quark loops are suppressed by N_c. This is easy to understand because there are simply more gluons around than quarks. Gluons[14] will not always lead to the maximal power of N_c either. By inspection, it can easily be seen that diagrams like that in Figure C.1b in which the color loops are entangled are subleading. This can be extended to a rule - all diagrams which possess self-intersecting color lines when embedded in a plane are subleading in the large-N_c limit. Since these diagrams cannot be topologically embedded in a plane, they are referred to as 'nonplanar'. This rule works for loops, but must be amended to include correlation functions with several external particles. Here, there exist diagrams which can be drawn such that they appear either planar or nonplanar depending on preference. However, when all external lines are drawn *directly* to the *same* point at infinity, only the planar diagrams will contribute to the leading large-N_c behavior. This rule is illustrated in Figure C.1c.

Beyond our rule for leading diagrammatic contributions, not many results of large-N_c physics affect the topics presented here. However, in hadronic physics this limit represents an extremely useful simplification. Among other things, it constrains meson-baryon couplings via uniterity relations, allows one to justify certain hadronic models, like the Skyrme model, and makes it easier to relate the macroscopic world of hadrons to the microscopic one of QCD. A new symmetry between the light quark flavor and spin becomes manifest in this limit. This 'spin-flavor' symmetry arises because baryons composed of $N_c \to \infty$ quarks do not care if you flip the spin of one or two. This leads to a symmetry between the low-lying baryonic spin excitations [93]. In particular, it relates the Δ resonance to the nucleon. Corrections to this result can also be systematically calculated, a situation hard to come by in hadronic physics. Furthermore, an even larger symmetry group encompasses the baryons if we simultaneously take some of the quark masses to infinity [94].

[14] In covariant gauges, one must also include ghosts. Since they appear as adjoints along with the gluons, their contributions can also give leading effects.

Appendix D

Quantum Mechanics

The fundamental postulate of quantum mechanics,

$$[x, p] = i\hbar \ , \tag{D.1}$$

is interpreted in quantum field theory as a statement about the commutator of a field operator with its conjugate momentum. The process known in the sixties as 'second quantization' involves promoting the wavefunctions of quantum mechanics to operators and imposing the 'canonical commutator' (D.1) on these wavefunctions and their conjugate momenta. The purpose of this appendix is to quickly review this process and obtain explicit expressions for the generators of the Poincaré group in a quantum field theory.

The analysis of Section D.2 closely follows that in S. Weinberg [88].

D.1 Canonical Quantization

In ordinary quantum mechanics, the Schrödinger equation,

$$i\hbar \frac{\partial}{\partial t}\phi(\vec{x}\,, t) = -\frac{\hbar^2}{2m}\nabla^2\phi(\vec{x}\,, t) + V(\vec{x}\,)\phi(\vec{x}\,, t) \tag{D.2}$$

is just the statement that

$$E = \frac{\vec{p}^{\,2}}{2m} + V(\vec{x}\,) \ , \tag{D.3}$$

along with the added understanding that the quantities E and $\vec{p}$ are operators on the function space that satisfy the commutation relations

$$[x^\mu, p^\nu] = -i\hbar g^{\mu\nu} \ . \tag{D.4}$$

Equation (D.2) follows from the solution

$$p^\mu = i\partial^\mu \tag{D.5}$$

of (D.4).

Unfortunately, a direct application of (D.5) to the relativistic version of (D.3),

$$p^2 = m^2 \ , \tag{D.6}$$

leads to a Hamiltonian whose spectrum is not bounded from below. Hence our theory does not possess any stable configurations. The solution to this dilemma has to do with interpretation. In non-relativistic quantum mechanics, the dynamical variables are $\vec{x}$ and t. One requires ϕ to be an eigenfunction of the energy operator, which depends on the position $\vec{x}$. In quantum field theory, the dynamical variables are the fields ϕ themselves. Since t is no longer a dynamical variable, we no longer have any need to bound its Fourier conjugate from below. This is the fundamental distinction between non-relativistic quantum mechanics and quantum field theory.

The dynamical variables of *any* quantum system satisfy canonical commutation relations with their conjugate momenta. In order to define this statement, we must introduce a lagrangian for our system. A lagrangian is dictated by the equations of motion it generates, so the first question we must ask ourselves is what kind of equation of motion we would like our fields to satisfy. At some point, we intend to consider freely propagating massive particles, so a logical choice for the equation of motion is (D.6),

$$\left(-\partial^2 - m^2\right)\phi = 0 \ . \tag{D.7}$$

This equation can be seen to follow from the classical lagrangian density[1]

$$\mathcal{L} = \frac{1}{2}\partial^\mu\phi\partial_\mu\phi - \frac{1}{2}m^2\phi^2 \ , \tag{D.8}$$

from which we obtain the conjugate momentum

$$\pi = \frac{\delta\mathcal{L}}{\delta\partial_0\phi} = \partial^0\phi \ . \tag{D.9}$$

[1]The method of going from lagrangian densities to equations of motion is discussed in the next section.

The canonical commutation relation[2]

$$[\phi(x), \pi(y)]\Big|_{x^0=y^0} = i\delta(\vec{x} - \vec{y}) \tag{D.10}$$

can be solved by expanding $\phi(x)$ in a complete basis of solutions to (D.7) with operator coefficients :[3]

$$\phi(x) = \int \frac{d^4p}{(2\pi)^4} \left[a_{\vec{p}}\, e^{-ip\cdot x} + a^\dagger_{\vec{p}}\, e^{ip\cdot x}\right] \sqrt{2p^0}\Theta(p^0)(2\pi)\delta\left(p^2 - m^2\right) \quad . \tag{D.11}$$

Consistency with (D.10) requires

$$\left[a_{\vec{p}}, a^\dagger_{\vec{p}'}\right] = (2\pi)^3\delta\left(\vec{p} - \vec{p}'\right) \quad . \tag{D.12}$$

Several comments are in order here. First, the canonical commutation relation (D.10) must be imposed only at equal times; any other choice would violate causality.[4] The presence of the δ-function is understood immediately once we realize that we are quantizing a continuously infinite number of field *densities* $\phi(\vec{x})$. Our commutation relations must be such that the integral over a small region about $\vec{x}$ gives one. Second, the form of equation (D.11) is dictated by the reality of $\phi(x)$. Complex fields will in general have different operators multiplying their positive- and negative-frequency solutions. Note that I have included the negative frequency solutions with the positive ones by changing the sign in the exponential.

The field ϕ, along with

$$\pi(x) = -i\int \frac{d^4p}{(2\pi)^4} \left[a_{\vec{p}}\, e^{-ip\cdot x} - a^\dagger_{\vec{p}}\, e^{ip\cdot x}\right] p^0\sqrt{2p^0}\Theta(p^0)(2\pi)\delta\left(p^2 - m^2\right) \quad , \tag{D.13}$$

can be used to form the *true* Hamiltonian of the system,

$$\begin{aligned} H &= \int d^3x \left[\pi(x)\partial^0\phi(x) - \mathcal{L}\right] \\ &= \int \frac{d^3p}{(2\pi)^3}\sqrt{\vec{p}^2 + m^2}\left[a^\dagger_{\vec{p}}\, a_{\vec{p}} + \frac{1}{2}(2\pi)^3\delta^{(3)}(0)\right] \quad . \end{aligned} \tag{D.14}$$

[2] $\hbar$ was reinstated at the beginning of the Appendix for nostalgic purposes; from now on, it will be given the value 1.

[3] The hermiticity of ϕ relates the two operator coefficients, as shown. The factor of $\sqrt{2p^0}$ is for later convenience. It belongs to the operators $a_{\vec{p}}$ and $a^\dagger_{\vec{p}}$, as the combination is invariant.

[4] Actually, this statement is not strictly true. Lorentz invariance implies that we can impose this relation for equal coordinate along any *timelike* vector we so choose. In the limit of large boosts, we can even quantize on the light cone. However, once we have chosen a direction and imposed an equal 'time' commutation relation, commutation relations at different times are fixed and can no longer be imposed.

The second term in this expression represents the sum of the zero-point energy of a continuously infinite number of quantum harmonic oscillators. This contribution to the energy is completely independent of the state we consider, and as such cannot contribute to any physical energy *differences*. For this reason, we will ignore it completely. The rest of the Hamiltonian involves the *energy* of a particle of momentum $\vec{p}$,

$$E_{\vec{p}} \equiv \sqrt{\vec{p}^{\,2} + m^2} \ , \tag{D.15}$$

and satisfies

$$\left[H, a_{\vec{p}}\right] = -E_{\vec{p}}\, a_{\vec{p}} \tag{D.16}$$

$$\left[H, a^\dagger_{\vec{p}}\right] = \ \ E_{\vec{p}}\, a^\dagger_{\vec{p}} \ . \tag{D.17}$$

Assuming a *ground state* $|0\rangle$ that satisfies

$$a_{\vec{p}}\,|0\rangle = 0 \ , \tag{D.18}$$

we can construct a whole tower of states

$$|n_1\vec{p}_1, n_2\vec{p}_2, \cdots, n_N\vec{p}_N\rangle \equiv \prod_{i=1}^{N} (2E_{\vec{p}_i})^{n_i/2}\, \frac{\left(a^\dagger_{\vec{p}_i}\right)^{n_i}}{\sqrt{n_i!}}\,|0\rangle \tag{D.19}$$

which have the *positive* energy

$$H\,|n_1\vec{p}_1, n_2\vec{p}_2, \cdots, n_N\vec{p}_N\rangle = \sum_{i=1}^{N} n_i E_{\vec{p}_i}\,|n_1\vec{p}_1, n_2\vec{p}_2, \cdots, n_N\vec{p}_N\rangle \ . \tag{D.20}$$

Our states are constructed to have the covariant normalization

$$\langle\vec{p}\,|\vec{p}\,'\rangle = 2E_{\vec{p}}(2\pi)^3\,\delta\,(\,\vec{p} - \,\vec{p}\,') \tag{D.21}$$

for each particle.[5]

These states have exactly the properties we expect of a state of n_i free particles of mass m and momentum $\vec{p}_i$. Apparently, we are encouraged to interpret the operators $a^\dagger_{\vec{p}}$ and $a_{\vec{p}}$ as *creating* and *destroying* a particle, or *field quantum*, of momentum $\vec{p}$. Quite unexpectedly, we have come up with a theory which does *not* have a fixed particle content. This fact, which

[5]For states with more than one particle of the same momentum, this normalization will lead to divergent δ-functions. However, any truly physical process will involve wavepackets rather than real momentum eigenstates. The averaging involved here will smear the δ-function.

stems *directly* from the change in wavefunction interpretation from innocent state vector to dynamical variable, represents a major shift in the way we view physics as a whole.

The field ϕ, known as the *Klein-Gordon* [95] field, is a scalar field because it is invariant under the action of the Lorentz group. Although this makes it a good starting place for a discussion on the quantization of field theories, its appearence in physical theories is rare at best.[6] Physical theories involve spinor and vector fields almost exclusively. For this reason, we now turn to the Dirac field.

According to the presentation of Section B.3, Dirac fields satisfy the equation of motion

$$\not{p}\,\psi(x) = i\not{\partial}\,\psi(x) = m\psi(x) \quad . \tag{D.22}$$

This equation follows from the lagrangian density

$$\mathcal{L} = \overline{\psi}\,(i\not{\partial} - m)\,\psi \quad , \tag{D.23}$$

from which we can deduce the momentum conjugate to ψ :

$$\frac{\delta\mathcal{L}}{\delta\partial_0\psi} = i\overline{\psi}\gamma^0 = i\psi^\dagger \quad . \tag{D.24}$$

Anticipating a field quantum for ψ, we use the complete basis of solutions to (D.22) found in Section B.3 to expand

$$\psi(x) = \int \frac{d^4p}{(2\pi)^4} \sum_s \left[b_{\vec{p},s} u(p,s) e^{-ip\cdot x} + d^\dagger_{\vec{p},s} v(p,s) e^{ip\cdot x} \right] \sqrt{2p^0}\Theta(p^0)(2\pi)\delta\left(p^2 - m^2\right) \quad . \tag{D.25}$$

We hope to identify the coefficients in this expansion with particle creation and annihilation operators.

The Dirac field is *not* hermitian, so we have no right to assume a relation between the creation and annihilation operators. However, as $e^{-ip\cdot x}(e^{+ip\cdot x})$ carries positive (negative) energy[7] $p^0(-p^0)$, we associate it with an annihilation (creation) operator.[8] With this in mind,

[6] The closest friend of the real Klein-Gordon field present in the Standard Model is the complex Higgs field. This is the only field in the Standard Model whose quanta have not yet (2000) been discovered experimentally (unless, of course, one counts the longitudinal modes of the massive gauge bosons...).

[7] Here, I mean energy in the sense of a wave, rather than energy in the sense of the Hamiltonian.

[8] When ψ acts on a physical state, we want the creation of a particle to take free energy away from the system and the destruction of a particle to give free energy to the system. In this sense, a system must use its wavelike (free) energy to 'make' a particle (bound energy).

we assume the commutators

$$\left[b_{\vec{p},s}, b^\dagger_{\vec{p}\,',s'}\right] = \delta_{ss'}(2\pi)^3\delta(\vec{p}-\vec{p}\,') = \left[d_{\vec{p},s}, d^\dagger_{\vec{p}\,',s'}\right] \quad , \tag{D.26}$$

with all others vanishing,[9] and check the canonical commutator

$$\left[\psi(x), \psi^\dagger(y)\right]\Big|_{x^0=y^0} = \delta(\vec{x}-\vec{y}) \ . \tag{D.27}$$

Unfortunately, our choice (D.26) does not cause this relation to be satisfied.[10] The culprit is the lack of sign between the positive- and negative-frequency components of

$$\psi^\dagger(x) = \int \frac{d^4p}{(2\pi)^4} \sum_s \left[b^\dagger_{\vec{p},s}\, u^\dagger(p,s)e^{ip\cdot x} + d_{\vec{p},s}\, v^\dagger(p,s)e^{-ip\cdot x}\right] \sqrt{2p^0}\Theta(p^0)(2\pi)\delta\left(p^2-m^2\right) \ . \tag{D.28}$$

This stems from the fact that there is only one derivative in the lagrangian density, (D.23), which is traced in turn to the form the translation operator takes in $SL(2,C)$. Hence this seems to be a fundamental property of spinor fields.

One solution to this problem is to simply generate the sign from the commutation relations. If we were to take

$$\left[d_{\vec{p},s}, d^\dagger_{\vec{p}\,',s'}\right] = -\delta_{ss'}(2\pi)^3\delta(\vec{p}-\vec{p}\,') \tag{D.29}$$

rather than (D.26), the fields would indeed satisfy the commutation relation (D.27). However, this solution is exactly as artificial as it seems. It only prolongs our suffering as it leads to a Hamiltonian the is not bounded from below. This fact is intuitively obvious. A commutation relation like (D.29) gives $d_{\vec{p},s}$ the role of creation operator. Hence the creation of a d-type quantum *gives* free energy to a state. This implies that all of our states will eventually *decay* to the state of *infinite* d quanta. Our aversion to this state of affairs was what led us to consider the fields as dynamical variables in the first place, so this 'solution' to our problem is clearly unacceptable.

Alternatively, we can *demand* that $d_{\vec{p},s}$ destroy quanta and search for other sources of the sign. To this end, we consider a state $|0\rangle$ that both b and d annihilate. We would also like $b^\dagger$ and $d^\dagger$ to create quanta, so we require that the state $dd^\dagger|0\rangle$ is proportional to $|0\rangle$. If the

[9] In the free theory, these are all uncoupled harmonic oscillators; operators from different oscillators should commute.

[10] One can check this directly using the completeness relations (B.122) and (B.156).

commutator $[d, d^\dagger] = 1$, then we arrive at the old result $dd^\dagger|0\rangle = |0\rangle$. However, the reason these two quantities are equal is that d annihilates $|0\rangle$. In a fundamental sense, it doesn't matter whether we *subtract* $d^\dagger d$ to form the *commutator* or *add* it to form the *anticommutator*. On the other hand, this difference is essential to the cancelation of Dirac structures in the canonical commutator. If we choose instead to impose a canonical *anti*commutator, we will find that

$$\left\{ b_{\vec{p},s}, b^\dagger_{\vec{p}',s'} \right\} = \delta_{ss'}(2\pi)^3\delta(\vec{p}-\vec{p}') = \left\{ d_{\vec{p},s}, d^\dagger_{\vec{p}',s'} \right\} \quad , \tag{D.30}$$

with all other anticommutators zero, satisfy all of our requirements without compromising the integrity of our vacuum.

The decision to impose canonical anticommutation relations on spinor fields is not extremely outrageous. As we have seen, these fields are *fundamentally* different from scalar and vector fields. In fact, looking at the classical fields, we would actually *expect* this behavior. Classical vector fields satisfy

$$\mathcal{A}^\mu g_{\mu\nu} \mathcal{B}^\nu = \mathcal{B}^\nu g_{\nu\mu} \mathcal{A}^\mu \quad , \tag{D.31}$$

while for spinor fields we have

$$\xi^\alpha \epsilon_{\alpha\beta} \eta^\beta = -\eta^\beta \epsilon_{\beta\alpha} \xi^\alpha \quad . \tag{D.32}$$

Hence in classical theories the vector degrees of freedom commute, while spinor degrees anticommute. From this point of view, it seems very natural that quantization imposes a commutator on vectors and an anticommutator on spinors. This argument also implies that *different* spinor fields should *anticommute* rather than commute, exactly as we found above.

Whatever the reason, the imposition of anticommutators on fermionic fields changes the theory drastically. In particular, the statement

$$\left\{ b^\dagger_{\vec{p},s}, b^\dagger_{\vec{p},s} \right\} = 0 \tag{D.33}$$

implies that the creation operator annihilates the one-particle state. Spinor states are either filled or empty; there can be no multiple occupation. This idea was first postulated by Pauli as a way to explain the observed atomic spectra. In non-relativistic quantum mechanics, it is imposed on the theory as an experimental input. In quantum field theory, it is promoted to a *consistency requirement*. This is one of the major triumphs of relativistic quantum mechanics. Another is the prediction of antiparticles. Since the quanta of the d operator are created by

ϕ while those of the b operator are destroyed, the absence of a 'b-on' is equivalent in some sense to the presence of a 'd-on'. This relationship is dubbed particle-antiparticle duality. By convention, we take the quanta destroyed by ψ as the particles, while those it creates the antiparticles. Thus the quark field ψ_q destroys quarks and creates antiquarks. Antiparticles were first predicted by P. A. M. Dirac [96] in 1931, and discovered in the form of the positron by C. D. Anderson in 1932 [97]. Dirac shared the 1933 Nobel prize in physics with E. Schrödinger for this advancement of the quantum theory, and Anderson won the Nobel Prize in 1936 for its experimental verification.

The Hamiltonian of the (correctly quantized) free Dirac field can be guessed from (D.20); it is a weighted sum over the 'occupation number' operators $b^\dagger b$ and $d^\dagger d$. Our particle states

$$|\,\vec{p},s\rangle = \sqrt{2E_{\vec{p}}}\,b^\dagger_{\vec{p},s}\,|0\rangle \tag{D.34}$$

are normalized such that

$$\big\langle\,\vec{p}\,',s'\big|\,\vec{p},s\big\rangle = 2E_{\vec{p}}\,\delta_{ss'}(2\pi)^3\delta(\,\vec{p}\,'-\vec{p}\,)\ . \tag{D.35}$$

Antiparticle states are normalized in the same way. Note that the action of our fields gives

$$\psi^\dagger(x)|0\rangle \;=\; \int\frac{d^3p}{(2\pi)^3}\,\frac{1}{2E_{\vec{p}}}\,\sum_s u^\dagger(p,s)e^{ip\cdot x}\,|\,\vec{p},s\rangle\ , \tag{D.36}$$

$$\psi(x)|\,\vec{p},s\rangle \;=\; u(p,s)e^{-ip\cdot x}|0\rangle+\cdots\ , \tag{D.37}$$

so $\psi^\dagger(x)$ creates a particle at spacetime point x and ψ changes the one-particle state into the vacuum while providing a wavefunction and some free energy.

Canonical quantization of ordinary vector fields goes through in the same way as for scalars. The only difference is the existence of a polarization vector ε^μ analogous to the wavefunction $u(p,s)$ for fermions. Gauge bosons, on the other hand, are much more nontrivial to quantize using the method presented above. In fact, quantization of non-abelian gauge bosons in covariant gauges requires the introduction of ghost fields, as discussed in Section 1.2. For these kinds of theories, it is much clearer to use the path integral approach presented there.

D.2 Spacetime Symmetries - The Generators of the Poincaré Group

In this section, we would like to study the implications of Poincaré symmetry on a quantum field theory. In all of the following, we will assume that the action relevant to our theory is invariant under Poincaré transformations of the fields, i.e.

$$\mathcal{S}[\Phi', (\partial_\mu \Phi)'] = \mathcal{S}[\Phi, \partial_\mu \Phi] \tag{D.38}$$

for any set of fields Φ' related to Φ by a Poincaré transformation. This is true of any theory whose lagrangian is a Lorentz scalar and does not depend explicitly on the spacetime coordinates x^μ.

In a classical theory, we expect such symmetries to lead to currents and charges which are conserved by physical processes. As mentioned in Section B.2, invariance under translation leads to conservation of momentum and that under Lorentz transformations leads to conservation of angular momentum. Let us see how this comes about. To begin with, we must define what we mean by a physical process. Classically, a physical process is one which makes the action stationary. Under an arbitrary infinitesimal variation of the fields, the action changes by the amount[11]

$$\begin{aligned} \delta\mathcal{S} &= \int d^4x \left[\frac{\delta\mathcal{L}}{\delta\Phi}\delta\Phi + \frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\delta\partial_\mu\Phi \right] && \text{(D.39)} \\ &= \int d^4x \left[\frac{\delta\mathcal{L}}{\delta\Phi}\delta\Phi + \frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\partial_\mu\delta\Phi \right] && \text{(D.40)} \\ &= \int d^4x \left[\frac{\delta\mathcal{L}}{\delta\Phi} - \partial_\mu\frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi} \right] \delta\Phi \ . && \text{(D.41)} \end{aligned}$$

In the last line, I have ignored a surface term. This contribution does become important in certain theories, most notably gravitational theories and topological models, but it will not

[11] These manipulations can be thought of in the following way. We first consider the action as a function of the *independent* fields Φ and Φ_μ. No one can prevent us from varying these fields independently. However, in the end we are interested only in the class of fields that satisfy the constraint $\Phi_\mu = \partial_\mu\Phi$. If we make independent variations, this constraint will not be preserved. On the other hand, if we correlate the field variations by imposing $\delta\Phi_\mu = \partial_\mu\delta\Phi$, field configurations which satisfy $\Phi_\mu = \partial_\mu\Phi$ will continue to satisfy it after the transformation. In this way, we can differentiate the lagrangian with respect to the fields Φ and $\partial_\mu\Phi$ as though they were independent and still obtain the proper constrained variations. Of course, this procedure is equivalent to the more orthodox method of Lagrange multipliers.

bother us here. From the last line, it is apparent that the requirement of a stationary action for *arbitrary* infinitesimal field variations implies the Euler-Lagrange equation

$$\frac{\delta \mathcal{L}}{\delta \Phi} - \partial_\mu \frac{\delta \mathcal{L}}{\delta \partial_\mu \Phi} = 0 \ . \tag{D.42}$$

This expression is know as the equation of motion of the theory. Each independent field in the theory has an associated equation of motion which is satisfied by all physical processes.

If we now specialize our variation to a translation of spacetime coordinates, we can work out the implications of translation invariance. Under a translation, we have the variations[12]

$$\begin{aligned} \delta\Phi(x) &= \epsilon^\mu(x)\partial_\mu\Phi(x) \\ \delta\partial_\mu\Phi(x) &= \partial_\mu\epsilon^\nu(x)\partial_\nu\Phi(x) \ . \end{aligned} \tag{D.43}$$

Substituting these expressions into (D.39), we see that $\mathcal{S}$ changes by the amount

$$\delta\mathcal{S} = \int d^4x \left[\epsilon^\nu(x)\left(\frac{\delta\mathcal{L}}{\delta\Phi}\partial_\nu\Phi + \frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\partial_\nu\partial_\mu\Phi\right) + \left(\frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\partial_\nu\Phi\right)\partial_\mu\epsilon^\nu(x)\right] \ . \tag{D.44}$$

The quantity multiplying ϵ^ν combines to form $\partial_\nu\mathcal{L}$, as can be seen via the chain rule, so we are left[13] with

$$\delta\mathcal{S} = \int d^4x\, \partial_\mu\epsilon_\nu(x)\left[\frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\partial^\nu\Phi - g^{\mu\nu}\mathcal{L}\right] \ . \tag{D.45}$$

Calling the quantity in brackets $\mathcal{T}^{\mu\nu}$, we see that invariance of the action under translation implies that

$$\partial_\mu\mathcal{T}^{\mu\nu} = 0 \ . \tag{D.46}$$

One often refers to this expression as the 'conservation' of $\mathcal{T}^{\mu\nu}$ since it implies that the vector $\int d^3\vec{x}\,\mathcal{T}^{0\nu}(x)$ does not depend on time :

$$\begin{aligned} \frac{d}{dt}\int d^3\vec{x}\,\mathcal{T}^{0\nu}(x) &= \int d^3\vec{x}\,\frac{\partial}{\partial x^0}\mathcal{T}^{0\nu}(x) \\ &= -\int d^3\vec{x}\,\frac{\partial}{\partial x^i}\mathcal{T}^{i\nu}(x) \\ &= -\oint dS_i\,\mathcal{T}^{i\nu}(x) \\ &= 0 \ . \end{aligned} \tag{D.47}$$

[12]The fact that the Pauli-Ljubanskii vector commutes with the generators of translation implies that this operation will not mix different elements of the same representation of the Poincaré group. In other words, we can translate each field independently. This will not be true of Lorentz rotations.

[13] after ignoring another surface term

The first equality requires the volume of integration to be independent of time. In the second, we have used the fact that the divergence of $\mathcal{T}$ is zero. The third is an application of the divergence theorem. The integration takes place on the boundary of the volume and dS_i represents an outward unit normal. In the fourth, we have assumed that this surface integral vanishes. If it does not, Equation (D.47) states that the change in this quantity with time is due only to the value of $\mathcal{T}^{i\nu}$ on the boundary. Physically, we can think of $\mathcal{T}^{i\nu}$ as the 'flow' of a 'charge density' $\mathcal{T}^{0\nu}$. This interpretation makes it easy to understand (D.47). It says that the change in total charge in a volume is equal to the amount that came in across the boundary. This is conservation in its most fundamental form. Since the conserved 'charge' here is due to translation invariance, we will call it momentum. Thus $\mathcal{T}^{0\nu}$ can be identified with a momentum density and $\mathcal{T}^{i\nu}$ with a momentum flow. The integrated charge, $\mathcal{P}^\mu \equiv \int d^3x \mathcal{T}^{0\mu}(x)$, is the conserved total momentum of the system. For these reasons, $\mathcal{T}$ is called the *canonical energy-momentum tensor* of our theory.

However nice this derivation is, it carries with a few problems. First, it is not explicitly covariant. This problem stems from the fact that we must choose a frame before we ask if a quantity changes with time. However, looking at the derivation, we certainly never used any specific features of the frame we were in. In fact, the whole derivation can be performed with any timelike vector v^μ replacing the coordinate t. The statement then becomes

$$\frac{d}{d\lambda}\int \epsilon_{\alpha\beta\gamma\delta}\, d\Sigma^{\alpha\beta\gamma} v^\delta v^*_\mu \mathcal{T}^{\mu\nu} = -\oint \epsilon_{\alpha\beta\gamma\delta}\, d\sigma^{\alpha\beta} v^\gamma \mathcal{T}^{\delta\nu} \quad , \tag{D.48}$$

where $d\lambda$ represents a change along the v^μ direction, $v^{*\,\mu}$ is a vector complementing v such that $v^* \cdot v = 1$, $d\Sigma^{\mu\nu\alpha}$ and $d\sigma^{\mu\nu}$ represent volume and area integration measures,[14] respectively, and the integration region cannot change as one traverses the v^μ direction. The physical content of this equation is identical to that of (D.47). In fact, due to the covariance of the result, we can always choose a frame in which $v^\mu = v^{*\,\mu} = (1, 0, 0, 0)$ and use (D.47).[15]

A more fundamental difficulty with this tensor is that it is not always symmetric. It turns out that defining a conserved angular momentum density requires a symmetric energy-

[14] These quantities are best described in the language of 3- and 2-forms, respectively. In both cases, the orientation of integration is contained in their definition. Here, the orientation of the volume integration must also be specified. It is timelike.

[15] unless we choose v^μ lightlike, as is done in light-cone perturbation theory [52].

momentum tensor. Non-relativistically, we would expect the angular momentum density to have the form

$$\vec{L} = \vec{x} \times \vec{P} \ . \tag{D.49}$$

Generalizing this equation to four dimensions, we write

$$J^{\mu\nu} = x^\mu P^\nu - x^\nu P^\mu \ . \tag{D.50}$$

Interpreting P as a momentum density, we see that this is just the time component of the tensor

$$\mathcal{M}^{\alpha\mu\nu} = x^\mu \mathcal{T}^{\alpha\nu} - x^\nu \mathcal{T}^{\alpha\mu} \ . \tag{D.51}$$

Taking the divergence of this object on α, we find that

$$\partial_\alpha \mathcal{M}^{\alpha\mu\nu} = \mathcal{T}^{\mu\nu} - \mathcal{T}^{\nu\mu} \ , \tag{D.52}$$

effectively ending all of our dreams of a conserved angular momentum.

This is not necessarily a failure. We have not yet told our calculations that the lagrangian for our theory is Lorentz invariant, and we cannot expect to get angular momentum conservation for free. If $\mathcal{T}^{\mu\nu}$ were manifestly symmetric, we would have the result that translationally invariant theories are automatically Lorentz invariant and possess conserved angular momenta. Lorentz invariant scalar theories do indeed possess symmetric canonical energy-momentum tensors. This is due to the simple fact that no antisymmetric two-index structure can be constructed in these theories.[16] However, somewhat more input is required in theories involving fields with nonzero intrinsic angular momentum.

The biggest problem with $\mathcal{T}^{\mu\nu}$ occurs in gauge theories like QCD : it is not gauge invariant. This fact destroys any hope we ever had of interpreting any of its components as physical objects. $\mathcal{T}$ may be conserved in a gauge theory, but it makes no sense. It seems that this devastating blow renders all of our work useless. However, all is not lost. Above, we argued that the canonical momentum density cannot be used to locally construct a conserved angular momentum density for fields with nontrivial spin. We have also seen that imposing local gauge invariance requires the introduction of a gauge field, which necessarily transforms like a

[16] The structure $\partial^\mu \phi \, \partial^\nu \chi$ can always be transformed away by defining new fields.

vector[17] since it is on the same footing as ∂_μ. Hence the inconsistency of the canonical energy-momentum tensor in any gauge theory is already expected on angular momentum grounds. This suggests that the gauge dependent part of $\mathcal{T}$ is related to its antisymmetric part.

If our theory truly is Lorentz invariant, we must be able to define a conserved angular momentum. There are several ways to do this. A straightforward approach is simply to substitute the infinitesimal form of a spacetime dependent Lorentz transformation into Eq.(D.39) and follow one's nose. This will certainly lead to a conserved total angular momentum, but the connection between angular momentum and momentum is completely obscured. As we saw above, any local correspondence between these two quantities is doomed unless the canonical energy-momentum tensor is symmetric. Furthermore, local angular momentum densities and currents defined in this way will not be gauge-invariant and therefore cannot have a physical interpretation. Alternatively, if we can somehow define a 'new and improved' symmetric energy-momentum tensor, we can make the connection between momentum and angular momentum explicit by constructing the angular momentum density locally. We will see that this construction also leads naturally to gauge-invariant local densities.

Looking back on our argument that the total momentum is conserved, we see that the only crucial property of $\mathcal{T}$ is that its divergence equals zero. The addition of any divergence-free tensor to $\mathcal{T}$ will yield another tensor from which conserved quantities can be derived. However, this argument does not show that the new quantities are the momenta of the system. The conservation of $\mathcal{P}$ is a *direct* consequence of translation invariance, as we saw above. This more than anything else promotes its identification with the momentum of the system. Alternative tensors may change the momentum density, $\mathcal{T}^{0\mu}$, allowing for a conserved angular momentum density, but must not change the integrated quantity $\mathcal{P}$. Both of these requirements are satisfied by the *Belinfanté* transformation,

$$\mathcal{T}^{\mu\nu} \rightarrow \mathcal{T}^{\mu\nu} + \partial_\alpha \chi^{[\alpha\mu]\nu} \quad . \tag{D.53}$$

The antisymmetry of χ under $\mu\alpha$ exchange guarantees the new tensor to be divergence-free.

[17] This is true for local *internal* symmetries. Gauge fields associated with *external* symmetries may have different transformation properties. For example, the gauge field associated with local Lorentz invariance transforms like a traceless rank-2 symmetric tensor. This causes the associated gauge boson, the graviton, to have spin-2.

Furthermore, the difference between the new conserved quantities and $\mathcal{P}$ is just the surface integral

$$\int d^3x\, \partial_\alpha \chi^{[\alpha 0]\nu} = \int d^3x\, \partial_i \chi^{[i0]\nu} = \oint dS_i\, \chi^{[i0]\nu} \quad , \tag{D.54}$$

which can certainly be chosen to vanish over a large enough region.[18] If there exists a χ such that the remaining tensor is symmetric, this tensor can be used to define angular momentum densities whose integrals represent the conserved total angular momentum of the system. Let us see how the assumption of Lorentz invariance leads to the existence of such a χ.

Under the action of the Lorentz group, a scalar lagrangian transforms only because its argument does :

$$\mathcal{L}(x) \to \mathcal{L}(\Lambda x) = \mathcal{L}(x) + \frac{1}{2}\left(\theta_{\mu\nu} J^{\mu\nu}\right)_{\alpha\beta} x^\beta \partial^\alpha \mathcal{L}(x) + \mathcal{O}(\theta^2) \quad . \tag{D.55}$$

This is a translation of the same form as that considered above. Since we have already taken the translation invariance of the action into account, we will ignore the change in spatial coordinate and focus on the other transformations induced by the Lorentz group.

The rotation generators do not commute with the Pauli-Ljubanskii vector, so they will in general mix different elements of irreducible representations of the Poincaré group; under an infinitesimal global Lorentz transformation, our fields transform as[19]

$$\begin{aligned} \Phi_m &\to \Phi'_m = \Phi_m - \frac{i}{2}\left(\theta_{\mu\nu}\Sigma^{\mu\nu}\right)_{mn} \Phi_n \\ \partial_\alpha \Phi_m &\to \partial_\alpha \Phi_m - \frac{1}{2}\left(\theta_{\mu\nu} J^{\mu\nu}\right)_{\alpha\beta} \partial^\beta \Phi_m - \frac{i}{2}\left(\theta_{\mu\nu}\Sigma^{\mu\nu}\right)_{mn} \partial_\alpha \Phi_n \end{aligned} \tag{D.56}$$

Here, m and n label different fields in our theory. Σ will depend on the representation the fields belong to (or, equivalently, the spin) and J is the generator of coordinate rotations,

$$\left(J^{\mu\nu}\right)_{\alpha\beta} = \left(\delta^\mu_\alpha \delta^\nu_\beta - \delta^\nu_\alpha \delta^\mu_\beta\right) \quad . \tag{D.57}$$

Under this transformation, the lagrangian itself must be invariant since we are ignoring its variation due to the coordinate translation. Hence Lorentz invariance implies

$$0 \quad = \quad -\frac{i}{2}\frac{\delta \mathcal{L}}{\delta \Phi}\Sigma^{\mu\nu}\Phi - \frac{i}{2}\frac{\delta \mathcal{L}}{\delta \partial_\alpha \Phi}\Sigma^{\mu\nu}\partial_\alpha \Phi - \frac{1}{2}\frac{\delta \mathcal{L}}{\delta \partial_\alpha \Phi}\left(J^{\mu\nu}\right)_{\alpha\beta}\partial^\beta \Phi$$

[18] barring topological effects...

[19] Note that the index on the derivative is treated like a vector index here.

$$
\begin{aligned}
&= \frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\partial^\nu\Phi - \frac{\delta\mathcal{L}}{\delta\partial_\nu\Phi}\partial^\mu\Phi \\
&\quad -i\left(\frac{\delta\mathcal{L}}{\delta\Phi}\Sigma^{\mu\nu}\Phi + \frac{\delta\mathcal{L}}{\delta\partial_\alpha\Phi}\Sigma^{\mu\nu}\partial_\alpha\Phi\right) \\
&= \mathcal{T}^{\mu\nu} - \mathcal{T}^{\nu\mu} - i\partial_\alpha\left(\frac{\delta\mathcal{L}}{\delta\partial_\alpha\Phi}\Sigma^{\mu\nu}\Phi\right) \quad ,
\end{aligned}
\tag{D.58}
$$

where in the last line I have used the equation of motion, (D.42), and the fact that $\Sigma^{\mu\nu}$ does not depend on x^μ.

This equation is rather interesting. Rather than giving us a divergence-free current from which we can construct a conserved angular momentum density, Lorentz invariance relates the antisymmetric part of $\mathcal{T}$ to a total divergence. We can use this to immediately construct the canonical angular momentum current density

$$
\tilde{\mathcal{M}}^{\alpha\mu\nu} \equiv x^\mu\mathcal{T}^{\alpha\nu} - x^\nu\mathcal{T}^{\alpha\mu} - i\frac{\delta\mathcal{L}}{\delta\partial_\alpha\Phi}\Sigma^{\mu\nu}\Phi \tag{D.59}
$$

which follows directly from Lorentz invariance. As is usually the case with canonical quantities, this object is easy to interpret : the first two terms are due to orbital angular momentum while the last is due entirely to intrinsic spin angular momentum. Unfortunately, $\tilde{\mathcal{M}}$ also displays one of the less favorable attributes of canonical quantities - it is gauge-dependent.

Alternatively, notice that the symmetric part of $\mathcal{T}$ can be written as

$$
\mathcal{T}^{(\mu\nu)} = \mathcal{T}^{\mu\nu} - \frac{i}{2}\partial_\alpha\left(\frac{\delta\mathcal{L}}{\delta\partial_\alpha\Phi}\Sigma^{\mu\nu}\Phi\right) \quad . \tag{D.60}
$$

This is reminiscent of a Belinfanté transformation, (D.53), but our version of χ is not antisymmetric in $\alpha\mu$. We can fix this by subtracting a term in which $\alpha \leftrightarrow \mu$, but the tensor we are left with is no longer symmetric. Fortunately, the term we need to add to restore the symmetry is automatically antisymmetric in $\mu\alpha$ due to the antisymmetry of Σ; the tensor

$$
\begin{aligned}
\Theta^{\mu\nu} \quad &\equiv \quad \mathcal{T}^{\mu\nu} - \frac{i}{2}\partial_\alpha\left[\frac{\delta\mathcal{L}}{\delta\partial_\alpha\Phi}\Sigma^{\mu\nu}\Phi\right. \\
&\qquad\qquad \left. - \frac{\delta\mathcal{L}}{\delta\partial_\mu\Phi}\Sigma^{\alpha\nu}\Phi - \frac{\delta\mathcal{L}}{\delta\partial_\nu\Phi}\Sigma^{\alpha\mu}\Phi\right]
\end{aligned}
\tag{D.61}
$$

is both symmetric and physically indistinguishable from $\mathcal{T}$. In gauge theories, we will also find it to be gauge-invariant. This is the tensor which can truly be interpreted as the momentum current density of our system. According to the above discussion, we can use Θ to form the angular momentum current density

$$
\mathcal{M}^{\alpha\mu\nu} = x^\mu\Theta^{\alpha\nu} - x^\nu\Theta^{\alpha\mu} \quad . \tag{D.62}
$$

This improved Belinfanté form differs from $\tilde{\mathcal{M}}$ locally, but one can show by direct substitution that the conserved total angular momentum

$$\mathcal{J}^{\mu\nu} = \int d^3x\, \mathcal{M}^{0\mu\nu}(x) \tag{D.63}$$

is unaffected.[20]

While the above discussion is perfectly fine in classical theories, some modification is clearly needed before these results can be applied to quantized fields. In the path-integral approach, the transition is quite clean. We begin with the vacuum matrix element

$$\left\langle \mathrm{T}\hat{\mathcal{O}}(x_1, \ldots, x_n) \right\rangle = \frac{1}{\mathcal{Z}[0]} \int \mathcal{D}[\Phi] \mathcal{O}(x_1, \ldots, x_n) e^{i\mathcal{S}[\Phi]} \ , \tag{D.64}$$

where $\hat{\mathcal{O}}$ is an arbitrary[21] collection of field operators depending on the spacetime coordinates $x_1, \ldots x_n$. Let us ignore the normalization factor $\mathcal{Z}[0]$ for the moment and focus on the path integral. Since Φ is nothing but a dummy variable of integration, it is clear that

$$\int \mathcal{D}[\Phi] \mathcal{O}(x_1, \ldots, x_n) e^{i\mathcal{S}[\Phi]} = \int \mathcal{D}[\Phi'] \mathcal{O}'(x_1, \ldots, x_n) e^{i\mathcal{S}[\Phi']} \ , \tag{D.65}$$

where $\mathcal{O}'$ is the functional $\mathcal{O}$ evaluated at the field configuration Φ'. Taking $\Phi'(x) = \Phi(x) + \delta\Phi(x)$, with $\delta\Phi$ infinitesimal but otherwise unconstrained, we rewrite (D.65) as[22]

$$\begin{aligned} \int \mathcal{D}[\Phi] \mathcal{O}(x_1, \ldots, x_n) e^{i\mathcal{S}[\Phi]} = \int \mathcal{D}[\Phi] \left| \det \left(\frac{\delta\Phi'(x)}{\delta\Phi(y)} \right) \right| \\ \times \left[\mathcal{O}(x_1, \ldots, x_n) + \frac{\delta\mathcal{O}}{\delta\Phi(x)} \delta\Phi(x) + i \frac{\delta\mathcal{S}[\Phi]}{\delta\Phi(x)} \delta\Phi(x) \mathcal{O}(x_1, \ldots, x_n) \right] e^{i\mathcal{S}[\Phi]} \end{aligned} \tag{D.66}$$

Let us consider the arbitrary variation

$$\delta\Phi(x) = \epsilon(x) \ . \tag{D.67}$$

In this case, our variation is merely a field-independent shift of the integration variable. Hence the Jacobian is trivial. This immediately leads to the relation

$$\int \mathcal{D}[\Phi] \left[\frac{\delta\mathcal{O}}{\delta\Phi(x)} \delta\Phi(x) + i \frac{\delta\mathcal{S}[\Phi]}{\delta\Phi(x)} \delta\Phi(x) \mathcal{O}(x_1, \ldots, x_n) \right] e^{i\mathcal{S}[\Phi]} = 0 \ . \tag{D.68}$$

[20] once again, ignoring surface terms.

[21] In a gauge theory, we will require it to be gauge invariant.

[22] Recall that repeated indices imply summation. Here, the repeated 'index' is the spacetime variable x. Also note that the determinant is inverted for Grassmann fields.

Writing

$$\frac{\delta \mathcal{S}[\Phi]}{\delta \Phi(x)} \delta\Phi(x) = \left[\frac{\delta \mathcal{L}(x)}{\delta \Phi(x)} - \partial_\mu \frac{\delta \mathcal{L}(x)}{\delta \partial_\mu \Phi(x)} \right] \epsilon(x) \quad , \tag{D.69}$$

and asserting the arbitrariness of $\epsilon(x)$, we see that

$$\left\langle \mathrm{T} \left\{ \left[\frac{\delta \mathcal{L}(x)}{\delta \Phi(x)} - \partial_\mu \frac{\delta \mathcal{L}(x)}{\delta \partial_\mu \Phi(x)} \right] \mathcal{O}(x_1, \ldots, x_n) \right\} \right\rangle = i \left\langle \mathrm{T} \frac{\delta \mathcal{O}(x_1, \ldots, x_n)}{\delta \Phi(x)} \right\rangle \quad . \tag{D.70}$$

The right-hand-side of this equation is a series of *contact terms* which involve $\delta(x - x_i)$ for some $i \in \{1, n\}$, while the left-hand-side is a Green's function involving the equation of motion operator

$$\frac{\delta \mathcal{L}}{\delta \Phi} - \partial_\mu \frac{\delta \mathcal{L}}{\delta \partial_\mu \Phi} \quad . \tag{D.71}$$

Equation (D.70) is an extremely useful relation. In particular, it tells us that all physical matrix elements of (D.71) vanish since in that case the operator $\mathcal{O}$ represents asymptotic fields and will have arguments only at infinity.

At this point, it is obvious how to proceed with other variations of the fields Φ. All we need is a transformation which leaves the measure of the path integral invariant[23] to arrive at an identity of the form

$$\langle \mathrm{T} \{ \mathcal{C}(x) \mathcal{O}(x_1, \ldots, x_n) \} \rangle = i \left\langle \mathrm{T} \frac{\delta \mathcal{O}(x_1, \ldots, x_n)}{\delta \Phi(x)} \right\rangle \quad , \tag{D.72}$$

where $\mathcal{C}$ is a collection of fields which is classically zero. In particular, the classical relations involving conservation of energy-momentum and angular momentum can easily be cast into this form since the transformations of translation and Lorentz rotation leave the measure invariant.[24]

The lagrangian density of massless QCD is

$$\mathcal{L} = \overline{\psi}\, i \not{\mathcal{D}}\, \psi - \frac{1}{4} \mathcal{F}^2 \quad , \tag{D.73}$$

from which we derive[25]

$$\frac{\delta \mathcal{L}}{\delta \psi} \quad = \quad -g \overline{\psi} \not{\mathcal{A}} \tag{D.75}$$

[23] Actually, useful identities can be derived even when the measure is not left invariant. One extremely well-known example is the derivation of the Adler-Bell-Jackiw anomaly [98].

[24] For fermions, we can see this as a consequence of the invariance of the product $\overline{\psi}\psi$, while for vector bosons it is due to the fact that $\det \Lambda = \pm 1$ for the Lorentz transformations introduced in Section B.1.

[25] These relations are *not* unique. In particular, the replacement $i\overrightarrow{\mathcal{D}} \rightarrow i\overleftrightarrow{\mathcal{D}}$ will affect the first four. However,

$$\frac{\delta\mathcal{L}}{\delta\partial_\mu\psi} = \overline{\psi}\, i\gamma^\mu \tag{D.76}$$

$$\frac{\delta\mathcal{L}}{\delta\overline{\overline{\psi}}} = i\not{\mathcal{D}}\,\psi \tag{D.77}$$

$$\frac{\delta\mathcal{L}}{\delta\partial_\mu\overline{\overline{\psi}}} = 0 \tag{D.78}$$

$$\frac{\delta\mathcal{L}}{\delta\mathcal{A}^a_\mu} = -g\overline{\psi}\gamma^\mu t^a\psi - gf^{abc}\mathcal{F}^{\mu\alpha}_b\mathcal{A}^c_\alpha \tag{D.79}$$

$$\frac{\delta\mathcal{L}}{\delta\partial_\mu\mathcal{A}^a_\nu} = -\mathcal{F}^{\mu\nu}_a \quad . \tag{D.80}$$

These relations, along with the Lorentz transformation properties

$$\Sigma^{\mu\nu}_q = \frac{1}{2}\sigma^{\mu\nu} \tag{D.81}$$

$$\Sigma^{\mu\nu}_g = -iJ^{\mu\nu} \quad , \tag{D.82}$$

contain the QCD equations of motion, energy-momentum tensor, angular momentum current density, and any other symmetry currents of[26] QCD. Some useful relations are summarized in Section A.4.

One very useful identity that can easily be proved at this point is the fundamental *Ward identity* of QCD. Taking

$$\delta\mathcal{A}^\mu_a(x) = \partial^\mu\epsilon_a(x) \quad , \tag{D.83}$$

---

the equations of motion and physical tensor densities are not affected by this ambiguity. We also note that the above derivation assumes commuting, rather than Grassmann, fields. To generalize our expressions, it is sufficient to consider Grassmann derivatives as coming *from the direction of the variation*. For example,

$$\begin{aligned}\frac{\delta\mathcal{L}}{\delta\Phi}\,\delta\Phi \quad &\rightarrow \quad \delta\overline{\psi}\,\frac{\overrightarrow{\delta}}{\delta\overline{\overline{\psi}}}\,\mathcal{L} \\ &\rightarrow \quad \mathcal{L}\,\frac{\overleftarrow{\delta}}{\delta\psi}\,\delta\psi \quad .\end{aligned} \tag{D.74}$$

[26] unrenormalized and un-fixed

(D.72) becomes[27]

$$\partial_\mu \left\langle \mathrm{T}\left\{ \left[g\overline{\psi}\gamma^\mu t^a \psi + g f^{abc} \mathcal{F}_b^{\mu\alpha} \mathcal{A}_\alpha^c \right] \mathcal{O}(x_1, \ldots, x_n) \right\} \right\rangle = -i\partial_\mu \left\langle \mathrm{T} \frac{\delta \mathcal{O}(x_1, \ldots, x_n)}{\delta \mathcal{A}_\mu^a(x)} \right\rangle \quad . \tag{D.84}$$

The expression in brackets is constructed to be the current the gluon field couples to, so this equation tells us that Green's functions involving a gluon current contracted with its momentum are reduced to contact terms. In particular, it implies transversality of the gluon propagator and conservation of the color charge. In this sense, the terms on the right-hand-side act as *sources* for the color current. Nonzero contact terms imply a color source, which will affect the total charge. Note that this identity is also obvious from the classical equation of motion of the gluon field,

$$\mathcal{D}_\mu^{ab} \mathcal{F}_b^{\mu\alpha} = g\overline{\psi}\gamma^\alpha t^a \psi \quad , \tag{D.85}$$

and the antisymmetry of the field strength.

An analogous identity implies the conservation of quark flavor (or, more generally, baryon and lepton number). Taking

$$\delta\psi(x) = i\alpha(x)\psi(x) \quad , \quad \delta\overline{\psi}(x) = -i\alpha(x)\overline{\psi}(x) \tag{D.86}$$

gives[28]

$$\partial_\mu \left\langle \mathrm{T}\left\{ \overline{\psi}(x)\gamma^\mu \psi(x) \mathcal{O}(x_1, \ldots, x_n) \right\} \right\rangle = \left\langle \mathrm{T}\left\{ \overline{\psi}(x) \frac{\delta \mathcal{O}(x_1, \ldots, x_n)}{\delta \overline{\psi}(x)} - \frac{\delta \mathcal{O}(x_1, \ldots, x_n)}{\delta \psi(x)} \psi(x) \right\} \right\rangle \quad . \tag{D.87}$$

This relation also finds use as the fundamental Ward identity of QED.

These identities have all been derived ignoring the necessary steps of renormalization and gauge-fixing. The fact that these unrenormalized and un-fixed Ward identities imply similar relations among the fully renormalized gauge-fixed Green's functions is nontrivial to show. Nonetheless, it has been done in the covariant gauges [41] and the light-cone gauge [44].

[27] Derivatives *always* belong *outside* the matrix element. The easiest way to see this involves the sources in the path integral approach. Since our sources bring down powers of the fields rather than their derivatives, we must differentiate *externally*. Among other things, this implies that derivatives will always generate contributions from the time-ordered product. These form the contact terms on the right-hand-side of our matrix element identities. As mentioned in Section 2.5, these contact terms can systematically be removed by considering the modified T* product. See [63].

[28] Once again, we must take care with these anticommuting derivatives.

Once these renormalized and gauge-fixed Ward identities are proven, the renormalizability of the whole theory is just a step away. Thanks to these identities, all of the ultraviolet divergences present in the theory are constrained to take forms which respect our symmetries. Since we have chosen the most general lagrangian consistent with the symmetry, this implies that all ultraviolet divergences can be absorbed into counterterms, which in turn implies renormalizability.

Appendix E

Dimensional Regularization

This Appendix is meant to familiarize the reader with the standard regularization procedure used in quantum field theory. Introduced by G. 't Hooft and M. Veltman in 1972 [10], this procedure exploits the analytic behavior of integrals as functions of their dimensionality. Since the ultraviolet (and infrared) divergences we meet in the text are in some sense accidents of the fact that we live in four dimensions, these integrals prove entirely convergent when evaluated in less (or more) spacetime dimensions. The subtleties associated with the actual meaning of a 3.5-dimensional space are completely avoided here; some discussion can be found in [40]. We will view dimensional regularization exclusively as a consistent *regularization procedure* for our amplitudes.

It can be shown quite generally that the coordinates

$$\begin{aligned}
x_1 &= \rho \cos\theta_1 \\
x_2 &= \rho \sin\theta_1 \cos\theta_2 \\
x_3 &= \rho \sin\theta_1 \sin\theta_2 \cos\theta_3 \\
\vdots &= \vdots \\
x_i &= \rho \prod_{j=1}^{i-1} \sin\theta_j \cos\theta_i \qquad \text{(E.1)} \\
\vdots &= \vdots \\
x_{(n-1)} &= \rho \prod_{j=1}^{n-2} \sin\theta_j \cos\theta_{(n-1)} \\
x_n &= \rho \prod_{j=1}^{n-1} \sin\theta_j \qquad \text{(E.2)}
\end{aligned}$$

span a Euclidean n-dimensional space for integer $n > 1$. The angles θ_i all take values from 0 to π except $\theta_{(n-1)}$, which takes values on $[0, 2\pi)$. The 2-norm of the vector $\mathbf{x}$ is given by ρ and the $(n-1)$ angles specify a location on the $(n-1)$-dimensional sphere defined by ρ. The line element in these coordinates

$$ds^2 = d\rho^2 + \rho^2\, d\theta_1^2 + \rho^2 \sum_{i=2}^{n-1} \left(\prod_{j=1}^{i-1} \sin^2 \theta_j \right) d\theta_i^2 \tag{E.3}$$

defines the volume element

$$d\,\mathbf{x} = \sqrt{\det\, g_{ij}}\, d\rho \prod_{i=1}^{n-1} d\theta_i = \rho^{n-1} \left(\prod_{i=1}^{n-1} \sin^{n-i-1} \theta_i \right) d\rho \prod_{j=1}^{n-1} d\theta_j \ . \tag{E.4}$$

The hyper-area of our $(n-1)$-dimensional sphere then becomes

$$\Omega_{(n-1)} = 2\pi \prod_{i=1}^{n-2} \int_0^\pi \sin^{n-i-1} \theta_i\, d\theta_i \ , \tag{E.5}$$

which upon using

$$\int_0^\pi \sin^n \theta d\theta = 2^n \frac{\Gamma^2\left(\frac{n+1}{2}\right)}{\Gamma(n+1)} = \sqrt{\pi}\, \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2}+1\right)} \tag{E.6}$$

reduces to

$$\Omega_{(n-1)} = \frac{2\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)} \ . \tag{E.7}$$

For any function $f(\mathbf{x})$ that depends only on the magnitude ρ of $\mathbf{x}$ we have

$$\int d\,\mathbf{x}\, f(\mathbf{x}) = \Omega_{(n-1)} \int d\rho\, \rho^{n-1} f(\rho) \ . \tag{E.8}$$

This expression is correct for any positive integer dimension n. Furthermore, these integrals respect shift invariance

$$\int d\,\mathbf{x}\, f(\mathbf{x}+\mathbf{y}) = \int d\,\mathbf{x}\, f(\mathbf{x}) \ , \tag{E.9}$$

for any constant vector $\mathbf{y}$, and the dilation identity

$$\int d\,\mathbf{x}\, f(a\,\mathbf{x}) = \frac{1}{|a|^n} \int d\,\mathbf{x}\, f(\mathbf{x}) \ , \tag{E.10}$$

where a is an arbitrary real constant.

We can *define* the functional

$$\mathcal{I}_d[f](\mathbf{y}_1, \ldots, \mathbf{y}_N) \tag{E.11}$$

as a *meromorphic* function of d that satisfies

$$\mathcal{I}_d[f](\mathbf{y}_1+\mathbf{y},\ldots,\mathbf{y}_N+\mathbf{y}) = \mathcal{I}_d[f](\mathbf{y}_1,\ldots,\mathbf{y}_N) \tag{E.12}$$

$$\mathcal{I}_d[f](a\mathbf{y}_1,\ldots,a\mathbf{y}_N) = |a|^d a^{\phi_f}\mathcal{I}_d[f](\mathbf{y}_1,\ldots,\mathbf{y}_N) \tag{E.13}$$

and takes the value

$$\mathcal{I}_n[f](\mathbf{y}_1,\ldots,\mathbf{y}_N) = \int d\mathbf{x} f(\mathbf{x},\mathbf{y}_1,\ldots,\mathbf{y}_N) \tag{E.14}$$

for $n \in \mathcal{Z}_+$.[1] The constant ϕ_f is a scale factor which depends only on f. This does *not* fix our functional uniquely. One can add *any* analytic function of d which vanishes at all positive integers to $\mathcal{I}_d$ to obtain an equally acceptable functional. This ambiguity will be resolved shortly.

The vectors $\{\mathbf{y}_i\}$ define at most N directions, so in $M > N$ integer dimensions $\mathcal{I}_M$ contains at most N directionally-dependent integrals. Doing these integrals explicitly gives

$$\mathcal{I}_M[f](\mathbf{y}_1,\ldots,\mathbf{y}_N) = \mathcal{I}_{M-N}\left[\mathcal{I}_N[f]\right](\mathbf{y}_1,\ldots,\mathbf{y}_N) \ , \tag{E.15}$$

where the remaining $(M-N)$ integrals are direction *independent*, i.e. of the form of (E.8). If $\mathcal{I}_d$ is to be a meromorphic function of d constrained by (E.14), it must also satisfy this reduction identity :

$$\mathcal{I}_d[f](\mathbf{y}_1,\ldots,\mathbf{y}_N) = \mathcal{I}_{d-N}\left[\mathcal{I}_N[f]\right](\mathbf{y}_1,\ldots,\mathbf{y}_N) \ . \tag{E.16}$$

This immediately implies that we need not consider directionally-dependent functions f. These can always be reduced to directionally-independent functions by doing an integer number of integrations.[2]

With this in mind, we can use (E.8) as a guideline to define a *unique* functional. For any given directionally-independent function $f(\rho)$, we define

$$\mathcal{I}_d[f] \equiv \Omega_{(d-1)} \int d\rho\, \rho^{d-1} f(\rho) \ , \tag{E.17}$$

where the integral on the right is to be evaluated in a suitable region[3] of the complex plane of d and analytically continued elsewhere. Since it has been constructed to look like an integral,

[1] unless this integral is not well-defined. We will see that these cases are handled quite nicely by the other properties of $\mathcal{I}_d$.

[2] although this is not always the most practical way to extract the value.

[3] This is taken to mean anywhere the integral is well-defined.

one often uses the notation

$$\mathcal{I}_d[f](\mathbf{y}_1,\cdots,\mathbf{y}_N) = \int d\mathbf{x}\, f(\mathbf{x},\mathbf{y}_1,\cdots,\mathbf{y}_N) \ , \tag{E.18}$$

but it must always be understood that this is simply a convenient way to write things. In this form, the shift and dilation invariance imply that one can manipulate the variable of integration as in a normal integral. However, nonlinear changes of variables require care; the interested reader is referred to [40].

Now that we know what a 'd-dimensional integral' means, let's compute some. First and foremost, *any* functional $\mathcal{I}_d[f]$ whose arguments $\{\mathbf{y}_i\}$ are all zero is identically zero.[4] This follows from the dilation property,

$$\mathcal{I}_d[f](\mathbf{0},\ldots,\mathbf{0}) = |a|^d a^{\phi_f}\mathcal{I}_d[f](\mathbf{0},\ldots,\mathbf{0}) \ , \tag{E.19}$$

and the analytic structure of $\mathcal{I}_d$ as a function of d. This result is somewhat striking since it implies that the integrals

$$\int d\mathbf{x}\,(\rho^2)^n \tag{E.20}$$

all take the value zero. However, as a consequence of (E.13), it is unalterable.[5] Using (E.19) in conjunction with (E.12), one can also easily show that

$$\mathcal{I}_d[f](\mathbf{y},\ldots,\mathbf{y}) = 0 \ . \tag{E.21}$$

Apparently, $\mathcal{I}_d$ must have *two* vectors on which to depend if it is to be nonzero.[6]

Throughout the above, I have ignored the dependence of $\mathcal{I}_d$ on scalar quantities. This dependence must, of course, be taken into account if one is to have a general functional. In particular, the direction-independent integrals we have asserted to be the only necessary consideration depend *exclusively* on scalar parameters. Since scalar quantities are not affected by shifts in the integration variable, there is no analogue of (E.12) for these quantities. However, these parameters *are* affected by dilation :

$$\mathcal{I}_d[f](a\lambda_1,\ldots,a\lambda_N) = |a|^d a^{\phi_f}\mathcal{I}_d[f](\lambda_1,\ldots,\lambda_N) \ . \tag{E.22}$$

[4] We need all scalar arguments to be zero as well; see below.

[5] Integrals in positive integer dimensions do not satisfy this equation. As a consequence, they are ill-defined.

[6] Note that these vectors can be taken as $\mathbf{y}$ and $\mathbf{0}$ if one wishes...

This relation again implies that parameterless integrals are necessarily zero. However, the lack of shift invariance means that we only need one parameter to obtain a finite result. In quantum field theory, we will find that the only truly d-dimensional integral one needs to compute is

$$\int \frac{d^{\,d}k}{(2\pi)^d} \frac{\left(k^2\right)^\alpha}{\left(k^2+M^2\right)^\beta} = \frac{1}{(4\pi)^{d/2}} \frac{2}{\Gamma\left(\frac{d}{2}\right)} \int_0^\infty dk \, \frac{\left(k^2\right)^{\alpha+(d-1)/2}}{\left(k^2+M^2\right)^\beta} \quad . \tag{E.23}$$

For sufficiently small odd d and integer α, β, one can extend the integration region to negative infinity and take the integral by contour. If we are very strong, we can show that this gives[7]

$$\frac{1}{(4\pi)^{d/2}} \left(M^2\right)^{d/2+\alpha-\beta} \, \frac{\Gamma(d/2+\alpha)\Gamma(\beta-\alpha-d/2)}{\Gamma(d/2)\Gamma(\beta)} \quad . \tag{E.24}$$

Analytically continuing this result to the full complex plane of α, β, and d gives the required general integral.

Dimensional regularization is a procedure in which one systematically replaces all four-dimensional loop-momentum integrals by d-dimensional ones. In this way, the ultraviolet divergences present in our theory are represented by poles at $d = 4$. As mentioned above this process is viewed exclusively as a way to exhibit ultraviolet (and infrared) divergences explicitly as well-defined quantities. From this perspective, one would expect no changes in the theory to be necessary. Unfortunately, small things have a way of infiltrating every aspect of a quantum field theory. Interpreting a loop-momentum integral as d-dimensional implies in turn that the loop-momenta are themselves d-dimensional, which implies that the metric used to form invariant products between them is d-dimensional. This fact changes the whole structure of the spacetime symmetry group of our theory.

Looking back at Section B.1, we see that things are not so bad. Almost all of the constructs of that section can be accommodated in d dimensions.[8] We have to write

$$g_{\mu\nu} g^{\mu\nu} = d \tag{E.25}$$

rather than four and we find ourselves with $d(d-1)/2$ rotation generators rather than six. We can easily keep track of such factors. The only change in the above formalism is that f must now be allowed to depend on d, which we would've had to consider at higher orders

[7] A much more elegant derivation is given in [40].

[8] To minimize the change in structure to our group, we add only spacelike coordinates.

anyway.[9] These modifications will affect physical amplitudes when multiplied by poles at $d = 4$. Renormalized quantities are guaranteed to be finite, but the divergences are certainly not guaranteed to leave no trace.

The intimate relation between $SO(3,1)$ and $SL(2,C)$ requires us to continue our Dirac space into d dimensions as well. The smallest dimensionality in which we can find d γ-matrices which satisfy the Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} \tag{E.27}$$

is $2^{d/2} \times 2^{d/2}$, so we should modify the normalizations

$$\gamma^\mu \gamma_\mu = d \tag{E.28}$$

$$\text{Tr } \mathbf{1} = 2^{d/2} \ . \tag{E.29}$$

Since the trace appears in all amplitudes, it constitutes an overall factor. As such, it can be *chosen* not to leave a residue by including its effects with the ultraviolet divergence (along with the γ_E and $\log 4\pi$ in the $\overline{\text{MS}}$ scheme, for example). For this reason, people usually just take $\text{Tr } \mathbf{1} = 4$.

In addition to the above changes, we must somehow make a connection between the d-dimensional *Euclidean* space our integrals are defined in and the d-dimensional *Minkowskii* space in which our theory is defined. The simple solution to this is to rotate our energy integration axis 90° in its complex plane. This has the effect

$$k^0 \to i k_E^0 \tag{E.30}$$

$$k^2 \to -k_E^2 \ . \tag{E.31}$$

Since we have constructed our propagators to have a well-defined pole structure, this rotation can be justified for our Feynman integrals by Cauchy's theorem.[10] All we must do is keep

[9] This does cause a slight problem with our derivation of (E.19) in the case $\phi = -d$. The result of this is that the integral

$$\int \frac{d^d k}{(k^2)^{d/2}} \tag{E.26}$$

is not uniquely defined. We cannot take its integer-dimension value since this is also not well-defined, yet we cannot prove that it is zero using the scaling argument. Even (E.24) is poorly defined in this case. However, since it is not determined, we are free to *define* it as zero.

[10] We can perform this shift provided we don't cross any poles in the process. One can check explicitly that

track of the sign and the factor of i. At the end of the calculation, we simply rotate back to obtain amplitudes with physical momenta.

Unfortunately, all of the sectors of our theory are not so readily expressed in d dimensions. One can trace all of the difficulties to the Levi-Cevita tensor, $\epsilon^{\mu\nu\alpha\beta}$. Looking back at its definition (B.19), we see that this object appears because of the relation between the determinant of Λ and its components. A similar object can be defined in n dimensions as a totally antisymmetric n-index tensor. Its invariance is based on its equal treatment of *all* of the dimensions of our space. We have no analogue of this in d dimensions. Mathematicians tell us that if we *really* want to take a non-integer-dimensional space seriously, its vectors will have an infinite number of components,[11] an infinite number of which vanish as d approaches a positive integer. Since we do not intend to introduce tensors with an infinite number of indices into our calculations (and have no idea how we would if we did intend!), we cannot determine a definition for ϵ in our dimensionally regularized theory. As before, this means that we are free to define one however we wish, *provided that it is consistent.*[12] Our studies of renormalization in Section 1.4 imply that we can choose whatever regularization scheme we want, as long as we stick with our scheme once it is chosen. The most obvious choice is simply to keep the four-dimensional definition. This carries the price of breaking the full rotational symmetry of the d-dimensional theory, but it is difficult to imagine a definition that will not.

When products of two ϵ tensors appear, we can pretend not to break the symmetry by eliminating the ϵ's via

$$\epsilon^{\mu\nu\alpha\beta}\epsilon_{\lambda\rho\sigma\tau} = -\delta^\mu_\lambda\delta^\nu_\rho\delta^\alpha_\sigma\delta^\beta_\tau \quad + \quad \text{permutations} \ . \tag{E.32}$$

Once we have everything in terms of δ's, (E.25) makes us feel like we are truly working in d dimensions. However, the reduction identity (E.32) is derived in four dimensions by working out the implications of the statement, "If the product of two epsilons is not zero, then each index on one must have a twin on the other.". This statement is obviously not generalizable to d dimensions. On the other hand, one can use (E.32) to define an ϵ scheme. This option

this leads to no problems here, unless one quantizes the theory in axial gauge with a pathological prescription for the spurious singularities. See Section 1.6.

[11] some discussion can be found in [40].

[12] It also should not do irreparable damage to Lorentz invariance.

is described extensively in [99]. Yet another scheme which considers ϵ a totally antisymmetric tensor with four indices in the full d-dimensional space has been brought forth in connection with a non-associative Dirac-Clifford algebra [100].

In the fermion sector, the Levi-Cevita symbol's ambassador, γ_5, poses similar difficulties. Because of the role γ_5 plays in fermion helicity projection operators and the prominence of fermions in real world applications, different choices for γ_5's extension have been extensively studied in the literature. The schemes most often seen are Bardeen's anticommuting γ_5 [101] and 't Hooft and Veltman's γ_5 prescription [10]. The first of these defines γ_5 through the relation

$$\{\gamma^\mu, \gamma_5\} = 0 \ , \tag{E.33}$$

which is taken to be valid for *all* μ. Although this is a natural extension, it is algebraically inconsistent with a nonzero trace. Beginning with

$$\begin{aligned} d \operatorname{Tr} \gamma_5 &= \operatorname{Tr} \gamma^\lambda \gamma_\lambda \gamma_5 \\ &= -\operatorname{Tr} \gamma^\lambda \gamma_5 \gamma_\lambda \qquad \text{(E.34)} \\ &= -d \operatorname{Tr} \gamma_5 \ , \qquad \text{(E.35)} \end{aligned}$$

one can show that

$$d(d-2)(d-4)\cdots(d-2n) \operatorname{Tr} \gamma_5 \gamma^{\mu_1} \gamma^{\mu_2} \cdots \gamma^{\mu_{2n}} = 0 \tag{E.36}$$

for any integer n. This is expected from the relation γ_5 has with ϵ; the purpose of γ_5 is to force the presence of *all* directions. However, a γ_5 which does not give finite traces is useless to us. It has been suggested that dropping some other property of the spinor algebra may lead to a consistent scheme. Examples are the algebra's associativity [100] and the trace's cyclicity [102].

The 't Hooft and Veltman scheme is analogous to the scheme introduced above for the Levi-Cevita symbol. One simply clings to the four-dimensional definition of γ_5,

$$\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 \ , \tag{E.37}$$

in d dimensions. This scheme has the advantage of being fully consistent. However, it explicitly breaks the full symmetry of our d-dimensional spacetime. In particular, its commutation

relations with the γ^μ read

$$\{\gamma^\mu, \gamma_5\} = 0 \ ; \quad \mu \in \{0, 3\} \tag{E.38}$$

$$[\gamma^\mu, \gamma_5] = 0 \ ; \quad \mu \notin \{0, 3\} \ . \tag{E.39}$$

This reckless destruction is not without its price. In order to ensure that the nonsinglet axial current is conserved, one must perform an extra finite renormalization. However, once this is done, we are guaranteed a consistent result.

In some sense, all of this discussion of γ_5 is academic for application to QCD. Since γ_5 does not appear in the QCD lagrangian, there is no real reason to introduce it. In a vector theory, γ_5 functions mainly as a projector to help simplify calculations. On the other hand, a theory like the Standard Model whose gauge bosons couple to a partially axial current requires a truly consistent way to separate the positive- and negative helicity components of the Dirac field. Here, a scheme like that of 't Hooft and Veltman's may do irreparable damage to the gauge theory and lead to nonrenormalizability. All of the standard proofs that the Standard Model is renormalizable assume an ultraviolet regulator that does not break the gauge symmetry. However, here it is certainly not clear that dimensional regularization offers such a regulator. Studies of the Ward identities in such theories can be found in [103].

Aside from the difficulties associated with parity-odd structures, dimensional regularization has proven itself an extremely useful tool. The simplicity of its integrals coupled with its respect for symmetry allow one to trust the correctness of one's calculation. One other major advantage of this prescription is the fact that it can be used to regulate infrared as well as ultraviolet divergences. Historically, one has had to keep infrared divergences at bay by introducing offshellness or gauge boson masses. Since the former does not allow the calculation of onshell physical matrix elements while the latter does violence to non-abelian gauge symmetries, these options are clearly not acceptable. Dimensional regularization allows one to calculate onshell matrix elements without breaking the gauge symmetry through gauge masses.

However, one cannot simply regulate *both* infrared *and* ultraviolet divergences simultaneously. This is obvious in a massless theory, where all corrections would give exactly zero due to lack of scale. The correct procedure begins with a calculation of offshell Green's functions. These will have ultraviolet divergences which are to be regulated dimensionally by taking $d < 4$. Through renormalization, we can re-express our amplitude in terms of renormalized

quantities which are ultraviolet finite. At this point, dimensional regularization is no longer needed to regularize *ultraviolet* physics, so we are free to analytically continue our amplitude to $d > 4$ where it is well-suited to handle *infrared* physics. It is now safe to take the onshell limit without worrying about unregularized divergences.

A list of useful formulae for d-dimensional integration, along with some helpful properties of the Γ-function, is given in Section A.3.

Bibliography

[1] The original references are

A. Einstein and M. Grossman, Z. Phys. **62**(1913)225

and

A. Einstein, Ann. Phys., 4, **49**(1916)769.

For a standard modern treatment, see for example

C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, Freeman and Co., New York, 1973.

[2] S. L. Glashow, Nucl. Phys. **22**(1961)579.

[3] S. Weinberg, Phys. Rev. Lett. **19**(1967)1264.

[4] A. Salam in *Elementary Particle Theory*, ed. N. Swartholm, Almquist and Wissell, Stockholm, 1968.

[5] For a review of recent precision QED experiments and calculations, see

T. Kinoshita, ed., *Quantum Electrodynamics*, World Scientific, Singapore, 1990

and

T. Kinoshita in *History of Original Ideas and Basic Discoveries in Particle Physics*, H. Newman and T. Ypsilantis, eds., Plenum Press, New York, 1995.

[6] H. Dehmelt, P. Schwinberg, and R. VanDyck, Jr., Phys. Rev. Lett. **59**(1987)26.

[7] H. Becquerel, Comptes Rendus **122**(1896)420; **122**(1896)501.

[8] M. S. Curie, Comtes Rendus **126**(1898)1101;

P. and S. Curie and G. Bémont, Comptes Rendus **127**(1898)1215.

[9] A. Pais, *Inward Bound*, Clarendon Press, Oxford, 1986.

[10] G. 't Hooft, Nucl. Phys. **B35**(1971)167;
G. 't Hooft and M. Veltman, Nucl. Phys. **B44**(1972)189.

[11] These bosons were discovered in the UA1 and UA2 experiments at the CERN $p\overline{p}$ collider in 1982-3;
UA1 Collaboration, G. Arnison et al., Phys. Lett. **B122**(1983)103 and Phys. Lett. **B126**(1983)476;
UA2 Collaboration, M. Banner et al., Phys. Lett. **B122**(1983)476 and P. Bagnaia et al., Phys. Lett. **B129**(1983)130.

[12] H. Geiger and E. Marsden, Proc. Roy. Soc. **A82**(1909)495.

[13] J. J. Thompson, Phil. Mag. **48**(1899)547;
Kelvin, Phil. Mag. **6**(1902)257;
J. J. Thompson, Phil. Mag. **6**(1903)673;
J. J. Thompson, Proc. Cambr. Phil. Soc. **13**(1904)49.

[14] E. Rutherford, Phil. Mag. **21**(1911)669.

[15] E. Rutherford, Phil. Mag. **37**(1919)581.

[16] J. Chadwick, Nature **129**(1932)312.

[17] N. Bohr, Dansk. Vid. Selsk. Virksomh., June 1935-May 1936, p. 39;
N. Bohr, Nature **138**(1936)695.

[18] See for example J. M. Blatt and V. Weisskopf, *Theoretical Nuclear Physics*, Wiley, New York, NY, 1952, Chap. 14.

[19] H. Yukawa, Proc. Phys. Math. Soc. (Japan) **17**(1935)48.

[20] C.M.C. Lattes, H. Muirhead, G.P.S. Occhialini, and C.F. Powell, Nature, **159**(1947)694.

[21] M. Gell-Mann, Phys. Lett. **8**(1964)214.

[22] G. Zweig, CERN Report No. 81821 TH401 (1964), unpublished.

[23] W. Pauli, Z. Phys. **31**(1925)765.

[24] O. W. Greenberg, Phys. Rev. Lett. **13**(1964)598;
M. Y. Han and Y. Nambu, Phys. Rev. **B139**(1965)1006.

[25] M. Gell-Mann, "The Symmetry Group of Vector and Axial Currents", *Physics 1* (1964), pp.63-75.

[26] E. D. Bloom, et al., Phys. Rev. Lett **23**(1969)930.

[27] R. P. Feynman, Phys. Rev. Lett. **23**(1969)1415.

[28] C. G. Callan and D. J. Gross, Phys. Rev. Lett. **21**(1968)311;
J. D. Bjorken, Phys. Rev. **179**(1969)1547.

[29] K. G. Wilson, Phys. Rev. **179**(1969)1499.

[30] C. G. Callan and D. J. Gross, Phys. Rev. **D8**(1973)4383.

[31] S. Coleman and D. J. Gross, Phys. Rev. Lett. **31**(1973)851.

[32] D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30**(1973)1343.

[33] H. D. Politzer, Phys. Rev. Lett. **30**(1973)3633.

[34] X. Ji, Phys. Rev. Lett. **78**(1997)610.

[35] X. Ji and J. Osborne, Phys. Rev. **D58**(1998)094018.

[36] A. Ali, V. M. Braun, and G. Hiller, Phys. Lett. **B266**(1991)117.

[37] X. Ji and J. Osborne, *Eur. Phys. J.* **C9**(1999)487.

[38] I have found
M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley, Reading, MA, 1995
extremely helpful for a general discussion. More advanced topics can be found in
R. J. Rivers, *Path Integral Methods in Quantum Field Theory*, Cambridge University, Cambridge, 1987.

[39] The original papers on the 'proper' prescription are

S. Mandelstam, Nucl. Phys. **B213**(1983)149;

G. Leibbrandt, Phys. Rev. **D29**(1984)1699.

A standard reference on this material is

A. Bassetto, G. Nardelli, and R. Soldati, *Yang-Mills Theories in Algebraic Noncovariant Gauges : Quantization and Renormalization*, World Scientific, Singapore (1991).

[40] J. Collins, *Renormalization*, Cambridge University, Cambridge, 1984.

[41] G. 't Hooft, Nucl. Phys. **B33**(1971)173;

B. W. Lee and J. Zinn-Justin, Phys. Rev. **D5**(1972)3121,3137;

A. A. Slavnov, Theor. Math. Phys. **10**(1972)99;

J. C. Taylor, Nucl. Phys. **B33**(1971)436;

C. Becchi, A. Rouet and R. Stora, Ann. Phys. **98**(1976)287;

M. Z. Iofa and I. V. Tyutin, Theor. Math. Phys. **27**(1976)316.

[42] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley, Reading, MA, 1995, Chapter 12.

[43] See for example

R. K. Ellis, W. J. Stirling and B. R. Webber, *QCD and Collider Physics*, Cambridge Univeristy Press, Cambridge, UK, 1996.

[44] A. Bassetto, M. Dalbosco, I. Lazzizzera and R. Soldati, Phys. Rev. **D36**(1987)3138;

S. Mandelstam, Nucl. Phys. **213**(1983)149;

G. Leibbrandt, Phys. Rev. **D29**(1984)1699.

[45] G. Curci, W. Furmanski and R. Petronzio, Nucl. Phys. **B175**(1980)27.

[46] W. Furmanski and R. Petronzio, Phys. Lett. **B97**(1980)437;

R. K. Ellis and W. Vogelsang, hep/ph-9602356;

W. Vogelsang, Nucl. Phys. **B475**(1996)47.

[47] G. Sterman, Phys. Rev. **D17**(1978)1789, 2773.

[48] S. Libby and G. Sterman, Phys. Rev. **D18**(1978)3252, 4737.

[49] J. C. Collins and G. Sterman, Nucl. Phys. **B185**(1981)172.

[50] L.D. Landau, Nucl. Phys. **13**(1959)181.

[51] H. Lehmann, K. Symanzik, and W. Zimmermann, Nuovo Cim. **1**(1955)205.

[52] J. C. Collins and D. E. Soper, Nucl. Phys. **194B**(1982)445.

[53] J. S. Bell and R. Jackiw, Nuovo Cimento **60A**(1969)47;
S. L. Adler, Phys. Rev. **117**(1969)2426.

[54] V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**(1972)438;
L. N. Lipatov, Sov. J. Nucl. Phys. **20**(1975)95;
G. Altarelli and G. Parisi, Nucl. Phys. **B126**(1977)298;
Yu. L. Dokshitzer, Sov. Phys. JETP **46**(1977)641.

[55] J. C. Collins, D. E. Soper and G. Sterman, in *Perturbative QCD*, ed. A. H. Mueller, World Scientific, Singapore, 1989.

[56] J. C. Collins, L. Frankfurt, and M. Strikman, Phys. Rev. **D56**(1997)2982.

[57] R. K. Ellis, W. Furmanski, R. Petronzio, Nucl. Phys. **B207**(1982)1; **B212**(1983)29;
J. Qiu, Phys. Rev. **D42**(1990)30.

[58] S. Coleman and R.E. Norton, Nuovo Cim. **28**(1965)438.

[59] T. Kinoshita, J. Math. Phys. **3**(1962)650;
T. D. Lee and M. Nauenberg, Phys. Rev. **B133**(1964)1549.

[60] G. Sterman, *An Introduction Quantum Field Theory,* Cambridge University Press, Cambridge, 1993.

[61] W. A. Bardeen, A. J. Buras, D. W. Duke and T. Muta, Phys. Rev. **D18** (1978)3998.

[62] R. Mertig and W. L. van Neerven, Z. Phys. **C70**(1996)637.

[63] C. Itzykson and J. B. Zuber, *Quantum Field Theory*, McGraw-Hill Inc., New York, NY, 1980.

[64] R. L. Jaffe and X. Ji, Phys. Rev. Lett. **67**(1991)552;
X. Ji, Nucl. Phys. **B402**(1993)217.

[65] D. Gross and F. Wilczek, Phys. Rev. **D9**(1974)980.

[66] X. Ji, J. Tang and P. Hoodbhoy, Phys. Rev. Lett. **76**(1996)740.

[67] A. V. Radyushkin, Phys. Lett. **B380**(1996)417; Phys. Lett. **B385**(1996)333; Phys. Rev. **D56**(1997)5524.

[68] X. Ji, Phys. Rev. **D55**(1997)7114.

[69] J. Bartels and M. Loewe, Z. Phys. **C12**(1982)263;
B. Geyer et al., Z. Phys. **C26**(1985)591;
T. Braunschweig et al., Z. Phys. **C33**(1986)275;
F.-M. Dittes et al., Phys. Lett. **B209**(1988)325;
I. I. Balitsky and V. M. Braun, Nucl. Phys. **B311**(1989)541;
P. Jain and J. P. Ralston, in *Future Directions in Particle and Nuclear Physics at Multi-GeV Hadron Beam Facilities,* BNL, March, 1993;
I. I. Balitsky and A. V. Radyushkin, Phys. Lett. **B413**(1997)114;
J. Blümlein, B. Geyer, and D. Rabaschik, Phys. Lett. **B406**(1997)161;
X. Ji, W. Melnitchouk, X. Song, Phys. Rev. **D56**(1997)5511;
L. L. Frankfurt, A. Freund, V. Guzey, and M. Strikman, hep-ph/9703449;
V. Yu. Petrov et al., hep-ph/9710270;
A. V. Belitsky, B. Geyer, D. Müller and A. Schäfer, hep-ph/9710427;
L. Mankiewicz, G. Piller, E. Stein, M. Vanttinen, T. Weigl, hep-ph/9712251.

[70] Z. Chen, Columbia preprint CU-TP-835, hep-ph/9705279.

[71] K. Watanabe, Prog. Th. Phys. **67**(1982)1834.

[72] D. Müller, D. Robaschik, B. Geyer, F.M.Dittes, and J. Horejsi, Fortschr. Phys. **42**(1994)101.

[73] D. Müller, Phys. Rev. **D58**(1998)054005;
A. V. Belitsky and D. Müller, Phys. Lett. **B417**(1998)129.

[74] L. Mankiewicz, G. Piller, E. Stein, M. Vanttinen, T. Weigl, TUM-T39-97-31, e-Print Archive: hep-ph/9712251.

[75] X. Ji and J. Osborne, Phys. Rev. **D57**(1998)1337.

[76] P. Hoodbhoy and X. Ji, Phys. Rev. **D58**(1998)054006.

[77] G. Altarelli, R. D. Ball, S. Forte and G. Ridolfi, Nucl. Phys. **B496**(1997)337.

[78] X. Ji, J. Phys. **G24**(1998)1.

[79] S. A. Larin, Phys. Lett. **B303**(1993)113.

[80] The original references are
P. Carruthers, Phys. Rev. **D2**(1970)2265;
R. J. Crewther, Phys. Rev. **D3**(1971)3152.
For a general discussion, see
M. E. Peskin and D. V. Schroder, *An Introduction to Quantum Field Theory*, Addison-Wesley, Reading, MA, 1995, Chapter 19.

[81] M. Stratmann, Z. Phys. **C60**(1993)763; K. Abe et al., Phys. Rev. **D58**(1998)112003.

[82] I. I. Balitsky, V. M. Braun, Y. Koike, and T. Tanaka, Phys. Rev. Lett. **77**(1996)3078; A. V. Belitsky, Phys. Lett. **B405**(1997)312.

[83] J. Kogut and D. Soper, Phys. Rev. **D1**(1970)2901;
E. Tomboulis, Phys. Rev. **D8**(1973)2736.

[84] E. V. Shuryak and A. I. Vainshtein, Nucl. Phys. **B199**(1982)951.

[85] A. P. Bukhvostov, E. A. Kuraev, and L. N. Lipatov, JETP Lett. **37**(1983)482;
A. P. Bukhvostov, E. A. Kuraev, and L. N. Lipatov, Sov. Phys. JETP **60**(1984)22;
P. G. Ratcliffe, Nucl. Phys. **B264**(1986)493;
I. I. Balitsky and V. M. Braun, Nucl. Phys. **B311**(1988/89)541;
X. Ji and C. Chou, Phys. Rev. **D42**(1990)3637;
J. Kodaira et al., Phys. Lett. **B387**(1996)855.

[86] R. L. Jaffe and X. Ji, Phys. Rev. Lett. **67**(1991)552;
R. L. Jaffe and X. Ji, Phys. Rev. Lett., Nucl. Phys. **B375**(1992)527.

[87] X. Ji, Nucl. Phys. **B402**(1993)217.

[88] S. Weinberg, The Theory of Quantum Fields Vol. I, Cambridge University, Cambridge, 1995.

[89] E. P. Wigner, Ann. Math. **40**(1939)149.

[90] M. Hamermesh, *Group Theory and Its Application to Physical Problems*, Addison-Wesley Publishing Co., Reading, MA, 1962.

[91] H. M. Chan and S. T. Tsou, *Some Elementary Gauge Theory Concepts*, World Scientific, Singapore (1993).

[92] G. 't Hooft, Nucl. Phys. **B72**(1974)461;
E. Witten, Nucl. Phys. **B160**(1979)57;
A. V. Manohar, hep-ph/9802419.

[93] R. Dashen and A.V. Manohar, Phys. Lett. **B315**(1993)425; **B315**(1993)438.

[94] T. D. Cohen and C. K. Chow, hep-ph/0003131.

[95] O. Klein, Z. Phys. **37**(1926)895;
W. Gordon, Z. Phys. **40**(1926)117.

[96] P. A. M. Dirac, Proc. Roy. Socy. **133**(1931)60.

[97] C. D. Anderson, Science **76**(1932)238; Phys. Rev. **43**(1933)491.

[98] K. Fujikawa, Phys. Rev. Lett. **42**(1979)1195;
K. Fujikawa, Phys. Rev. **D21**(1980)2848.

[99] P. Breitenlohner and D. Maison, Commun. Math. Phys. **52**(1977)11.

[100] S. Okubo, Prog. Theor. Phys. Supp. **86**(1986)287.

[101] W. A. Bardeen, R. Gastmans and B. Lautrup, Nucl. Phys. **B46**(1972)319.

[102] J. G. Körner, D. Kreimer and K. Schilcher, Z. Phys. **C54**(1992)503.

[103] B. A. Ovrut, Nucl. Phys. **B213**(1983)241. R. Ferrari, A LeYaouanc, L. Oliver and J. C. Raynal, Phys. Rev. **D52**(1995)3036.